# Student Solutions Manual

# PROBABILITY AND STATISTICS

FOR ENGINEERING AND THE SCIENCE

NINTH EDITION

JAY L. DEVORE

# MATT CARLTON

www.konkur.in

# **Student Solutions Manual**







California Polytechnic State University

Prepared by

# **Matt Carlton**

California Polytechnic State University



Australia • Brazil • Mexico • Singapore • United Kingdom • United States

Telegram: @uni\_k

No. In a subscription of party

Printed at the United States o



#### © 2016 Cengage Learning

#### WCN: 01-100-101

ALL RIGHTS RESERVED. No part of this work covered by the copyright herein may be reproduced, transmitted, stored, or used in any form or by any means graphic, electronic, or mechanical, including but not limited to photocopying, recording, scanning, digitizing, taping, Web distribution, information networks, or information storage and retrieval systems, except as permitted under Section 107 or 108 of the 1976 United States Copyright Act, without the prior written permission of the publisher.

For product information and technology assistance, contact us at Cengage Learning Customer & Sales Support, 1-800-354-9706.

For permission to use material from this text or product, submit all requests online at www.cengage.com/permissions Further permissions questions can be emailed to permissionrequest@cengage.com.

#### ISBN: 978-1-305-26059-7

Cengage Learning 20 Channel Center Street Boston, MA 02210 USA

Cengage Learning is a leading provider of customized learning solutions with office locations around the globe, including Singapore, the United Kingdom, Australia, Mexico, Brazil, and Japan. Locate your local office at: www.cengage.com/global.

Cengage Learning products are represented in Canada by Nelson Education, Ltd.

To learn more about Cengage Learning Solutions, visit www.cengage.com.

Purchase any of our products at your local college store or at our preferred online store www.cengagebrain.com.

Printed in the United States of America Print Number: 01 Print Year: 2014 www.konkur.in

# CONTENTS

Chapter 1	Overview and Descriptive Statistics	1
Chapter 2	Probability	22
Chapter 3	Discrete Random Variables and Probability Distributions	41
Chapter 4	Continuous Random Variables and Probability Distributions	57
Chapter 5	Joint Probability Distributions and Random Samples	79
Chapter 6	Point Estimation	94
Chapter 7	Statistical Intervals Based on a Single Sample	100
Chapter 8	Tests of Hypotheses Based on a Single Sample	108
Chapter 9	Inferences Based on Two Samples	119
Chapter 10	The Analysis of Variance	135
Chapter 11	Multifactor Analysis of Variance	142
Chapter 12	Simple Linear Regression and Correlation	157
Chapter 13	Nonlinear and Multiple Regression	174
Chapter 14	Goodness-of-Fit Tests and Categorical Data Analysis	196
Chapter 15	Distribution-Free Procedures	205
Chapter 16	Quality Control Methods	209



#### Compared by branching the state of the state

CHAPTER 1

## Section 1.1

- a. Los Angeles Times, Oberlin Tribune, Gainesville Sun, Washington Post
- b. Duke Energy, Clorox, Seagate, Neiman Marcus
- c. Vince Correa, Catherine Miller, Michael Cutler, Ken Lee
- d. 2.97, 3.56, 2.20, 2.97

3.

1.

- a. How likely is it that more than half of the sampled computers will need or have needed warranty service? What is the expected number among the 100 that need warranty service? How likely is it that the number needing warranty service will exceed the expected number by more than 10?
- b. Suppose that 15 of the 100 sampled needed warranty service. How confident can we be that the proportion of *all* such computers needing warranty service is between .08 and .22? Does the sample provide compelling evidence for concluding that more than 10% of all such computers need warranty service?

#### 5.

- a. No. All students taking a large statistics course who participate in an SI program of this sort.
- b. The advantage to randomly allocating students to the two groups is that the two groups should then be fairly comparable before the study. If the two groups perform differently in the class, we might attribute this to the treatments (SI and control). If it were left to students to choose, stronger or more dedicated students might gravitate toward SI, confounding the results.
- c. If all students were put in the treatment group, there would be no firm basis for assessing the effectiveness of SI (nothing to which the SI scores could reasonably be compared).
- 7. One could generate a simple random sample of all single-family homes in the city, or a stratified random sample by taking a simple random sample from each of the 10 district neighborhoods. From each of the selected homes, values of all desired variables would be determined. This would be an enumerative study because there exists a finite, identifiable population of objects from which to sample.

1

9.

a. There could be several explanations for the variability of the measurements. Among them could be measurement error (due to mechanical or technical changes across measurements), recording error, differences in weather conditions at time of measurements, etc.

b. No, because there is no sampling frame.

## Section 1.2

11.

3L	1	
3H	56678	
4L	000112222234	
4H	5667888	stem: tenths
5L	144	leaf: hundredths
SH	58	
6L	2	
5H	6678	
7L	nue surve velen	
7H	5	
	<ul> <li>A second sec second second sec</li></ul>	

The stem-and-leaf display shows that .45 is a good representative value for the data. In addition, the display is not symmetric and appears to be positively skewed. The range of the data is .75 - .31 = .44, which is comparable to the typical value of .45. This constitutes a reasonably large amount of variation in the data. The data value .75 is a possible outlier.

13.

a.

	to monore the second many shows offer	
12	2	stem: tens
12	445	leaf: ones
12	6667777	
12	889999	
13	00011111111	
13	222222222333333333333333333	
13	444444444444444445555555555555555555555	555555555
13	6666666666667777777777	
13	888888888888999999	
14	0000001111	
14	2333333	
14	444	
14	77 The second second second second	

The observations are highly concentrated at around 134 or 135, where the display suggests the typical value falls.



The histogram of ultimate strengths is symmetric and unimodal, with the point of symmetry at approximately 135 ksi. There is a moderate amount of variation, and there are no gaps or outliers in the distribution.

French

00234566

1

2356

1369

2 16

223558

	American	
- History	dad Darres have	8
	755543211000	9
	9432	10
	6630	11
	850	12
	8	13

American movie times are unimodal strongly positively skewed, while French movie times appear to be bimodal. A typical American movie runs about 95 minutes, while French movies are typically either around 95 minutes or around 125 minutes. American movies are generally shorter than French movies and are less variable in length. Finally, both American and French movies occasionally run very long (outliers at 162 minutes and 158 minutes, respectively, in the samples).

Telegram: @uni\_k

- 17. The sample size for this data set is n = 7 + 20 + 26 + ... + 3 + 2 = 108.
  - a. "At most five bidders" means 2, 3, 4, or 5 bidders. The proportion of contracts that involved at most 5 bidders is (7 + 20 + 26 + 16)/108 = 69/108 = .639. Similarly, the proportion of contracts that involved at least 5 bidders (5 through 11) is equal to (16 + 11 + 9 + 6 + 8 + 3 + 2)/108 = .509.
  - **b.** The number of contracts with between 5 and 10 bidders, inclusive, is 16 + 11 + 9 + 6 + 8 + 3 = 53, so the proportion is 53/108 = .491. "Strictly" between 5 and 10 means 6, 7, 8, or 9 bidders, for a proportion equal to (11 + 9 + 6 + 8)/108 = .34/108 = .315.
  - c. The distribution of number of bidders is positively skewed, ranging from 2 to 11 bidders, with a typical value of around 4-5 bidders.



19.

- a. From this frequency distribution, the proportion of wafers that contained at least one particle is (100-1)/100 = .99, or 99%. Note that it is much easier to subtract 1 (which is the number of wafers that contain 0 particles) from 100 than it would be to add all the frequencies for 1, 2, 3,... particles. In a similar fashion, the proportion containing at least 5 particles is (100 1-2-3-12-11)/100 = .71, or, 71%.
- **b.** The proportion containing between 5 and 10 particles is (15+18+10+12+4+5)/100 = 64/100 = .64, or 64%. The proportion that contain strictly between 5 and 10 (meaning strictly *more* than 5 and strictly *less* than 10) is (18+10+12+4)/100 = 44/100 = .44, or 44%.
- c. The following histogram was constructed using Minitab. The histogram is *almost* symmetric and unimodal; however, the distribution has a few smaller modes and has a very slight positive skew.



#### 21.

**a.** A histogram of the y data appears below. From this histogram, the number of subdivisions having no cul-de-sacs (i.e., y = 0) is 17/47 = .362, or 36.2%. The proportion having at least one cul-de-sac ( $y \ge 1$ ) is (47 - 17)/47 = 30/47 = .638, or 63.8%. Note that subtracting the number of cul-de-sacs with y = 0 from the total, 47, is an easy way to find the number of subdivisions with  $y \ge 1$ .



**b.** A histogram of the z data appears below. From this histogram, the number of subdivisions with at most 5 intersections (i.e.,  $z \le 5$ ) is 42/47 = .894, or 89.4%. The proportion having fewer than 5 intersections (i.e.,  $z \le 5$ ) is 39/47 = .830, or 83.0%.





The distribution of tantrum durations is unimodal and heavily positively skewed. Most tantrums last between 0 and 11 minutes, but a few last more than half an hour! With such heavy skewness, it's difficult to give a representative value.

25. The transformation creates a much more symmetric, mound-shaped histogram.



Histogram of original data:

23. Note: since the class intervals have unequal length, we must use a *density scale*.



27.

a. The endpoints of the class intervals overlap. For example, the value 50 falls in both of the intervals 0-50 and 50-100.

**b.** The lifetime distribution is positively skewed. A representative value is around 100. There is a great deal of variability in lifetimes and several possible candidates for outliers.

Class Interval	Frequency	<b>Relative Frequency</b>	
0-< 50	9	0.18	
50-<100	19	0.38	
100-<150	11	0.22	
150-<200	4	0.08	
200-<250	2	0.04	
250-<300	2	0.04	
300-<350	1	0.02	
350-<400	1	0.02	
400-<450	0	0.00	
450-<500	0	0.00	
500-<550	1	0.02	
	50	1.00	

10. A Manual and the second provides the straight plat and the second s second se second s

www.konkur.in





c. There is much more symmetry in the distribution of the transformed values than in the values themselves, and less variability. There are no longer gaps or obvious outliers.

<b>Class Interval</b>	Frequency	Relative Frequency	
2.25-<2.75	2	0.04	
2.75-<3.25	2	0.04	
3.25-<3.75	3	0.06	
3.75-<4.25	8	0.16	
4.25-<4.75	18	0.36	
4.75-<5.25	10	0.20	
5.25-<5.75	4	0.08	
5.75-<6.25	3	0.06	



**d.** The proportion of lifetime observations in this sample that are less than 100 is .18 + .38 = .56, and the proportion that is at least 200 is .04 + .04 + .02 + .02 = .14.

29.

# Chapter 1: Overview and Descriptive Statistics

	Physical Activity	Frequency	l Fi	Relative			
	A	28	of Jungers	.28	No.		
	B	19		.19			
	C	18		.18			
	D	17		.17			
	Е	9		.09			
	F	9		.09			
		100	in the loss	1.00	the month of		
		5.6.102		in the second	i how to egod	in removing the	
30	0-1				182	The second	
						t search sectorities with	
2:	STUSIO STORAGE						
						WE TE LIVE TO WAT	
20	-		Sto mes			2) 153=16585=	
jount 15	5-		15.51	28. Lud			
-				Just 2 Lit		t a atom nights :	
10	)-	2000 000	and the	14 210 21	in statt	and the man into here	
						statute of a cales	
-	5-	The Sulton	1.5.14		10-10	The month of the	
10	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1		Arost		a derive alle	anner the Jorn on	
	A	В ( Туре	of Physical .	D E Activity	i add to lo Fues	te anna i mharr baile i	
			1.211	Cristo I -	CALD shows	here two transitions	
	Class	Fre	quency	Cum. Fre	eq. Cur	n. Rel. Freq.	
36. 21	0.0-<4.	.0	2	2	el altradio	0.050	
	4.0-<8.	.0	14	16		0.400	
	8.0-<12	2.0	11	27		0.675	
	120 <1/	60	8	35		0.875	

31.

16.0-<20.0

20.0-<24.0

24.0-<28.0

0.975

0.975

1.000

C. J. 204 carries decreased used is reaches (2011) (i.e. by 1.204 - 1.011 - 1.201), its farm of the 2 middle values in this sections of (b more area). My, the reaction of I charm.

39

39

40

4

0

1

9

#### Section 1.3

#### 33.

- a. Using software,  $\bar{x} = 640.5$  (\$640,500) and  $\tilde{x} = 582.5$  (\$582,500). The average sale price for a home in this sample was \$640,500. Half the sales were for less than \$582,500, while half were for more than \$582,500.
- b. Changing that one value lowers the sample mean to 610.5 (\$610,500) but has no effect on the sample median.
- c. After removing the two largest and two smallest values,  $\overline{x}_{ir(20)} = 591.2$  (\$591,200).
- d. A 10% trimmed mean from removing just the highest and lowest values is  $\bar{x}_{\nu(10)} = 596.3$ . To form a 15% trimmed mean, take the average of the 10% and 20% trimmed means to get  $\bar{x}_{\nu(15)} = (591.2 + 596.3)/2 = 593.75$  (\$593,750).

#### 35. The sample size is n = 15.

- a. The sample mean is  $\bar{x} = 18.55/15 = 1.237 \ \mu\text{g/g}$  and the sample median is  $\tilde{x} = \text{the 8}^{\text{th}}$  ordered value = .56  $\mu$ g/g. These values are very different due to the heavy positive skewness in the data.
- b. A 1/15 trimmed mean is obtained by removing the largest and smallest values and averaging the remaining 13 numbers: (.22 + ... + 3.07)/13 = 1.162. Similarly, a 2/15 trimmed mean is the average of the middle 11 values: (.25 + ... + 2.25)/11 = 1.074. Since the average of 1/15 and 2/15 is .1 (10%), a 10% trimmed mean is given by the midpoint of these two trimmed means: (1.162 + 1.074)/2 = 1.118 μg/g.
- c. The median of the data set will remain .56 so long as that's the 8<sup>th</sup> ordered observation. Hence, the value .20 could be increased to as high as .56 without changing the fact that the 8<sup>th</sup> ordered observation is .56. Equivalently, .20 could be increased by as much as .36 without affecting the value of the sample median.
- 37.  $\overline{x} = 12.01$ ,  $\overline{x} = 11.35$ ,  $\overline{x}_{tr(10)} = 11.46$ . The median or the trimmed mean would be better choices than the mean because of the outlier 21.9.
- 39.

**a.** 
$$\Sigma x_i = 16.475$$
 so  $\overline{x} = \frac{16.475}{16} = 1.0297$ ;  $\widetilde{x} = \frac{(1.007 + 1.011)}{2} = 1.009$ 

**b.** 1.394 can be decreased until it reaches 1.011 (i.e. by 1.394 - 1.011 = 0.383), the largest of the 2 middle values. If it is decreased by more than 0.383, the median will change.

x/n = 7/10 = .7

- 41.
- **b.**  $\overline{x} = .70 =$  the sample proportion of successes
- c. To have x/n equal .80 requires x/25 = .80 or x = (.80)(25) = 20. There are 7 successes (S) already, so another 20 7 = 13 would be required.
- 43. The median and certain trimmed means can be calculated, while the mean cannot the exact values of the "100+" observations are required to calculate the mean.  $\tilde{x} = \frac{(57+79)}{2} = 68.0$ ,

 $\overline{x}_{tr(20)} = 66.2, \ \overline{x}_{tr(30)} = 67.5.$ 

#### Section 1.4

2.

- 45.
- **a.**  $\overline{x} = 115.58$ . The deviations from the mean are 116.4 115.58 = .82, 115.9 115.58 = .32, 114.6 115.58 = -.98, 115.2 115.58 = -.38, and 115.8 115.58 = .22. Notice that the deviations from the mean sum to zero, as they should.

**b.** 
$$s^2 = [(.82)^2 + (.32)^2 + (-.98)^2 + (-.38)^2 + (.22)^2]/(5-1) = 1.928/4 = .482$$
, so  $s = .694$ .

- c.  $\Sigma x_i^2 = 66795.61$ , so  $s^2 = S_{xx}/(n-1) = (\Sigma x_i^2 (\Sigma x_i)^2 / n)/(n-1) = (66795.61 (577.9)^2/5)/4 = 1.928/4 = .482.$
- **d.** The new sample values are: 16.4 15.9 14.6 15.2 15.8. While the new mean is 15.58, all the deviations are the same as in part (a), and the variance of the transformed data is identical to that of part (b).

47.

- **a.** From software,  $\tilde{x} = 14.7\%$  and  $\bar{x} = 14.88\%$ . The sample average alcohol content of these 10 wines was 14.88\%. Half the wines have alcohol content below 14.7% and half are above 14.7% alcohol.
- **b.** Working long-hand,  $\Sigma(x_i \overline{x})^2 = (14.8 14.88)^2 + ... + (15.0 14.88)^2 = 7.536$ . The sample variance equals  $s^2 = \Sigma(x_i \overline{x})^2 = 7.536/(10 1) = 0.837$ .
- c. Subtracting 13 from each value will not affect the variance. The 10 new observations are 1.8, 1.5, 3.1, 1.2, 2.9, 0.7, 3.2, 1.6, 0.8, and 2.0. The sum and sum of squares of these 10 new numbers are  $\sum y_i = 18.8$  and  $\sum y_i^2 = 42.88$ . Using the sample variance shortcut, we obtain  $s^2 = [42.88 (18.8)^2/10]/(10 1) = 7.536/9 = 0.837$  again.

49.

a. 
$$\Sigma x_i = 2.75 + \dots + 3.01 = 56.80$$
,  $\Sigma x_i^2 = 2.75^2 + \dots + 3.01^2 = 197.8040$ 

**b.** 
$$s^2 = \frac{197.8040 - (56.80)^2 / 17}{16} = \frac{8.0252}{16} = .5016, \ s = .708$$

51.

3.

From software,  $s^2 = 1264.77 \text{ min}^2$  and s = 35.56 min. Working by hand,  $\Sigma x = 2563$  and  $\Sigma x^2 = 368501$ , so

$$s^2 = \frac{368501 - (2563)^2 / 19}{19 - 1} = 1264.766 \text{ and } s = \sqrt{1264.766} = 35.564$$

**b.** If y = time in hours, then y = cx where  $c = \frac{1}{60}$ . So,  $s_y^2 = c^2 s_x^2 = (\frac{1}{60})^2 1264.766 = .351 \text{ hr}^2$ and  $s_y = cs_x = (\frac{1}{60})35.564 = .593 \text{ hr}.$ 

53.

- a. Using software, for the sample of balanced funds we have  $\overline{x} = 1.121, \overline{x} = 1.050, s = 0.536$ ; for the sample of growth funds we have  $\overline{x} = 1.244, \overline{x} = 1.100, s = 0.448$ .
- b. The distribution of expense ratios for this sample of balanced funds is fairly symmetric, while the distribution for growth funds is positively skewed. These balanced and growth mutual funds have similar median expense ratios (1.05% and 1.10%, respectively), but expense ratios are generally higher for growth funds. The lone exception is a balanced fund with a 2.86% expense ratio. (There is also one unusually low expense ratio in the sample of balanced funds, at 0.09%.)



55.

a.

Lower half of the data set: 325 325 334 339 356 356 359 359 363 364 364 366 369, whose median, and therefore the lower fourth, is 359 (the 7<sup>th</sup> observation in the sorted list).

Upper half of the data set: 370 373 373 374 375 389 392 393 394 397 402 403 424, whose median, and therefore the upper fourth is 392.

So,  $f_s = 392 - 359 = 33$ .

- b. inner fences: 359 1.5(33) = 309.5, 392 + 1.5(33) = 441.5
   To be a mild outlier, an observation must be below 309.5 or above 441.5. There are none in this data set. Clearly, then, there are also no extreme outliers.
  - c. A boxplot of this data appears below. The distribution of escape times is roughly symmetric with no outliers. Notice the box plot "hides" the fact that the distribution contains two gaps, which can be seen in the stem-and-leaf display.



**d.** Not until the value x = 424 is lowered below the upper fourth value of 392 would there be any change in the value of the upper fourth (and, thus, of the fourth spread). That is, the value x = 424 could not be decreased by more than 424 - 392 = 32 seconds.

57.

**a.**  $f_s = 216.8 - 196.0 = 20.8$ 

inner fences: 196 - 1.5(20.8) = 164.6, 216.8 + 1.5(20.8) = 248outer fences: 196 - 3(20.8) = 133.6, 216.8 + 3(20.8) = 279.2Of the observations listed, 125.8 is an extreme low outlier and 250.2 is a mild high outlier.

**b.** A boxplot of this data appears below. There is a bit of positive skew to the data but, except for the two outliers identified in part (a), the variation in the data is relatively small.



59.

www.konkur.in

a. If you aren't using software, don't forget to *sort* the data first! *ED*: median = .4, lower fourth = (.1 + .1)/2 = .1, upper fourth = (2.7 + 2.8)/2 = 2.75, fourth spread = 2.75 - .1 = 2.65

*Non-ED*: median = (1.5 + 1.7)/2 = 1.6, lower fourth = .3, upper fourth = 7.9, fourth spread = 7.9 - .3 = 7.6.

b. ED: mild outliers are less than .1 - 1.5(2.65) = -3.875 or greater than 2.75 + 1.5(2.65) = 6.725. Extreme outliers are less than .1 - 3(2.65) = -7.85 or greater than 2.75 + 3(2.65) = 10.7. So, the two largest observations (11.7, 21.0) are extreme outliers and the next two largest values (8.9, 9.2) are mild outliers. There are no outliers at the lower end of the data.

*Non-ED*: mild outliers are less than .3 - 1.5(7.6) = -11.1 or greater than 7.9 + 1.5(7.6) = 19.3. Note that there are no mild outliers in the data, hence there cannot be any extreme outliers, either.

c. A comparative boxplot appears below. The outliers in the ED data are clearly visible. There is noticeable positive skewness in both samples; the Non-ED sample has more variability then the Ed sample; the typical values of the ED sample tend to be smaller than those for the Non-ED sample.



61. Outliers occur in the 6a.m. data. The distributions at the other times are fairly symmetric. Variability and the "typical" gasoline-vapor coefficient values increase somewhat until 2p.m., then decrease slightly.

#### **Supplementary Exercises**

63. As seen in the histogram below, this noise distribution is bimodal (but close to unimodal) with a positive skew and no outliers. The mean noise level is 64.89 dB and the median noise level is 64.7 dB. The fourth spread of the noise measurements is about 70.4 - 57.8 = 12.6 dB.



65.

a. The histogram appears below. A representative value for this data would be around 90 MPa. The histogram is reasonably symmetric, unimodal, and somewhat bell-shaped with a fair amount of variability ( $s \approx 3$  or 4 MPa).



**b.** The proportion of the observations that are at least 85 is 1 - (6+7)/169 = .9231. The proportion less than 95 is 1 - (13+3)/169 = .9053.

- c. 90 is the midpoint of the class 89 <91, which contains 43 observations (a relative frequency of 43/169 = .2544). Therefore about half of this frequency, .1272, should be added to the relative frequencies for the classes to the left of x = 90. That is, the approximate proportion of observations that are less than 90 is .0355 + .0414 + .1006 + .1775 + .1272 = .4822.
- a. Aortic root diameters for males have mean 3.64 cm, median 3.70 cm, standard deviation 0.269 cm, and fourth spread 0.40. The corresponding values for females are x̄ = 3.28 cm, x̃ = 3.15 cm, s = 0.478 cm, and f<sub>s</sub> = 0.50 cm. Aortic root diameters are typically (though not universally) somewhat smaller for females than for males, and females show more variability. The distribution for males is negatively skewed, while the distribution for females is positively skewed (see graphs below).



**b.** For females (n = 10), the 10% trimmed mean is the average of the middle 8 observations:  $\overline{x}_{tr(10)} = 3.24$  cm. For males (n = 13), the 1/13 trimmed mean is 40.2/11 = 3.6545, and the 2/13 trimmed mean is 32.8/9 = 3.6444. Interpolating, the 10% trimmed mean is  $\overline{x}_{tr(10)} = 0.7(3.6545) + 0.3(3.6444) = 3.65$  cm. (10% is three-tenths of the way from 1/13 to 2/13).

69.

a.

67.

$$\overline{y} = \frac{\sum y_i}{n} = \frac{\sum (ax_i + b)}{n} = \frac{a\sum x_i + \sum b}{n} = \frac{a\sum x_i + nb}{n} = a\overline{x} + b$$
$$s_y^2 = \frac{\sum (y_i - \overline{y})^2}{n-1} = \frac{\sum (ax_i + b - (a\overline{x} + b))^2}{n-1} = \frac{\sum (ax_i - a\overline{x})^2}{n-1}$$
$$= \frac{a^2 \sum (x_i - \overline{x})^2}{n-1} = a^2 s_x^2$$

16

.872°F

$$x = {}^{\circ}C, y = {}^{\circ}F$$
  
$$\overline{y} = \frac{9}{5}(87.3) + 32 = 189.14 {}^{\circ}F$$
  
$$s_{y} = \sqrt{s_{y}^{2}} = \sqrt{\left(\frac{9}{5}\right)^{2}(1.04)^{2}} = \sqrt{3.5044} = 1$$

71.

- a. The mean, median, and trimmed mean are virtually identical, which suggests symmetry. If there are outliers, they are balanced. The range of values is only 25.5, but half of the values are between 132.95 and 138.25.
- **b.** See the comments for (a). In addition, using 1.5(Q3 Q1) as a yardstick, the two largest and three smallest observations are mild outliers.



73. From software,  $\overline{x} = .9255$ , s = .0809;  $\overline{x} = .93$ ,  $f_s = .1$ . The cadence observations are slightly skewed (mean = .9255 strides/sec, median = .93 strides/sec) and show a small amount of variability (standard deviation = .0809, fourth spread = .1). There are no apparent outliers in the data.



75.

a. The median is the same (371) in each plot and all three data sets are very symmetric. In addition, all three have the same minimum value (350) and same maximum value (392). Moreover, all three data sets have the same lower (364) and upper quartiles (378). So, all three boxplots will be *identical*. (Slight differences in the boxplots below are due to the way Minitab software interpolates to calculate the quartiles.)



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

www.konkur.in

**b.** A comparative dotplot is shown below. These graphs show that there are differences in the variability of the three data sets. They also show differences in the way the values are distributed in the three data sets, especially big differences in the presence of gaps and clusters.



c. The boxplot in (a) is not capable of detecting the differences among the data sets. The primary reason is that boxplots give up some detail in describing data because they use only five summary numbers for comparing data sets.

77.

a

leaf = 1.00 44444444577888999 0001111111124455669999 stem = 0.11 2 1234457 3 11355 4 17 5 3 6 7 67 8 1 HI 10.44, 13.41

 0.5

 0.4

 0.4

 0.4

 0.4

 0.5

 0.4

 0.5

 0.5

 0.6

 0.6

 0.6

 0.6

 0.6

 0.6

 0.6

 0.6

 0.6

 0.6

 0.6

 0.6

 0.6

 0.6

 0.7

 0.6

 0.6

 0.7

 0.6

 0.7

 0.6

 0.7

 0.7

 0.7

 0.7

 0.7

 0.7

 0.7

 0.7

 0.7

 0.7

 0.7

 0.7

 0.7

 0.7

 0.7

 0.7

 0.7

 0.7

 0.7

 0.7

 0.7

 0.7

 0.7

 0.7

 0.7

 0.7

 0.7

 0.7

 0.7

 0.7

 0.7

 0.7

 0.7

 0.7
 </

Since the intervals have unequal width, you must use a density scale.

c. Representative depths are quite similar for the three types of soils — between 1.5 and 2. Data from the C and CL soils shows much more variability than for the other two types. The boxplots for the first three types show substantial positive skewness both in the middle 50% and overall. The boxplot for the SYCL soil shows negative skewness in the middle 50% and mild positive skewness overall. Finally, there are multiple outliers for the first three types of soils, including extreme outliers.

$$\sum_{i=1}^{n+1} x_i = \sum_{i=1}^n x_i + x_{n+1} = n\overline{x}_n + x_{n+1}, \text{ so } \overline{x}_{n+1} = \frac{1}{n+1} \sum_{i=1}^{n+1} x_i = \frac{n\overline{x}_n + x_{n+1}}{n+1}.$$

**b.** In the second line below, we artificially add and subtract  $n\overline{x}_n^2$  to create the term needed for the sample variance:

$$ns_{n+1}^{2} = \sum_{i=1}^{n+1} (x_{i} - \overline{x}_{n+1})^{2} = \sum_{i=1}^{n+1} x_{i}^{2} - (n+1)\overline{x}_{n+1}^{2}$$
$$= \sum_{i=1}^{n} x_{i}^{2} + x_{n+1}^{2} - (n+1)\overline{x}_{n+1}^{2} = \left[\sum_{i=1}^{n} x_{i}^{2} - n\overline{x}_{n}^{2}\right] + n\overline{x}_{n}^{2} + x_{n+1}^{2} - (n+1)\overline{x}_{n+1}^{2}$$
$$= (n-1)s_{n}^{2} + \left\{x_{n+1}^{2} + n\overline{x}_{n}^{2} - (n+1)\overline{x}_{n+1}^{2}\right\}$$

Substitute the expression for  $\overline{x}_{n+1}$  from part (a) into the expression in braces, and it

simplifies to  $\frac{n}{n+1}(x_{n+1}-\overline{x}_n)^2$ , as desired.

c. First,  $\overline{x}_{16} = \frac{15(12.58) + 11.8}{16} = \frac{200.5}{16} = 12.53$ . Then, solving (b) for  $s_{n+1}^2$  gives  $s_{n+1}^2 = \frac{n-1}{n} s_n^2 + \frac{1}{n+1} (x_{n+1} - \overline{x}_n)^2 = \frac{14}{15} (.512)^2 + \frac{1}{16} (11.8 - 12.58)^2 = .238$ . Finally, the standard deviation is  $s_{16} = \sqrt{.238} = .532$ .



79.

8

b.

www.konkur.in

81. Assuming that the histogram is unimodal, then there is evidence of positive skewness in the data since the median lies to the left of the mean (for a symmetric distribution, the mean and median would coincide).

For more evidence of skewness, compare the distances of the 5<sup>th</sup> and 95<sup>th</sup> percentiles from the median: median  $-5^{th}$  %ile = 500 -400 = 100, while 95<sup>th</sup> %ile - median = 720 -500 = 220. Thus, the largest 5% of the values (above the 95th percentile) are further from the median than are the lowest 5%. The same skewness is evident when comparing the 10<sup>th</sup> and 90<sup>th</sup> percentiles to the median, or comparing the maximum and minimum to the median.

83.

**a.** When there is perfect symmetry, the smallest observation  $y_1$  and the largest observation  $y_n$  will be equidistant from the median, so  $y_n - \tilde{x} = \tilde{x} - y_1$ . Similarly, the second-smallest and second-largest will be equidistant from the median, so  $y_{n-1} - \tilde{x} = \tilde{x} - y_2$ , and so on. Thus, the first and second numbers in each pair will be equal, so that each point in the plot will fall exactly on the 45° line.

When the data is positively skewed,  $y_n$  will be much further from the median than is  $y_1$ , so  $y_n - \tilde{x}$  will considerably exceed  $\tilde{x} - y_1$  and the point  $(y_n - \tilde{x}, \tilde{x} - y_1)$  will fall considerably below the 45° line, as will the other points in the plot.

**b.** The median of these n = 26 observations is 221.6 (the midpoint of the 13<sup>th</sup> and 14<sup>th</sup> ordered values). The first point in the plot is (2745.6 - 221.6, 221.6 - 4.1) = (2524.0, 217.5). The others are: (1476.2, 213.9), (1434.4, 204.1), (756.4, 190.2), (481.8, 188.9), (267.5, 181.0), (208.4, 129.2), (112.5, 106.3), (81.2, 103.3), (53.1, 102.6), (53.1, 92.0), (33.4, 23.0), and (20.9, 20.9). The first number in each of the first seven pairs greatly exceeds the second number, so each of those points falls well below the 45° line. A substantial positive skew (stretched upper tail) is indicated.



# **CHAPTER 2**

#### Section 2.1

a.

1.

www.konkur.in

S= {1324, 1342, 1423, 1432, 2314, 2341, 2413, 2431, 3124, 3142, 4123, 4132, 3214, 3241, 4213, 4231}.

- c. Event *B* contains the outcomes where 2 is first or second:  $B = \{2314, 2341, 2413, 2431, 3214, 3241, 4213, 4231\}.$
- **d.** The event  $A \cup B$  contains the outcomes in A or B or both:  $A \cup B = \{1324, 1342, 1423, 1432, 2314, 2341, 2413, 2431, 3214, 3241, 4213, 4231\}$ .  $A \cap B = \emptyset$ , since 1 and 2 can't both get into the championship game.  $A' = \mathscr{S} - A = \{2314, 2341, 2413, 2431, 3124, 3142, 4123, 4132, 3214, 3241, 4213, 4231\}$ .
- 3.
- **a.**  $A = \{SSF, SFS, FSS\}.$
- **b.**  $B = \{SSS, SSF, SFS, FSS\}.$
- c. For event C to occur, the system must have component 1 working (S in the first position), then at least one of the other two components must work (at least one S in the second and third positions): C = {SSS, SSF, SFS}.
- **d.**  $C' = \{SFF, FSS, FSF, FFS, FFF\}.$   $A \cup C = \{SSS, SSF, SFS, FSS\}.$   $A \cap C = \{SSF, SFS\}.$   $B \cup C = \{SSS, SSF, SFS, FSS\}.$  Notice that B contains C, so  $B \cup C = B$ .  $B \cap C = \{SSS, SSF, SFS\}.$  Since B contains C,  $B \cap C = C$ .

**b.** Event A contains the outcomes where 1 is first in the list:  $A = \{1324, 1342, 1423, 1432\}.$ 

5. **a.** The  $3^3 = 27$  possible outcomes are numbered below for later reference.

Outcome Number	Outcome	Outcome Number	Outcome
1	111	15	223
2	112	16	231
3	113	17	232
4	121	18	233
5	122	19	311
6	123	20	312
7	131	21	313
8	132	22	321
9	133	23	322
10	211	24	323
11	212	25	331
12	213	26	332
13	221	27	333
14	222		-

b. Outcome numbers 1, 14, 27 above.

c. Outcome numbers 6, 8, 12, 16, 20, 22 above.

d. Outcome numbers 1, 3, 7, 9, 19, 21, 25, 27 above.

#### 7.

a. S= {BBBAAAA, BBABAAA, BBAABAA, BBAAABAA, BBAAABA, BBAAAAB, BABBAAA, BABABAA, BABAABA, BABAAAB, BAABBAA, BAABABA, BAABAAB, BAABAAB, BAAABBA, BAAABAB, BAAAABB, ABBBAAA, ABBABAA, ABBAABA, ABBAAAB, ABBABAA, ABABBAA, ABABABA, ABABAAB, ABAABBA, ABAABAB, ABAAABB, AABBBAA, AABBABA, AABBAAB, AABBAAB, AABABBA, AABABAB, AABAABB, AAABBBAA, AAABBAB, AAABABB, AAAABBB}.

- b. AAAABBB, AAABABB, AAABBAB, AABAABB, AABAABB, AABABAB.
- 9.
- **a.** In the diagram on the left, the shaded area is  $(A \cup B)'$ . On the right, the shaded area is A', the striped area is B', and the intersection  $A' \cap B'$  occurs where there is both shading and stripes. These two diagrams display the same area.





23

**b.** In the diagram below, the shaded area represents  $(A \cap B)'$ . Using the right-hand diagram from (a), the <u>union</u> of A' and B' is represented by the areas that have either shading <u>or</u> stripes (or both). Both of the diagrams display the same area.



### Section 2.2

11.

**a.** .07.

- **b.** .15 + .10 + .05 = .30.
- c. Let A = the selected individual owns shares in a stock fund. Then P(A) = .18 + .25 = .43. The desired probability, that a selected customer does <u>not</u> shares in a stock fund, equals P(A') = 1 P(A) = 1 .43 = .57. This could also be calculated by adding the probabilities for all the funds that are not stocks.

13.

- **a.**  $A_1 \cup A_2 =$  "awarded either #1 or #2 (or both)": from the addition rule,  $P(A_1 \cup A_2) = P(A_1) + P(A_2) - P(A_1 \cap A_2) = .22 + .25 - .11 = .36.$
- **b.**  $A'_1 \cap A'_2 =$  "awarded neither #1 or #2": using the hint and part (a),  $P(A'_1 \cap A'_2) = P((A_1 \cup A_2)') = 1 - P(A_1 \cup A_2) = 1 - .36 = .64.$
- c.  $A_1 \cup A_2 \cup A_3 =$  "awarded at least one of these three projects": using the addition rule for 3 events,  $P(A_1 \cup A_2 \cup A_3) = P(A_1) + P(A_2) + P(A_3) - P(A_1 \cap A_2) - P(A_1 \cap A_3) - P(A_2 \cap A_3) + P(A_1 \cap A_2 \cap A_3) =$  .22 + .25 + .28 - .11 - .05 - .07 + .01 = .53.
  - **d.**  $A'_1 \cap A'_2 \cap A'_3 =$  "awarded none of the three projects":  $P(A'_1 \cap A'_2 \cap A'_3) = 1 - P(\text{awarded at least one}) = 1 - .53 = .47.$

www.konkur.in

e.  $A'_1 \cap A'_2 \cap A_3 =$  "awarded #3 but neither #1 nor #2": from a Venn diagram,  $P(A'_1 \cap A'_2 \cap A_3) = P(A_3) - P(A_1 \cap A_3) - P(A_2 \cap A_3) + P(A_1 \cap A_2 \cap A_3) =$ .28 - .05 - .07 + .01 = .17. The last term addresses the "double counting" of the two subtractions.



f.  $(A'_1 \cap A'_2) \cup A_3 =$  "awarded neither of #1 and #2, or awarded #3": from a Venn diagram,  $P((A'_1 \cap A'_2) \cup A_3) = P(\text{none awarded}) + P(A_3) = .47 \text{ (from d)} + .28 = 75.$ 



Alternatively, answers to a-f can be obtained from probabilities on the accompanying Venn diagram:



- a. Let *E* be the event that at most one purchases an electric dryer. Then *E'* is the event that at least two purchase electric dryers, and P(E') = 1 P(E) = 1 .428 = .572.
- b. Let A be the event that all five purchase gas, and let B be the event that all five purchase electric. All other possible outcomes are those in which at least one of each type of clothes dryer is purchased. Thus, the desired probability is 1 [P(A) P(B)] = 1 [.116 + .005] = .879.

17.

15.

www.konkur.in

- a. The probabilities do not add to 1 because there are other software packages besides SPSS and SAS for which requests could be made.
- **b.** P(A') = 1 P(A) = 1 .30 = .70.
- c. Since A and B are mutually exclusive events,  $P(A \cup B) = P(A) + P(B) = .30 + .50 = .80$ .
- **d.** By deMorgan's law,  $P(A' \cap B') = P((A \cup B)') = 1 P(A \cup B) = 1 .80 = .20$ . In this example, deMorgan's law says the event "neither A nor B" is the complement of the event "either A or B." (That's true regardless of whether they're mutually exclusive.)
- 19. Let A be that the selected joint was found defective by inspector A, so  $P(A) = \frac{724}{10,000}$ . Let B be analogous for inspector B, so  $P(B) = \frac{751}{10,000}$ . The event "at least one of the inspectors judged a joint to be defective is  $A \cup B$ , so  $P(A \cup B) = \frac{1159}{10,000}$ .
  - **a.** By deMorgan's law,  $P(\text{neither } A \text{ nor } B) = P(A' \cap B') = 1 P(A \cup B) = 1 \frac{1159}{10,000} = \frac{8841}{10,000} = .8841.$
  - **b.** The desired event is  $B \cap A'$ . From a Venn diagram, we see that  $P(B \cap A') = P(B) P(A \cap B)$ . From the addition rule,  $P(A \cup B) = P(A) + P(B) P(A \cap B)$  gives  $P(A \cap B) = .0724 + .0751 .1159 = .0316$ . Finally,  $P(B \cap A') = P(B) P(A \cap B) = .0751 .0316 = .0435$ .
- 21. In what follows, the first letter refers to the auto deductible and the second letter refers to the homeowner's deductible.
  - **a.** P(MH) = .10.
  - **b.**  $P(\text{low auto deductible}) = P(\{LN, LL, LM, LH\}) = .04 + .06 + .05 + .03 = .18$ . Following a similar pattern, P(low homeowner's deductible) = .06 + .10 + .03 = .19.
  - c.  $P(\text{same deductible for both}) = P(\{LL, MM, HH\}) = .06 + .20 + .15 = .41.$
  - **d.** P(deductibles are different) = 1 P(same deductible for both) = 1 .41 = .59.
  - e.  $P(\text{at least one low deductible}) = P(\{LN, LL, LM, LH, ML, HL\}) = .04 + .06 + .05 + .03 + .10 + .03 = .31.$
  - **f.** P(neither deductible is low) = 1 P(at least one low deductible) = 1 .31 = .69.

- 23. Assume that the computers are numbered 1-6 as described and that computers 1 and 2 are the two laptops. There are 15 possible outcomes: (1,2) (1,3) (1,4) (1,5) (1,6) (2,3) (2,4) (2,5) (2,6) (3,4) (3,5) (3,6) (4,5) (4,6) and (5,6).
  - **a.**  $P(\text{both are laptops}) = P(\{(1,2)\}) = \frac{1}{15} = .067.$
  - **b.**  $P(\text{both are desktops}) = P(\{(3,4), (3,5), (3,6), (4,5), (4,6), (5,6)\}) = \frac{6}{15} = .40.$
  - c. P(at least one desktop) = 1 P(no desktops) = 1 P(both are laptops) = 1 .067 = .933.
  - **d.** P(at least one of each type) = 1 P(both are the same) = 1 [P(both are laptops) + P(both are desktops)] = 1 [.067 + .40] = .533.
- 25. By rearranging the addition rule,  $P(A \cap B) = P(A) + P(B) P(A \cup B) = .40 + .55 .63 = .32$ . By the same method,  $P(A \cap C) = .40 + .70 .77 = .33$  and  $P(B \cap C) = .55 + .70 .80 = .45$ . Finally, rearranging the addition rule for 3 events gives

 $P(A \cap B \cap C) = P(A \cup B \cup C) - P(A) - P(B) - P(C) + P(A \cap B) + P(A \cap C) + P(B \cap C) = .85 - .40 - .55 - .70 + .32 + .33 + .45 = .30.$ 

These probabilities are reflected in the Venn diagram below.



- **a.**  $P(A \cup B \cup C) = .85$ , as given.
- **b.**  $P(\text{none selected}) = 1 P(\text{at least one selected}) = 1 P(A \cup B \cup C) = 1 .85 = .15.$
- c. From the Venn diagram, P(only automatic transmission selected) = .22.
- **d.** From the Venn diagram, P(exactly one of the three) = .05 + .08 + .22 = .35.
- 27. There are 10 equally likely outcomes: {A, B} {A, Co} {A, Cr} {A,F} {B, Co} {B, Cr} {B, F} {Co, Cr} {Co, F} and {Cr, F}.
  a. P({A, B}) = 1/10 = .1.
  - **b.**  $P(\text{at least one } C) = P(\{A, Co\} \text{ or } \{A, Cr\} \text{ or } \{B, Co\} \text{ or } \{B, Cr\} \text{ or } \{Co, Cr\} \text{ or } \{Co, F\} \text{ or } \{Cr, F\}) = \frac{7}{10} = .7.$
  - c. Replacing each person with his/her years of experience,  $P(\text{at least 15 years}) = P(\{3, 14\} \text{ or } \{6, 10\} \text{ or } \{6, 14\} \text{ or } \{7, 10\} \text{ or } \{7, 14\} \text{ or } \{10, 14\}) = \frac{6}{10} = .6.$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

CERMIN



151,200 ways to choose six right-handers for the other positions. The total number of possibilities is = (60)(151,200) = 9,072,000.

35.

Section 2.3

- There are  $\binom{10}{5} = 252$  ways to select 5 workers from the day shift. In other words, of all the ways to a. select 5 workers from among the 24 available, 252 such selections result in 5 day-shift workers. Since the grand total number of possible selections is  $\begin{pmatrix} 24 \\ 5 \end{pmatrix} = 42504$ , the probability of randomly selecting 5 day-shift workers (and, hence, no swing or graveyard workers) is 252/42504 = .00593.
- Similar to **a**, there are  $\begin{pmatrix} 8 \\ 5 \end{pmatrix} = 56$  ways to select 5 swing-shift workers and  $\begin{pmatrix} 6 \\ 5 \end{pmatrix} = 6$  ways to select 5 b.

graveyard-shift workers. So, there are 252 + 56 + 6 = 314 ways to pick 5 workers from the same shift. The probability of this randomly occurring is 314/42504 = .00739.

P(at least two shifts represented) = 1 - P(all from same shift) = 1 - .00739 = .99261.c.

www.konkur.in

d. There are several ways to approach this question. For example, let  $A_1 =$  "day shift is unrepresented,"  $A_2 =$  "swing shift is unrepresented," and  $A_3 =$  "graveyard shift is unrepresented." Then we want  $P(A_1 \cup A_2 \cup A_3)$ .

 $N(A_1) = N(\text{day shift unrepresented}) = N(\text{all from swing/graveyard}) = \binom{8+6}{5} = 2002,$ 

since there are 8 + 6 = 14 total employees in the swing and graveyard shifts. Similarly,

 $N(A_2) = {\binom{10+6}{5}} = 4368 \text{ and } N(A_3) = {\binom{10+8}{5}} = 8568. \text{ Next, } N(A_1 \cap A_2) = N(\text{all from graveyard}) = 6$ 

from b. Similarly,  $N(A_1 \cap A_3) = 56$  and  $N(A_2 \cap A_3) = 252$ . Finally,  $N(A_1 \cap A_2 \cap A_3) = 0$ , since at least one shift must be represented. Now, apply the addition rule for 3 events:

$$P(A_1 \cup A_2 \cup A_3) = \frac{2002 + 4368 + 8568 - 6 - 56 - 252 + 0}{42504} = \frac{14624}{42504} = .3441.$$

37.

**a.** By the Fundamental Counting Principle, with  $n_1 = 3$ ,  $n_2 = 4$ , and  $n_3 = 5$ , there are (3)(4)(5) = 60 runs.

- **b.** With  $n_1 = 1$  (just one temperature),  $n_2 = 2$ , and  $n_3 = 5$ , there are (1)(2)(5) = 10 such runs.
- c. For each of the 5 specific catalysts, there are (3)(4) = 12 pairings of temperature and pressure. Imagine we separate the 60 possible runs into those 5 sets of 12. The number of ways to select exactly one run

from each of these 5 sets of 12 is  $\binom{12}{1}^5 = 12^5$ . Since there are  $\binom{60}{5}$  ways to select the 5 runs overall, the desired probability is  $\binom{12}{1}^5 / \binom{60}{5} = 12^5 / \binom{60}{5} = .0456$ .

39. In a-c, the size of the sample space is  $N = \begin{pmatrix} 5+6+4\\ 3 \end{pmatrix} = \begin{pmatrix} 15\\ 3 \end{pmatrix} = 455.$ 

**a.** There are four 23W bulbs available and 5+6 = 11 non-23W bulbs available. The number of ways to select exactly two of the former (and, thus, exactly one of the latter) is  $\binom{4}{2}\binom{11}{1} = 6(11) = 66$ . Hence, the probability is  $\frac{66}{455} = .145$ .

- **b.** The number of ways to select three 13W bulbs is  $\binom{5}{3} = 10$ . Similarly, there are  $\binom{6}{3} = 20$  ways to select three 18W bulbs and  $\binom{4}{3} = 4$  ways to select three 23W bulbs. Put together, there are 10 + 20 + 4 = 34 ways to select three bulbs of the same wattage, and so the probability is 34/455 = .075.
  - c. The number of ways to obtain one of each type is  $\binom{5}{1}\binom{6}{1}\binom{4}{1} = (5)(6)(4) = 120$ , and so the probability is 120/455 = .264.

d. Rather than consider many different options (choose 1, choose 2, etc.), re-frame the problem this way: at least 6 draws are required to get a 23W bulb iff a random sample of <u>five</u> bulbs fails to produce a 23W bulb. Since there are 11 non-23W bulbs, the chance of getting no 23W bulbs in a sample of size 5

$$s\binom{11}{5} / \binom{15}{5} = 462/3003 = .154.$$

41.

8.

www.konkur.in

 $(10)(10)(10)(10) = 10^4 = 10,000$ . These are the strings 0000 through 9999.

b. Count the number of prohibited sequences. There are (i) 10 with all digits identical (0000, 1111, ..., 9999); (ii) 14 with sequential digits (0123, 1234, 2345, 3456, 4567, 5678, 6789, and 7890, plus these same seven descending); (iii) 100 beginning with 19 (1900 through 1999). That's a total of 10 + 14 + 100 = 124 impermissible sequences, so there are a total of 10,000 - 124 = 9876 permissible sequences.

The chance of randomly selecting one is just  $\frac{9876}{10,000} = .9876$ .

- c. All PINs of the form 8xx1 are legitimate, so there are (10)(10) = 100 such PINs. With someone randomly selecting 3 such PINs, the chance of guessing the correct sequence is 3/100 = .03.
- **d.** Of all the PINs of the form 1xx1, eleven is prohibited: 1111, and the ten of the form 19x1. That leaves 89 possibilities, so the chances of correctly guessing the PIN in 3 tries is 3/89 = .0337.

43.

There are  $\binom{52}{5} = 2,598,960$  five-card hands. The number of 10-high straights is  $(4)(4)(4)(4)(4)(4) = 4^5 = 1024$ (any of four 6s, any of four 7s, etc.). So, P(10 high straight) =  $\frac{1024}{2,598,960} = .000394$ . Next, there ten "types of straight: A2345, 23456, ..., 910JQK, 10JQKA. So,  $P(\text{straight}) = 10 \times \frac{1024}{2,598,960} = .00394$ . Finally, there are only 40 straight flushes: each of the ten sequences above in each of the 4 suits makes (10)(4) = 40. So,  $P(\text{straight flush}) = \frac{40}{2,598,960} = .00001539$ .

### Section 2.4

#### 45.

**a.** P(A) = .106 + .141 + .200 = .447, P(C) = .215 + .200 + .065 + .020 = .500, and  $P(A \cap C) = .200$ .

**b.**  $P(A/C) = \frac{P(A \cap C)}{P(C)} = \frac{.200}{.500} = .400$ . If we know that the individual came from ethnic group 3, the

probability that he has Type A blood is .40.  $P(C|A) = \frac{P(A \cap C)}{P(A)} = \frac{.200}{.447} = .447$ . If a person has Type A blood, the probability that he is from ethnic group 3 is .447.

c. Define D = "ethnic group 1 selected." We are asked for P(D/B'). From the table,  $P(D \cap B') = .082 + .106 + .004 = .192$  and P(B') = 1 - P(B) = 1 - [.008 + .018 + .065] = .909. So, the desired probability is  $P(D/B') = \frac{P(D \cap B')}{P(B')} = \frac{.192}{.909} = .211$ .
47.

a. Apply the addition rule for three events:  $P(A \cup B \cup C) = .6 + .4 + .2 - .3 - .15 - .1 + .08 = .73$ .

**b.** 
$$P(A \cap B \cap C') = P(A \cap B) - P(A \cap B \cap C) = .3 - .08 = .22$$

c.  $P(B|A) = \frac{P(A \cap B)}{P(A)} = \frac{.3}{.6} = .50$  and  $P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{.3}{.4} = .75$ . Half of students with Visa cards also

have a MasterCard, while three-quarters of students with a MasterCard also have a Visa card.

d. 
$$P(A \cap B \mid C) = \frac{P([A \cap B] \cap C)}{P(C)} = \frac{P(A \cap B \cap C)}{P(C)} = \frac{.08}{.2} = .40.$$
  
e.  $P(A \cup B \mid C) = \frac{P([A \cup B] \cap C)}{P(C)} = \frac{P([A \cap C] \cup [B \cap C])}{P(C)}$ . Use a distributive law:  
 $= \frac{P(A \cap C) + P(B \cap C) - P([A \cap C] \cap [B \cap C])}{P(C)} = \frac{P(A \cap C) + P(B \cap C) - P(A \cap B \cap C)}{P(C)} = \frac{.15 + .1 - .08}{.2} = .85.$ 

49.

- **a.** P(small cup) = .14 + .20 = .34. P(decaf) = .20 + .10 + .10 = .40.
- **b.**  $P(\text{decaf} | \text{small}) = \frac{P(\text{small} \cap \text{decaf})}{P(\text{small})} = \frac{.20}{.34} = .588.58.8\%$  of all people who purchase a small cup of coffee choose decaf.
- c.  $P(\text{small} | \text{decaf}) = \frac{P(\text{small} \cap \text{decaf})}{P(\text{decaf})} = \frac{.20}{.40} = .50.50\%$  of all people who purchase decaf coffee choose the small size.

#### 51.

- **a.** Let  $A = \text{child has a food allergy, and } R = \text{child has a history of severe reaction. We are told that <math>P(A) = .08$  and  $P(R \mid A) = .39$ . By the multiplication rule,  $P(A \cap R) = P(A) \times P(R \mid A) = (.08)(.39) = .0312$ .
- **b.** Let M = the child is allergic to multiple foods. We are told that P(M | A) = .30, and the goal is to find P(M). But notice that M is actually a subset of A: you can't have multiple food allergies without having at least one such allergy! So, apply the multiplication rule again:  $P(M) = P(M \cap A) = P(A) \times P(M | A) = (.08)(.30) = .024.$

53.  $P(B|A) = \frac{P(A \cap B)}{P(A)} = \frac{P(B)}{P(A)} = \frac{.05}{.60} = .0833$  (since B is contained in A,  $A \cap B = B$ ).

55. Let  $A = \{\text{carries Lyme disease}\}\ \text{and } B = \{\text{carries HGE}\}\$ . We are told P(A) = .16, P(B) = .10, and  $P(A \cap B \mid A \cup B) = .10$ . From this last statement and the fact that  $A \cap B$  is contained in  $A \cup B$ ,  $P(A \cap B) \Rightarrow P(A \cap B) \Rightarrow P(A \cap B) = .10P(A \cup B) = .10[P(A) + P(B) - P(A \cap B)] = .10[.10 + .16 - P(A \cap B)] \Rightarrow$ 

$$10 = \frac{1}{P(A \cup B)} \Rightarrow P(A \cap B) = .101(A \cup B) = .101(A) + 1(B) +$$

Finally, the desired probability is  $P(A \mid B) = \frac{P(A \cap B)}{P(B)} = \frac{.02364}{.10} = .2364.$ 

- 57. P(B | A) > P(B) iff P(B | A) + P(B' | A) > P(B) + P(B'|A) iff 1 > P(B) + P(B'|A) by Exercise 56 (with the letters switched). This holds iff 1 P(B) > P(B' | A) iff P(B') > P(B' | A), QED.
- 59. The required probabilities appear in the tree diagram below.



- **a.**  $P(A_2 \cap B) = .21$ .
- **b.** By the law of total probability,  $P(B) = P(A_1 \cap B) + P(A_2 \cap B) + P(A_3 \cap B) = .455$ .
- c. Using Bayes' theorem,  $P(A_1 \mid B) = \frac{P(A_1 \cap B)}{P(B)} = \frac{.12}{.455} = .264$ ;  $P(A_2 \mid B) = \frac{.21}{.455} = .462$ ;  $P(A_3 \mid B) = 1 \frac{.21}{.455} = .462$ ;  $P(A_3 \mid B) = 1 \frac{.21}{.455} = .462$ ;  $P(A_3 \mid B) = 1 \frac{.21}{.455} = .264$ ;  $P(A_2 \mid B) = \frac{.21}{.455} = .264$ ;  $P(A_3 \mid B) = \frac{.21}{.455} = .264$ ;  $P(A_3$

.264 - .462 = .274. Notice the three probabilities sum to 1.

61. The initial ("prior") probabilities of 0, 1, 2 defectives in the batch are .5, .3, .2. Now, let's determine the probabilities of 0, 1, 2 defectives in the sample based on these three cases.

• If there are 0 defectives in the batch, clearly there are 0 defectives in the sample. P(0 def in sample | 0 def in batch) = 1.

• If there is 1 defective in the batch, the chance it's discovered in a sample of 2 equals 2/10 = .2, and the probability it isn't discovered is 8/10 = .8.

P(0 def in sample | 1 def in batch) = .8, P(1 def in sample | 1 def in batch) = .2.

If there are 2 defectives in the batch, the chance both are discovered in a sample of 2 equals

 $\frac{2}{10} \times \frac{1}{9} = .022$ ; the chance neither is discovered equals  $\frac{8}{10} \times \frac{7}{9} = .622$ ; and the chance exactly 1 is

discovered equals 1 - (.022 + .622) = .356.

P(0 def in sample | 2 def in batch) = .622, P(1 def in sample | 2 def in batch) = .356,

P(2 def in sample | 2 def in batch) = .022.

These calculations are summarized in the tree diagram below. Probabilities at the endpoints are intersectional probabilities, e.g.  $P(2 \text{ def in batch} \cap 2 \text{ def in sample}) = (.2)(.022) = .0044$ .

.50

 $(0) \stackrel{.5}{(1) .3} \quad (0) \stackrel{.8}{(2)} \quad .24 \\ (1) \stackrel{.6}{(1)} \quad .06 \\ (1) \stackrel{.622}{(2)} \quad .06 \\ .622 \quad (0) \stackrel{.1244}{(2)} \quad .0712 \\ .022 \\ (2) \quad .0044 \\ (2) \quad .0044 \\ (2) \quad .0044 \\ (2) \quad .0044 \\ (3) \stackrel{.6}{(2)} \quad .06 \\ .0$ 

- **a.** Using the tree diagram and Bayes' rule,  $P(0 \text{ def in batch} \mid 0 \text{ def in sample}) = \frac{.5}{.5 + .24 + .1244} = .578$   $P(1 \text{ def in batch} \mid 0 \text{ def in sample}) = \frac{.24}{.5 + .24 + .1244} = .278$  $P(2 \text{ def in batch} \mid 0 \text{ def in sample}) = \frac{.1244}{.5 + .24 + .1244} = .144$
- **b.** P(0 def in batch | 1 def in sample) = 0  $P(1 \text{ def in batch} | 1 \text{ def in sample}) = \frac{.06}{.06 + .0712} = .457$  $P(2 \text{ def in batch} | 1 \text{ def in sample}) = \frac{.0712}{.06 + .0712} = .543$

63.

a.



- **b.** From the top path of the tree diagram,  $P(A \cap B \cap C) = (.75)(.9)(.8) = .54$ .
- **c.** Event  $B \cap C$  occurs twice on the diagram:  $P(B \cap C) = P(A \cap B \cap C) + P(A' \cap B \cap C) = .54 + (.25)(.8)(.7) = .68.$
- **d.**  $P(C) = P(A \cap B \cap C) + P(A' \cap B \cap C) + P(A \cap B' \cap C) + P(A' \cap B' \cap C) = .54 + .045 + .14 + .015 = .74.$

e. Rewrite the conditional probability first:  $P(A | B \cap C) = \frac{P(A \cap B \cap C)}{P(B \cap C)} = \frac{.54}{.68} = .7941$ .

65. A tree diagram can help. We know that P(day) = .2, P(1-night) = .5, P(2-night) = .3; also, P(purchase | day) = .1, P(purchase | 1-night) = .3, and P(purchase | 2-night) = .2.

Apply Bayes' rule: e.g.,  $P(\text{day} | \text{purchase}) = \frac{P(\text{day} \cap \text{purchase})}{P(\text{purchase})} = \frac{(.2)(.1)}{(.2)(.1) + (.5)(.3) + (.3)(.2)} = \frac{.02}{.23} = .087.$ 

Similarly,  $P(1-\text{night} | \text{purchase}) = \frac{(.5)(.3)}{.23} = .652 \text{ and } P(2-\text{night} | \text{purchase}) = .261.$ 

67. Let T denote the event that a randomly selected person is, in fact, a terrorist. Apply Bayes' theorem, using P(T) = 1,000/300,000 = .0000033:

 $P(T|+) = \frac{P(T)P(+|T)}{P(T)P(+|T) + P(T')P(+|T')} = \frac{(.0000033)(.99)}{(.0000033)(.99) + (1 - .0000033)(1 - .999)} = .003289.$  That is to

say, roughly 0.3% of all people "flagged" as terrorists would be actual terrorists in this scenario.

69. The tree diagram below summarizes the information in the exercise (plus the previous information in Exercise 59). Probabilities for the branches corresponding to paying with credit are indicated at the far right. ("extra" = "plus")



a.  $P(\text{plus} \cap \text{fill} \cap \text{credit}) = (.35)(.6)(.6) = .1260.$ 

1

- **b.**  $P(\text{premium} \cap \text{no fill} \cap \text{credit}) = (.25)(.5)(.4) = .05.$
- c. From the tree diagram,  $P(\text{premium} \cap \text{credit}) = .0625 + .0500 = .1125$ .
- **d.** From the tree diagram,  $P(\text{fill} \cap \text{credit}) = .0840 + .1260 + .0625 = .2725$ .
- e. P(credit) = .0840 + .1400 + .1260 + .0700 + .0625 + .0500 = .5325.

F. 
$$P(\text{premium} | \text{credit}) = \frac{P(\text{premium} \cap \text{credit})}{P(\text{credit})} = \frac{.1125}{.5325} = .2113$$
.

Section 2.5

- 71.
- **a.** Since the events are independent, then A' and B' are independent, too. (See the paragraph below Equation 2.7.) Thus, P(B'|A') = P(B') = 1 .7 = .3.
- **b.** Using the addition rule,  $P(A \cup B) = P(A) + P(B) P(A \cap B) = .4 + .7 (.4)(.7) = .82$ . Since A and B are independent, we are permitted to write  $P(A \cap B) = P(A)P(B) = (.4)(.7)$ .

c. 
$$P(AB' | A \cup B) = \frac{P(AB' \cap (A \cup B))}{P(A \cup B)} = \frac{P(AB')}{P(A \cup B)} = \frac{P(A)P(B')}{P(A \cup B)} = \frac{(.4)(1-.7)}{.82} = \frac{.12}{.82} = .146$$

73. From a Venn diagram,  $P(B) = P(A' \cap B) + P(A \cap B) = P(B) \Rightarrow P(A' \cap B) = P(B) - P(A \cap B)$ . If A and B are independent, then  $P(A' \cap B) = P(B) - P(A)P(B) = [1 - P(A)]P(B) = P(A')P(B)$ . Thus, A' and B are independent.

Alternatively, 
$$P(A' | B) = \frac{P(A' \cap B)}{P(B)} = \frac{P(B) - P(A \cap B)}{P(B)} = \frac{P(B) - P(A)P(B)}{P(B)} = 1 - P(A) = P(A').$$

75. Let event E be the event that an error was signaled incorrectly.
We want P(at least one signaled incorrectly) = P(E<sub>1</sub> ∪ ... ∪ E<sub>10</sub>). To use independence, we need intersections, so apply deMorgan's law: = P(E<sub>1</sub> ∪ ... ∪ E<sub>10</sub>) = 1 - P(E'<sub>1</sub> ∩ ... ∩ E'<sub>10</sub>) . P(E') = 1 - .05 = .95, so for 10 independent points, P(E'<sub>1</sub> ∩ ... ∩ E'<sub>10</sub>) = (.95)...(.95) = (.95)<sup>10</sup>. Finally, P(E<sub>1</sub> ∪ E<sub>2</sub> ∪ ... ∪ E<sub>10</sub>) = 1 - (.95)<sup>10</sup> = .401. Similarly, for 25 points, the desired probability is 1 - (P(E'))<sup>25</sup> = 1 - (.95)<sup>25</sup> = .723.

77. Let p denote the probability that a rivet is defective.

a.  $.15 = P(\text{seam needs reworking}) = 1 - P(\text{seam doesn't need reworking}) = 1 - P(\text{no rivets are defective}) = 1 - P(1^{\text{st}} \text{ isn't def} \cap ... \cap 25^{\text{th}} \text{ isn't def}) = 1 - (1 - p)...(1 - p) = 1 - (1 - p)^{25}.$ Solve for  $p: (1 - p)^{25} = .85 \Rightarrow 1 - p = (.85)^{1/25} \Rightarrow p = 1 - .99352 = .00648.$ 

**b.** The desired condition is  $.10 = 1 - (1 - p)^{25}$ . Again, solve for  $p: (1 - p)^{25} = .90 \Rightarrow p = 1 - (.90)^{1/25} = 1 - .99579 = .00421$ .

79. Let  $A_1$  = older pump fails,  $A_2$  = newer pump fails, and  $x = P(A_1 \cap A_2)$ . The goal is to find x. From the Venn diagram below,  $P(A_1) = .10 + x$  and  $P(A_2) = .05 + x$ . Independence implies that  $x = P(A_1 \cap A_2) = P(A_1)P(A_2) = (.10 + x)(.05 + x)$ . The resulting quadratic equation,  $x^2 - .85x + .005 = 0$ , has roots x = .0059 and x = .8441. The latter is impossible, since the probabilities in the Venn diagram would then exceed 1. Therefore, x = .0059.



35

81. Using the hints, let  $P(A_i) = p$ , and  $x = p^2$ . Following the solution provided in the example,  $P(\text{system lifetime exceeds } t_0) = p^2 + p^2 - p^4 = 2p^2 - p^4 = 2x - x^2$ . Now, set this equal to .99:  $2x - x^2 = .99 \Rightarrow x^2 - 2x + .99 = 0 \Rightarrow x = 0.9 \text{ or } 1.1 \Rightarrow p = 1.049 \text{ or } .9487$ . Since the value we want is a probability and cannot exceed 1, the correct answer is p = .9487.



83.

- We'll need to know P(both detect the defect) = 1 P(at least one doesn't) = 1 .2 = .8. **a.**  $P(1^{st} detects \cap 2^{nd} doesn't) = P(1^{st} detects) - P(1^{st} does \cap 2^{nd} does) = .9 - .8 = .1.$ Similarly,  $P(1^{st} doesn't \cap 2^{nd} does) = .1$ , so P(exactly one does) = .1 + .1 = .2.
- b. P(neither detects a defect) = 1 [P(both do) + P(exactly 1 does)] = 1 [.8+.2] = 0. That is, under this model there is a 0% probability neither inspector detects a defect. As a result, P(all 3 escape) = (0)(0)(0) = 0.

85.

- **a.** Let  $D_1$  = detection on 1<sup>st</sup> fixation,  $D_2$  = detection on 2<sup>nd</sup> fixation.  $P(\text{detection in at most 2 fixations}) = P(D_1) + P(D'_1 \cap D_2)$ ; since the fixations are independent,  $P(D_1) + P(D'_1 \cap D_2) = P(D_1) + P(D'_1) P(D_2) = p + (1-p)p = p(2-p).$
- **b.** Define  $D_1, D_2, ..., D_n$  as in **a**. Then  $P(\text{at most } n \text{ fixations}) = P(D_1) + P(D'_1 \cap D_2) + P(D'_1 \cap D'_2 \cap D_3) + ... + P(D'_1 \cap D'_2 \cap \cdots \cap D'_{n-1} \cap D_n) = p + (1-p)p + (1-p)^2p + ... + (1-p)^{n-1}p = p[1 + (1-p) + (1-p)^2 + ... + (1-p)^{n-1}] = p \cdot \frac{1-(1-p)^n}{1-(1-p)} = 1-(1-p)^n$ .

Alternatively,  $P(\text{at most } n \text{ fixations}) = 1 - P(\text{at least } n+1 \text{ fixations are required}) = 1 - P(\text{no detection in } 1^{\text{st}} n \text{ fixations}) = 1 - P(D'_1 \cap D'_2 \cap \dots \cap D'_n) = 1 - (1-p)^n$ .

- c.  $P(\text{no detection in 3 fixations}) = (1-p)^3$ .
- **d.**  $P(\text{passes inspection}) = P(\{\text{not flawed}\} \cup \{\text{flawed and passes}\})$ = P(not flawed) + P(flawed and passes)
  - = .9 +  $P(\text{flawed}) P(\text{passes} | \text{flawed}) = .9 + (.1)(1-p)^3$ .
- e. Borrowing from d,  $P(\text{flawed} | \text{passed}) = \frac{P(\text{flawed} \cap \text{passed})}{P(\text{passed})} = \frac{.1(1-p)^3}{.9+.1(1-p)^3}$ . For p = .5,

$$P(\text{flawed} | \text{passed}) = \frac{.1(1-.5)^3}{.9+.1(1-.5)^3} = .0137$$

87.

## Chapter 2: Probability

- **a.** Use the information provided and the addition rule:  $P(A_1 \cup A_2) = P(A_1) + P(A_2) - P(A_1 \cap A_2) \Rightarrow P(A_1 \cap A_2) = P(A_1) + P(A_2) - P(A_1 \cup A_2) = .55 + .65 - .80$ = .40.
- **b.** By definition,  $P(A_2 | A_3) = \frac{P(A_2 \cap A_3)}{P(A_3)} = \frac{.40}{.70} = .5714$ . If a person likes vehicle #3, there's a 57.14% chance s/he will also like vehicle #2.
- c. No. From b,  $P(A_2 | A_3) = .5714 \neq P(A_2) = .65$ . Therefore,  $A_2$  and  $A_3$  are not independent. Alternatively,  $P(A_2 \cap A_3) = .40 \neq P(A_2)P(A_3) = (.65)(.70) = .455$ .
- **d.** The goal is to find  $P(A_2 \cup A_3 | A_1')$ , i.e.  $\frac{P([A_2 \cup A_3] \cap A_1')}{P(A_1')}$ . The denominator is simply 1 .55 = .45. There are several ways to calculate the numerator; the simplest approach using the information
  - provided is to draw a Venn diagram and observe that  $P([A_2 \cup A_3] \cap A_1) = P(A_1 \cup A_2 \cup A_3) P(A_1) = P(A_1 \cup A_2 \cup A_3) P(A_2 \cup A_3) P(A_3 \cup A_3) P(A_3 \cup A_3) P(A_3 \cup A_3) P(A_3 \cup A_3) = P(A_3 \cup A_3) P(A_3 \cup A_3)$
  - .88 .55 = .33. Hence,  $P(A_2 \cup A_3 \mid A_1') = \frac{.33}{.45} = .7333.$
- 89. The question asks for P(exactly) one tag lost  $| \text{ at most}}$  one tag lost) =  $P((C_1 \cap C_2) \cup (C_1' \cap C_2) | (C_1 \cap C_2)')$ . Since the first event is contained in (a subset of) the second event, this equals  $\frac{P((C_1 \cap C_2') \cup (C_1' \cap C_2))}{P((C_1 \cap C_2)')} = \frac{P(C_1 \cap C_2') + P(C_1' \cap C_2)}{1 - P(C_1 \cap C_2)} = \frac{P(C_1)P(C_2') + P(C_1')P(C_2)}{1 - P(C_1)P(C_2)}$ by independence =

$$\frac{\pi(1-\pi)+(1-\pi)\pi}{1-\pi^2} = \frac{2\pi(1-\pi)}{1-\pi^2} = \frac{2\pi}{1+\pi}.$$

# Supplementary Exercises

91.

- **a.**  $P(\text{line 1}) = \frac{500}{1500} = .333;$  $P(\text{crack}) = \frac{.50(500) + .44(400) + .40(600)}{1500} = \frac{.666}{1500} = .444.$
- **b.** This is one of the percentages provided: P(blemish | line 1) = .15.
- c.  $P(\text{surface defect}) = \frac{.10(500) + .08(400) + .15(600)}{1500} = \frac{.172}{1500};$   $P(\text{line } 1 \cap \text{surface defect}) = \frac{.10(500)}{1500} = \frac{.50}{1500};$ so,  $P(\text{line } 1 | \text{surface defect}) = \frac{.50 / 1500}{172 / 1500} = \frac{.50}{172} = .291.$

93. Apply the addition rule:  $P(A \cup B) = P(A) + P(B) - P(A \cap B) \Rightarrow .626 = P(A) + P(B) - .144$ . Apply independence:  $P(A \cap B) = P(A)P(B) = .144$ .

So, P(A) + P(B) = .770 and P(A)P(B) = .144. Let x = P(A) and y = P(B). Using the first equation, y = .77 - x, and substituting this into the second equation yields x(.77 - x) = .144 or  $x^2 - .77x + .144 = 0$ . Use the quadratic formula to solve:

$$x = \frac{.77 \pm \sqrt{(-.77)^2 - (4)(1)(.144)}}{2(1)} = \frac{.77 \pm .13}{2} = .32 \text{ or } .45. \text{ Since } x = P(A) \text{ is assumed to be the larger}$$
  
probability,  $x = P(A) = .45$  and  $y = P(B) = .32$ .

95.

- **a.** There are 5! = 120 possible orderings, so  $P(BCDEF) = \frac{1}{120} = .0833$ .
- b. The number of orderings in which F is third equals 4×3×1\*×2×1 = 24 (\*because F must be here), so P(F is third) = <sup>24</sup>/<sub>120</sub> = .2. Or more simply, since the five friends are ordered completely at random, there is a ½ chance F is specifically in position three.

c. Similarly, 
$$P(F \text{ last}) = \frac{4 \times 3 \times 2 \times 1 \times 1}{120} = .2$$

- **d.**  $P(\text{F hasn't heard after 10 times}) = P(\text{not on } \#1 \cap \text{not on } \#2 \cap \dots \cap \text{not on } \#10) = \frac{4}{5} \times \dots \times \frac{4}{5} = \left(\frac{4}{5}\right)^n = .1074.$
- 97. When three experiments are performed, there are 3 different ways in which detection can occur on exactly 2 of the experiments: (i) #1 and #2 and not #3; (ii) #1 and not #2 and #3; and (iii) not #1 and #2 and #3. If the impurity is present, the probability of exactly 2 detections in three (independent) experiments is (.8)(.8)(.2) + (.8)(.2)(.8) + (.2)(.8)(.8) = .384. If the impurity is absent, the analogous probability is 3(.1)(.1)(.9) = .027. Thus, applying Bayes' theorem, P(impurity is present | detected in exactly 2 out of 3)

99. Refer to the tree diagram below.



a. P(pass inspection) = P(pass initially ∪ passes after recrimping) =
 P(pass initially) + P(fails initially ∩ goes to recrimping ∩ is corrected after recrimping) =
 .95 + (.05)(.80)(.60) (following path "bad-good-good" on tree diagram) = .974.

- **b.**  $P(\text{needed no recrimping} | \text{passed inspection}) = \frac{P(\text{passed initially})}{P(\text{passed inspection})} = \frac{.95}{.974} = .9754$ .
- **101.** Let  $A = 1^{\text{st}}$  functions,  $B = 2^{\text{nd}}$  functions, so P(B) = .9,  $P(A \cup B) = .96$ ,  $P(A \cap B) = .75$ . Use the addition rule:  $P(A \cup B) = P(A) + P(B) - P(A \cap B) \Rightarrow .96 = P(A) + .9 - .75 \Rightarrow P(A) = .81$ .

Therefore,  $P(B \mid A) = \frac{P(B \cap A)}{P(A)} = \frac{.75}{.81} = .926.$ 

**103.** A tree diagram can also help here. **a.**  $P(E_1 \cap L) = P(E_1)P(L / E_1) = (.40)(.02) = .008.$ 

**b.** The law of total probability gives  $P(L) = \sum P(E_i)P(L | E_i) = (.40)(.02) + (.50)(.01) + (.10)(.05) = .018$ .

2. 
$$P(E'_1 | L') = 1 - P(E_1 | L') = 1 - \frac{P(E_1 \cap L')}{P(L')} = 1 - \frac{P(E_1)P(L' | E_1)}{1 - P(L)} = 1 - \frac{(.40)(.98)}{1 - .018} = .60$$

105. This is the famous "Birthday Problem" in probability.

1

a. There are  $365^{10}$  possible lists of birthdays, e.g. (Dec 10, Sep 27, Apr 1, ...). Among those, the number with zero matching birthdays is  $P_{10,365}$  (sampling ten birthdays without replacement from 365 days. So,

$$P(\text{all different}) = \frac{P_{10,365}}{365^{10}} = \frac{(365)(364)\cdots(356)}{(365)^{10}} = .883. P(\text{at least two the same}) = 1 - .883 = .117$$

- **b.** The general formula is  $P(\text{at least two the same}) = 1 \frac{P_{k,365}}{365^k}$ . By trial and error, this probability equals .476 for k = 22 and equals .507 for k = 23. Therefore, the smallest k for which k people have at least a 50-50 chance of a birthday match is 23.
- c. There are 1000 possible 3-digit sequences to end a SS number (000 through 999). Using the idea from a,  $P(\text{at least two have the same SS ending}) = 1 \frac{P_{10,1000}}{1000^{10}} = 1 .956 = .044.$

Assuming birthdays and SS endings are independent, P(at least one "coincidence") = P(birthday coincidence) = .117 + .044 - (.117)(.044) = .156.

107. P(detection by the end of the nth glimpse) = 1 - P(not detected in first n glimpses) =

$$1 - P(G'_1 \cap G'_2 \cap \dots \cap G'_n) = 1 - P(G'_1)P(G'_2) \cdots P(G'_n) = 1 - (1 - p_1)(1 - p_2) \dots (1 - p_n) = 1 - \prod_{i=1}^{n} (1 - p_i)$$

109.

a

. 
$$P(\text{all in correct room}) = \frac{1}{4!} = \frac{1}{24} = .0417.$$

b. The 9 outcomes which yield completely incorrect assignments are: 2143, 2341, 2413, 3142, 3412, 3421, 4123, 4321, and 4312, so  $P(\text{all incorrect}) = \frac{9}{24} = .375$ .

Outcome	s = 0	<i>s</i> = 1	<i>s</i> = 2	<i>s</i> = 3	Outcome	s = 0	s = 1	<i>s</i> = 2	<i>s</i> = 3	1. 1 1
1234	1	4	4	4	3124	3	1	4	4	
1234	1	3	3	3	3142	3	1	4	2	
1245	1	4	4	4	3214	3	2	1	4	
1342	1	2	2	2	3241	3	2	1	1	
1423	1	3	3	3	3412	3	1	1	2	
1423	1	2	2	2	3421	3	2	2	1	
2124	2	1	4	4	4123	4	1	3	3	
2134	2	1	3	3	4132	4	1	2	2	
2145	2	1	1	4	4213	4	2	1	3	
2314	2	1	1	1	4231	4	2	1	1	
2341	2	1	1	3	4312	4	3	1	2	
2413	2	1	1	1	4321	4	3	2	1	

111. Note: s = 0 means that the very first candidate interviewed is hired. Each entry below is the candidate hired for the given policy and outcome.

From the table, we derive the following probability distribution based on s:

5	0	1	2	3	
P(hire #1)	6	11	10	6	
1 (1110)	24	24	24	24	

Therefore s = 1 is the best policy.

113.  $P(A_1) = P(\text{draw slip 1 or } 4) = \frac{1}{2}; P(A_2) = P(\text{draw slip 2 or } 4) = \frac{1}{2};$   $P(A_3) = P(\text{draw slip 3 or } 4) = \frac{1}{2}; P(A_1 \cap A_2) = P(\text{draw slip } 4) = \frac{1}{4};$   $P(A_2 \cap A_3) = P(\text{draw slip } 4) = \frac{1}{4}; P(A_1 \cap A_3) = P(\text{draw slip } 4) = \frac{1}{4}.$ Hence  $P(A_1 \cap A_2) = P(A_1)P(A_2) = \frac{1}{4}; P(A_2 \cap A_3) = P(A_2)P(A_3) = \frac{1}{4};$  and  $P(A_1 \cap A_3) = P(A_1)P(A_3) = \frac{1}{4}.$  Thus, there exists pairwise independence. However,  $P(A_1 \cap A_2 \cap A_3) = P(\text{draw slip } 4) = \frac{1}{4} \neq \frac{1}{8} = P(A_1)P(A_2)P(A_3),$  so the events are not mutually independent.

Comparing the state was been very block and the lattice of the state of the

**CHAPTER 3** 

# Section 3.1

1.

S:	FFF	SFF	FSF	FFS	FSS	SFS	SSF	SSS	
X:	0	1	1	1	2	2	2	3	

- 3. Examples include: M = the difference between the large and the smaller outcome with possible values 0, 1, 2, 3, 4, or 5; T = 1 if the sum of the two resulting numbers is even and T = 0 otherwise, a Bernoulli random variable. See the back of the book for other examples.
- 5. No. In the experiment in which a coin is tossed repeatedly until a H results, let Y = 1 if the experiment terminates with at most 5 tosses and Y = 0 otherwise. The sample space is infinite, yet Y has only two possible values. See the back of the book for another example.
- 7.
- a. Possible values of X are 0, 1, 2, ..., 12; discrete.
- b. With n = # on the list, values of Y are 0, 1, 2, ..., N; discrete.
- c. Possible values of U are 1, 2, 3, 4, ...; discrete.
- **d.** Possible values of X are  $(0, \infty)$  if we assume that a rattlesnake can be arbitrarily short or long; not discrete.
- e. Possible values of Z are all possible sales tax percentages for online purchases, but there are only finitely-many of these. Since we could list these different percentages  $\{z_1, z_2, ..., z_N\}$ , Z is discrete.
- f. Since 0 is the smallest possible pH and 14 is the largest possible pH, possible values of Y are [0, 14]; not discrete.
- g. With m and M denoting the minimum and maximum possible tension, respectively, possible values of X are [m, M]; not discrete.
- h. The number of possible tries is 1, 2, 3, ...; each try involves 3 racket spins, so possible values of X are 3, 6, 9, 12, 15, ...; discrete.
- **a.** Returns to 0 can occur only after an even number of tosses, so possible X values are 2, 4, 6, 8, .... Because the values of X are enumerable, X is discrete.
- **b.** Now a return to 0 is possible after any number of tosses greater than 1, so possible values are 2, 3, 4, 5, .... Again, X is discrete.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

9





Therefore, the complete cdf of X is

$$F(x) = \begin{cases} 0 & x < 0\\ .30 & 0 \le x < \\ .90 & 1 \le x < \\ 1 & 2 \le x \end{cases}$$

17.

a. p(2) = P(Y=2) = P(first 2 batteries are acceptable) = P(AA) = (.9)(.9) = .81.

- **b.**  $p(3) = P(Y=3) = P(UAA \text{ or } AUA) = (.1)(.9)^2 + (.1)(.9)^2 = 2[(.1)(.9)^2] = .162.$
- c. The fifth battery must be an *A*, and exactly one of the first four must also be an *A*. Thus,  $p(5) = P(AUUUA \text{ or } UAUUA \text{ or } UUUAA) = 4[(.1)^3(.9)^2] = .00324$ .
- **d.**  $p(y) = P(\text{the } y^{\text{th}} \text{ is an } A \text{ and so is exactly one of the first } y 1) = (y 1)(.1)^{y-2}(.9)^2$ , for y = 2, 3, 4, 5, ...
- 19. p(0) = P(Y=0) = P(both arrive on Wed) = (.3)(.3) = .09; p(1) = P(Y=1) = P((W,Th) or (Th,W) or (Th,Th)) = (.3)(.4) + (.4)(.3) + (.4)(.4) = .40; p(2) = P(Y=2) = P((W,F) or (Th,F) or (F,W) or (F,Th) or (F,F)) = .32;p(3) = 1 - [.09 + .40 + .32] = .19.

21.

**a.** First, 1 + 1/x > 1 for all x = 1, ..., 9, so  $\log(1 + 1/x) > 0$ . Next, check that the probabilities sum to 1:  $\sum_{x=1}^{9} \log_{10}(1+1/x) = \sum_{x=1}^{9} \log_{10}\left(\frac{x+1}{x}\right) = \log_{10}\left(\frac{2}{1}\right) + \log_{10}\left(\frac{3}{2}\right) + \dots + \log_{10}\left(\frac{10}{9}\right); \text{ using properties of logs,}$ this equals  $\log_{10}\left(\frac{2}{1} \times \frac{3}{2} \times \dots \times \frac{10}{9}\right) = \log_{10}(10) = 1.$ 

- **b.** Using the formula  $p(x) = \log_{10}(1 + 1/x)$  gives the following values: p(1) = .301, p(2) = .176, p(3) = .125, p(4) = .097, p(5) = .079, p(6) = .067, p(7) = .058, p(8) = .051, p(9) = .046. The distribution specified by Benford's Law is <u>not</u> uniform on these nine digits; rather, lower digits (such as 1 and 2) are much more likely to be the lead digit of a number than higher digits (such as 8 and 9).
- c. The jumps in F(x) occur at 0, ..., 8. We display the cumulative probabilities here:  $F(1) = .301, F(2) = .477, F(3) = .602, F(4) = .699, F(5) = .778, F(6) = .845, F(7) = .903, F(8) = .954, F(9) = 1. So, F(x) = 0 \text{ for } x < 1; F(x) = .301 \text{ for } 1 \le x < 2; F(x) = .477 \text{ for } 2 \le x < 3; \text{ etc.}$

**d.** 
$$P(X \le 3) = F(3) = .602; P(X \ge 5) = 1 - P(X \le 5) = 1 - P(X \le 4) = 1 - F(4) = 1 - .699 = .301.$$

#### 23.

**a.** 
$$p(2) = P(X=2) = F(3) - F(2) = .39 - .19 = .20.$$

- **b.**  $P(X > 3) = 1 P(X \le 3) = 1 F(3) = 1 .67 = .33.$
- c.  $P(2 \le X \le 5) = F(5) F(2-1) = F(5) F(1) = .92 .19 = .78.$
- **d.**  $P(2 < X < 5) = P(2 < X \le 4) = F(4) F(2) = .92 .39 = .53.$

25. 
$$p(0) = P(Y = 0) = P(B \text{ first}) = p;$$
  
 $p(1) = P(Y = 1) = P(G \text{ first, then } B) = (1 - p)p;$   
 $p(2) = P(Y = 2) = P(GGB) = (1 - p)^2p;$   
Continuing,  $p(y) = P(y \text{ Gs and then } a B) = (1 - p)^y p \text{ for } y = 0,1,2,3,...$ 

27.

2

a. The sample space consists of all possible permutations of the four numbers 1, 2, 3, 4:

outcome	x value	outcome	x value	outcome	x value
1234	4	2314	1	3412	0
1243	2	2341	0	3421	0
1324	2	2413	0	4132	1
1342	1	2431	1	4123	0
1423	1	3124	1	4213	1
1432	2	3142	0	4231	2
2134	2	3214	2	4312	0
2143	0	3241	1	4321	0

**b.** From the table in **a**, 
$$p(0) = P(X=0) = \frac{9}{24}$$
,  $p(1) = P(X=1) = \frac{8}{24}$ ,  $p(2) = P(Y=2) = \frac{6}{24}$   
 $p(3) = P(X=3) = 0$ , and  $p(4) = P(Y=4) = \frac{1}{24}$ .

# Section 3.3

29.

**a.**  $E(X) = \sum_{\text{all } x} xp(x) = 1(.05) + 2(.10) + 4(.35) + 8(.40) + 16(.10) = 6.45 \text{ GB}.$ 

- **b.**  $V(X) = \sum_{\text{all } x} (x \mu)^2 p(x) = (1 6.45)^2 (.05) + (2 6.45)^2 (.10) + \dots + (16 6.45)^2 (.10) = 15.6475.$
- c.  $\sigma = \sqrt{V(X)} = \sqrt{15.6475} = 3.956$  GB.
- **d.**  $E(X^2) = \sum_{\text{all},x} x^2 p(x) = 1^2 (.05) + 2^2 (.10) + 4^2 (.35) + 8^2 (.40) + 16^2 (.10) = 57.25$ . Using the shortcut formula,  $V(X) = E(X^2) \mu^2 = 57.25 (6.45)^2 = 15.6475$ .
- 31. From the table in Exercise 12, E(Y) = 45(.05) + 46(.10) + ... + 55(.01) = 48.84; similarly,  $E(Y^2) = 45^2(.05) + 46^2(.10) + ... + 55^2(.01) = 2389.84$ ; thus  $V(Y) = E(Y^2) - [E(Y)]^2 = 2389.84 - (48.84)^2 = 4.4944$  and  $\sigma_Y = \sqrt{4.4944} = 2.12$ .

One standard deviation from the mean value of Y gives  $48.84 \pm 2.12 = 46.72$  to 50.96. So, the probability Y is within one standard deviation of its mean value equals P(46.72 < Y < 50.96) = P(Y = 47, 48, 49, 50) = .12 + .14 + .25 + .17 = .68.

33.

**a.** 
$$E(X^2) = \sum_{x=0}^{1} x^2 \cdot p(x) = 0^2 (1-p) + 1^2 (p) = p$$

**b.** 
$$V(X) = E(X^2) - [E(X)]^2 = p - [p]^2 = p(1-p)$$

c.  $E(X^{79}) = 0^{79}(1-p) + 1^{79}(p) = p$ . In fact,  $E(X^n) = p$  for any non-negative power *n*.

35. Let  $h_3(X)$  and  $h_4(X)$  equal the net revenue (sales revenue minus order cost) for 3 and 4 copies purchased, respectively. If 3 magazines are ordered (\$6 spent), net revenue is 4-56=-2 if X=1, 2(4)-56=2 if X=2, 3(4)-56=56 if X=3, and also 6 if X=4, 5, or 6 (since that additional demand simply isn't met. The values of  $h_4(X)$  can be deduced similarly. Both distributions are summarized below.

x	1	2	3	4	5	6
$h_3(x)$	-2	2	6	6	6	6
$h_4(x)$	4	0	4	8	8	8
p(x)	1	$\frac{2}{15}$	3	$\frac{4}{15}$	3	$\frac{2}{15}$

Using the table,  $E[h_3(X)] = \sum_{x=1}^{6} h_3(x) \cdot p(x) = (-2)(\frac{1}{15}) + \dots + (6)(\frac{2}{15}) = $4.93.$ Similarly,  $E[h_4(X)] = \sum_{x=1}^{6} h_4(x) \cdot p(x) = (-4)(\frac{1}{15}) + \dots + (8)(\frac{2}{15}) = $5.33.$ 

Therefore, ordering 4 copies gives slightly higher revenue, on the average.

37. Using the hint, 
$$E(X) = \sum_{x=1}^{n} x \cdot \left(\frac{1}{n}\right) = \frac{1}{n} \sum_{x=1}^{n} x = \frac{1}{n} \left[\frac{n(n+1)}{2}\right] = \frac{n+1}{2}$$
. Similarly  
 $E(X^2) = \sum_{x=1}^{n} x^2 \cdot \left(\frac{1}{n}\right) = \frac{1}{n} \sum_{x=1}^{n} x^2 = \frac{1}{n} \left[\frac{n(n+1)(2n+1)}{6}\right] = \frac{(n+1)(2n+1)}{6}$ , so  
 $V(X) = \frac{(n+1)(2n+1)}{6} - \left(\frac{n+1}{2}\right)^2 = \frac{n^2 - 1}{12}$ .

**39.** From the table,  $E(X) = \sum xp(x) = 2.3$ ,  $E(X^2) = 6.1$ , and  $V(X) = 6.1 - (2.3)^2 = .81$ . Each lot weighs 5 lbs, so the number of pounds left = 100 - 5X. Thus the expected weight left is E(100 - 5X) = 100 - 5E(X) = 88.5 lbs, and the variance of the weight left is  $V(100 - 5X) = V(-5X) = (-5)^2 V(X) = 25 V(X) = 20.25$ .

41. Use the hint: 
$$V(aX + b) = E[((aX + b) - E(aX + b))^2] = \sum [ax + b - E(aX + b)]^2 p(x) =$$

$$\sum [ax+b-(a\mu+b)]^2 p(x) = \sum [ax-a\mu]^2 p(x) = a^2 \sum (x-\mu)^2 p(x) = a^2 V(X).$$

43. With a = 1 and b = -c, E(X - c) = E(aX + b) = a E(X) + b = E(X) - c. When  $c = \mu$ ,  $E(X - \mu) = E(X) - \mu = \mu - \mu = 0$ ; i.e., the expected deviation from the mean is zero.

45

45 
$$a \le X \le b$$
 means that  $a \le x \le b$  for all x in the range of X. Hence  $ap(x) \le xp(x) \le bp(x)$  for all x, and

$$\sum ap(x) \le \sum xp(x) \le \sum bp(x)$$
$$a \ge p(x) \le \sum xp(x) \le b \ge p(x)$$
$$a \cdot 1 \le E(X) \le b \cdot 1$$
$$a \le E(X) \le b$$

# Section 3.4

47.

- **a.** B(4;15,.7) = .001.
- **b.** b(4;15,.7) = B(4;15,.7) B(3;15,.7) = .001 .000 = .001.
- c. Now p = .3 (multiple vehicles). b(6;15,.3) = B(6;15,.3) B(5;15,.3) = .869 .722 = .147.
- **d.**  $P(2 \le X \le 4) = B(4;15,.7) B(1;15,.7) = .001.$
- e.  $P(2 \le X) = 1 P(X \le 1) = 1 B(1;15,.7) = 1 .000 = 1.$
- f. The information that 11 accidents involved multiple vehicles is redundant (since n = 15 and x = 4). So, this is actually identical to **b**, and the answer is .001.
- 49. Let X be the number of "seconds," so  $X \sim Bin(6, .10)$ .

**a.** 
$$P(X=1) = {n \choose x} p^x (1-p)^{n-x} = {6 \choose 1} (.1)^1 (.9)^5 = .3543.$$

**b.**  $P(X \ge 2) = 1 - [P(X = 0) + P(X = 1)] = 1 - \left[\binom{6}{0}(.1)^{0}(.9)^{6} + \binom{6}{1}(.1)^{1}(.9)^{5}\right] = 1 - [.5314 + .3543] = 1 - [.5314 + .3544] = 1 -$ 

.1143.

- c. Either 4 or 5 goblets must be selected.
  - Select 4 goblets with zero defects:  $P(X=0) = \begin{pmatrix} 4 \\ 0 \end{pmatrix} (.1)^0 (.9)^4 = .6561.$

Select 4 goblets, one of which has a defect, and the 5<sup>th</sup> is good:  $\begin{bmatrix} 4\\1 \end{bmatrix} (.1)^1 (.9)^3 \end{bmatrix} \times .9 = .26244$ So, the desired probability is .6561 + .26244 = .91854.

- 51. Let X be the number of faxes, so  $X \sim Bin(25, .25)$ . a. E(X) = np = 25(.25) = 6.25.
  - **b.** V(X) = np(1-p) = 25(.25)(.75) = 4.6875, so SD(X) = 2.165.
  - c.  $P(X > 6.25 + 2(2.165)) = P(X > 10.58) = 1 P(X \le 10.58) = 1 P(X \le 10) = 1 B(10;25,.25) = .030.$

- 53. Let "success" = has at least one citation and define X = number of individuals with at least one citation. Then  $X \sim Bin(n = 15, p = .4)$ .
  - a. If at least 10 have no citations (failure), then at most 5 have had at least one (success):  $P(X \le 5) = B(5;15,.40) = .403.$
  - **b.** Half of 15 is 7.5, so less than half means 7 or fewer:  $P(X \le 7) = B(7; 15, .40) = .787$ .

c. 
$$P(5 \le X \le 10) = P(X \le 10) - P(X \le 4) = .991 - .217 = .774.$$

55. Let "success" correspond to a telephone that is submitted for service while under warranty and must be replaced. Then  $p = P(\text{success}) = P(\text{replaced} | \text{submitted}) \cdot P(\text{submitted}) = (.40)(.20) = .08$ . Thus X, the number among the company's 10 phones that must be replaced, has a binomial distribution with n = 10 and

$$p = .08$$
, so  $P(X=2) = {\binom{10}{2}} (.08)^2 (.92)^8 = .1478$ .

- 57. Let X = the number of flashlights that work, and let event  $B = \{\text{battery has acceptable voltage}\}$ . Then P(flashlight works) = P(both batteries work) = P(B)P(B) = (.9)(.9) = .81. We have assumed here that the batteries' voltage levels are independent. Finally,  $X \sim \text{Bin}(10, .81)$ , so  $P(X \ge 9) = P(X = 9) + P(X = 10) = .285 + .122 = .407$ .
- 59. In this example,  $X \sim Bin(25, p)$  with p unknown. a.  $P(rejecting claim when p = .8) = P(X \le 15 when p = .8) = B(15; 25, .8) = .017.$ 
  - **b.**  $P(\underline{\text{not}} \text{ rejecting claim when } p = .7) = P(X > 15 \text{ when } p = .7) = 1 P(X \le 15 \text{ when } p = .7) = 1 B(15; 25, .7) = 1 .189 = .811.$ For p = .6, this probability is = 1 - B(15; 25, .6) = 1 - .575 = .425.
  - c. The probability of rejecting the claim when p = .8 becomes B(14; 25, .8) = .006, smaller than in a above. However, the probabilities of b above increase to .902 and .586, respectively. So, by changing 15 to 14, we're making it less likely that we will reject the claim when it's true (p really is  $\ge .8$ ), but more likely that we'll "fail" to reject the claim when it's false (p really is < .8).

61. If topic A is chosen, then n = 2. When n = 2,  $P(\text{at least half received}) = P(X \ge 1) = 1 - P(X = 0) = 1$ 

$$1 - \binom{2}{0} (.9)^0 (.1)^2 = .99.$$

If topic B is chosen, then n = 4. When n = 4,  $P(\text{at least half received}) = P(X \ge 2) = 1 - P(X \le 1) = 1$ 

$$1 - \left[\binom{4}{0}(.9)^{0}(.1)^{4} + \binom{4}{1}(.9)^{1}(.1)^{3}\right] = .9963.$$

Thus topic B should be chosen if p = .9.

However, if p = .5, then the probabilities are .75 for A and .6875 for B (using the same method as above), so now A should be chosen.

**a.** 
$$b(x; n, 1-p) = \binom{n}{x} (1-p)^x (p)^{n-x} = \binom{n}{n-x} (p)^{n-x} (1-p)^x = b(n-x; n, p).$$

Conceptually, P(x S's when P(S) = 1 - p) = P(n-x F's when P(F) = p), since the two events are identical, but the labels S and F are arbitrary and so can be interchanged (if P(S) and P(F) are also interchanged), yielding P(n-x S's when P(S) = 1 - p) as desired.

- **b.** Use the conceptual idea from **a**:  $B(x; n, 1-p) = P(\underline{\text{at most } x \text{ S's when } P(\text{S}) = 1-p) = P(\underline{\text{at least } n-x \text{ F's when } P(\text{F}) = p)$ , since these are the same event  $= P(\underline{\text{at least } n-x \text{ S's when } P(\text{S}) = p)$ , since the S and F labels are arbitrary  $= 1 P(\underline{\text{at most } n-x-1 \text{ S's when } P(\text{S}) = p) = 1 B(n-x-1; n, p)$ .
- c. Whenever p > .5, (1-p) < .5 so probabilities involving X can be calculated using the results **a** and **b** in combination with tables giving probabilities only for  $p \le .5$ .
- 65.
- **a.** Although there are three payment methods, we are only concerned with S = uses a debit card and F = does not use a debit card. Thus we can use the binomial distribution. So, if X = the number of customers who use a debit card,  $X \sim Bin(n = 100, p = .2)$ . From this, E(X) = np = 100(.2) = 20, and V(X) = npq = 100(.2)(1-.2) = 16.
- **b.** With S = doesn't pay with cash, n = 100 and p = .7, so  $\mu = np = 100(.7) = 70$ , and V = 21.
- 67.
- When n = 20 and p = .5,  $\mu = 10$  and  $\sigma = 2.236$ , so  $2\sigma = 4.472$  and  $3\sigma = 6.708$ . The inequality  $|X - 10| \ge 4.472$  is satisfied if either  $X \le 5$  or  $X \ge 15$ , or  $P(|X - \mu| \ge 2\sigma) = P(X \le 5 \text{ or } X \ge 15) = .021 + .021 = .042$ . The inequality  $|X - 10| \ge 6.708$  is satisfied if either  $X \le 3$  or  $X \ge 17$ , so  $P(|X - \mu| \ge 3\sigma) = P(X \le 3 \text{ or } X \ge 17) = .001 + .001 = .002$ .

# Section 3.5

- 69. According to the problem description, X is hypergeometric with n = 6, N = 12, and M = 7.
  - a.  $P(X=4) = \frac{\binom{7}{4}\binom{5}{2}}{\binom{12}{6}} = \frac{350}{924} = .379 \cdot P(X \le 4) = 1 P(X > 4) = 1 [P(X=5) + P(X=6)] = 1 [\frac{7}{5}\binom{5}{1} + \frac{\binom{7}{6}\binom{5}{0}}{\binom{12}{6}} = 1 [.114 + .007] = 1 .121 = .879.$ 
    - **b.**  $E(X) = n \cdot \frac{M}{N} = 6 \cdot \frac{7}{12} = 3.5; \ V(X) = \left(\frac{12-6}{12-1}\right) 6\left(\frac{7}{12}\right) \left(1-\frac{7}{12}\right) = 0.795; \ \sigma = 0.892. \ \text{So},$  $P(X > \mu + \sigma) = P(X > 3.5 + 0.892) = P(X > 4.392) = P(X = 5 \text{ or } 6) = .121 \ \text{(from part a)}.$
    - c. We can approximate the hypergeometric distribution with the binomial if the population size and the number of successes are large. Here, n = 15 and M/N = 40/400 = .1, so h(x;15, 40, 400) ≈ b(x;15, .10). Using this approximation, P(X≤5) ≈ B(5; 15, .10) = .998 from the binomial tables. (This agrees with the exact answer to 3 decimal places.)

71.

a. Possible values of X are 5, 6, 7, 8, 9, 10. (In order to have less than 5 of the granite, there would have to be more than 10 of the basaltic). X is hypergeometric, with n = 15, N = 20, and M = 10. So, the pmf of X is

$$p(x) = h(x; 15, 10, 20) = \frac{\binom{10}{x}\binom{10}{15-x}}{\binom{20}{15}}.$$

The pmf is also provided in table form below.

x
 5
 6
 7
 8
 9
 10

 
$$p(x)$$
 .0163
 .1354
 .3483
 .3483
 .1354
 .0163

**b.** P(all 10 of one kind or the other) = P(X=5) + P(X=10) = .0163 + .0163 = .0326.

c. 
$$\mu = n \cdot \frac{M}{N} = 15 \cdot \frac{10}{20} = 7.5; V(X) = \left(\frac{20 - 15}{20 - 1}\right) 15 \left(\frac{10}{20}\right) \left(1 - \frac{10}{20}\right) = .9868; \sigma = .9934.$$

 $\mu \pm \sigma = 7.5 \pm .9934 = (6.5066, 8.4934)$ , so we want P(6.5066 < X < 8.4934). That equals P(X = 7) + P(X = 8) = .3483 + .3483 = .6966.

73.

1

The successes here are the top M = 10 pairs, and a sample of n = 10 pairs is drawn from among the N a. = 20. The probability is therefore  $h(x; 10, 10, 20) = \frac{\binom{10}{x}\binom{10}{10-x}}{\binom{20}{10}}$ .

Let X = the number among the top 5 who play east-west. (Now, M = 5.) b. Then P(all of top 5 play the same direction) = P(X = 5) + P(X = 0) =

$$h(5; 10, 5, 20) + h(5; 10, 5, 20) = \frac{\binom{5}{5}\binom{15}{5}}{\binom{20}{10}} + \frac{\binom{5}{0}\binom{15}{10}}{\binom{20}{10}} = .033.$$

Generalizing from earlier parts, we now have N = 2n; M = n. The probability distribution of X is c.

hypergeometric:  $p(x) = h(x; n, n, 2n) = \frac{\binom{n}{x}\binom{n}{n-x}}{\binom{2n}{n}}$  for x = 0, 1, ..., n. Also,  $E(X) = n \cdot \frac{n}{2n} = \frac{1}{2}n \text{ and } V(X) = \left(\frac{2n-n}{2n-1}\right) \cdot n \cdot \frac{n}{2n} \cdot \left(1 - \frac{n}{2n}\right) = \frac{n^2}{4(2n-1)}.$ 

75. Let X = the number of boxes that do <u>not</u> contain a prize until you find 2 prizes. Then  $X \sim NB(2, .2)$ . a. With S = a female child and F = a male child, let X = the number of F's before the 2<sup>nd</sup> S. Then

$$P(X = x) = nb(x; 2, .2) = {\binom{x+2-1}{2-1}}(.2)^2(1-.2)^x = (x+1)(.2)^2(.8)^x.$$

- **b.**  $P(4 \text{ boxes purchased}) = P(2 \text{ boxes without prizes}) = P(X = 2) = nb(2; 2, .2) = (2 + 1)(.2)^2(.8)^2 = .0768.$
- c.  $P(\text{at most 4 boxes purchased}) = P(X \le 2) = \sum_{x=0}^{2} nb(x; 2, .8) = .04 + .064 + .0768 = .1808.$
- **d.**  $E(X) = \frac{r(1-p)}{p} = \frac{2(1-2)}{.2} = 8$ . The total number of boxes you expect to buy is 8 + 2 = 10.
- 77. This is identical to an experiment in which a single family has children until exactly 6 females have been born (since p = .5 for each of the three families). So,

 $p(x) = nb(x; 6, .5) = {\binom{x+5}{5}}(.5)^6(1-.5)^x = {\binom{x+5}{5}}(.5)^{6+x}$ . Also,  $E(X) = \frac{r(1-p)}{p} = \frac{6(1-.5)}{.5} = 6$ ; notice this is just 2 + 2 + 2, the sum of the expected number of males born to each family.

# Section 3.6

- 79. All these solutions are found using the cumulative Poisson table,  $F(x; \mu) = F(x; 1)$ . a.  $P(X \le 5) = F(5; 1) = .999$ .
  - **b.**  $P(X=2) = \frac{e^{-1}1^2}{2!} = .184$ . Or, P(X=2) = F(2; 1) F(1; 1) = .920 .736 = .184.
  - c.  $P(2 \le X \le 4) = P(X \le 4) P(X \le 1) = F(4; 1) F(1; 1) = .260.$
  - **d.** For X Poisson,  $\sigma = \sqrt{\mu} = 1$ , so  $P(X > \mu + \sigma) = P(X > 2) = 1 P(X \le 2) = 1 F(2; 1) = 1 .920 = .080$ .
- 81. Let  $X \sim \text{Poisson}(\mu = 20)$ . a.  $P(X \le 10) = F(10; 20) = .011$ .
  - **a.**  $P(X \le 10) = P(10, 20) = .011.$
  - **b.** P(X > 20) = 1 F(20; 20) = 1 .559 = .441.
  - c.  $P(10 \le X \le 20) = F(20; 20) F(9; 20) = .559 .005 = .554;$ P(10 < X < 20) = F(19; 20) - F(10; 20) = .470 - .011 = .459.
  - **d.**  $E(X) = \mu = 20$ , so  $\sigma = \sqrt{20} = 4.472$ . Therefore,  $P(\mu 2\sigma < X < \mu + 2\sigma) = P(20 8.944 < X < 20 + 8.944) = P(11.056 < X < 28.944) = P(X \le 28) P(X \le 11) = F(28; 20) F(11; 20) = .966 .021 = .945.$

83. The exact distribution of X is binomial with n = 1000 and p = 1/200; we can approximate this distribution by the Poisson distribution with  $\mu = np = 5$ . a.  $P(5 \le X \le 8) = F(8; 5) - F(4; 5) = .492$ .

**b.** 
$$P(X \ge 8) = 1 - P(X \le 7) = 1 - F(7; 5) = 1 - .867 = .133.$$

85.

- **a.**  $\mu = 8$  when t = 1, so  $P(X = 6) = \frac{e^{-8}8^6}{6!} = .122$ ;  $P(X \ge 6) = 1 F(5; 8) = .809$ ; and  $P(X \ge 10) = 1 F(9; 8) = .283$ .
- **b.** t = 90 min = 1.5 hours, so  $\mu = 12$ ; thus the expected number of arrivals is 12 and the standard deviation is  $\sigma = \sqrt{12} = 3.464$ .
- c. t = 2.5 hours implies that  $\mu = 20$ . So,  $P(X \ge 20) = 1 F(19; 20) = .530$  and  $P(X \le 10) = F(10; 20) = .011$ .

87.

- **a.** For a two hour period the parameter of the distribution is  $\mu = \alpha t = (4)(2) = 8$ , so  $P(X = 10) = \frac{e^{-8} 8^{10}}{10!} = .099$ .
- **b.** For a 30-minute period,  $\alpha t = (4)(.5) = 2$ , so  $P(X = 0) = \frac{e^{-2}2^0}{0!} = .135$ .
- c. The expected value is simply  $E(X) = \alpha t = 2$ .
- 89. In this example,  $\alpha$  = rate of occurrence = 1/(mean time between occurrences) = 1/.5 = 2. a. For a two-year period,  $\mu = \alpha t = (2)(2) = 4$  loads.
  - **b.** Apply a Poisson model with  $\mu = 4$ :  $P(X > 5) = 1 P(X \le 5) = 1 F(5; 4) = 1 .785 = .215$ .
  - c. For  $\alpha = 2$  and the value of t unknown, P(no loads occur during the period of length t) =  $P(X=0) = \frac{e^{-2t}(2t)^0}{0!} = e^{-2t}$ . Solve for t:  $e^{-2t} \le .1 \Rightarrow -2t \le \ln(.1) \Rightarrow t \ge 1.1513$  years.

91.

- **a.** For a quarter-acre (.25 acre) plot, the mean parameter is  $\mu = (80)(.25) = 20$ , so  $P(X \le 16) = F(16; 20) = .221$ .
- **b.** The expected number of trees is  $\alpha$ -(area) = 80 trees/acre (85,000 acres) = 6,800,000 trees.
- c. The area of the circle is  $\pi r^2 = \pi (.1)^2 = .01\pi = .031416$  square miles, which is equivalent to .031416(640) = 20.106 acres. Thus X has a Poisson distribution with parameter  $\mu = \alpha(20.106) = 80(20.106) = 1608.5$ . That is, the pmf of X is the function p(x; 1608.5).

# 93.

a. No events occur in the time interval  $(0, t + \Delta t)$  if and only if no events occur in (0, t) and no events occur in  $(t, t + \Delta t)$ . Since it's assumed the numbers of events in non-overlapping intervals are independent (Assumption 3),

 $\begin{array}{l}P(\text{no events in } (0, t + \Delta t)) = P(\text{no events in } (0, t)) \cdot P(\text{no events in } (t, t + \Delta t)) \Rightarrow\\P_0(t + \Delta t) = P_0(t) \cdot P(\text{no events in } (t, t + \Delta t)) = P_0(t) \cdot [1 - \alpha \Delta t - o(\Delta t)] \text{ by Assumption 2.}\end{array}$ 

- **b.** Rewrite **a** as  $P_0(t + \Delta t) = P_0(t) P_0(t)[\alpha\Delta t + o(\Delta t)]$ , so  $P_0(t + \Delta t) P_0(t) = -P_0(t)[\alpha\Delta t + o(\Delta t)]$  and  $\frac{P_0(t + \Delta t) - P_0(t)}{\Delta t} = -\alpha P_0(t) - P_0(t) \cdot \frac{o(\Delta t)}{\Delta t}$ . Since  $\frac{o(\Delta t)}{\Delta t} \to 0$  as  $\Delta t \to 0$  and the left-hand side of the equation converges to  $\frac{dP_0(t)}{dt}$  as  $\Delta t \to 0$ , we find that  $\frac{dP_0(t)}{dt} = -\alpha P_0(t)$ .
- c. Let  $P_0(t) = e^{-\alpha t}$ . Then  $\frac{dP_0(t)}{dt} = \frac{d}{dt} [e^{-\alpha t}] = -\alpha e^{-\alpha t} = -\alpha P_0(t)$ , as desired. (This suggests that the probability of zero events in (0, t) for a process defined by Assumptions 1-3 is equal to  $e^{-\alpha t}$ .)

**d.** Similarly, the product rule implies 
$$\frac{d}{dt} \left[ \frac{e^{-\alpha t} (\alpha t)^k}{k!} \right] = \frac{-\alpha e^{-\alpha t} (\alpha t)^k}{k!} + \frac{k \alpha e^{-\alpha t} (\alpha t)^{k-1}}{k!} = -\alpha \frac{e^{-\alpha t} (\alpha t)^k}{k!} + \alpha \frac{e^{-\alpha t} (\alpha t)^{k-1}}{(k-1)!} = -\alpha P_k(t) + \alpha P_{k-1}(t), \text{ as desired.}$$

# Supplementary Exercises

95.

a. We'll find p(1) and p(4) first, since they're easiest, then p(2). We can then find p(3) by subtracting the others from 1.

$$p(1) = P(\text{exactly one suit}) = P(\text{all } \bigstar) + P(\text{all } \bigstar) + P(\text{all } \bigstar) + P(\text{all } \bigstar) = P(\text{all } \bigstar)$$

$$4 \cdot P(\text{all} \bullet) = 4 \cdot \frac{\binom{13}{5}\binom{39}{0}}{\binom{52}{5}} = .00198 \text{, since there are } 13 \bullet \text{s and } 39 \text{ other cards.}$$

$$p(4) = 4 \cdot P(2 \bullet, 1 \bullet, 1 \bullet, 1 \bullet) = 4 \cdot \frac{\binom{13}{2}\binom{13}{1}\binom{13}{1}\binom{13}{1}}{\binom{52}{5}} = .26375$$

 $p(2) = P(\text{all } \forall \text{s and } \text{ s, with } \geq \text{ one of each}) + \ldots + P(\text{all } \text{ s and } \text{ s with } \geq \text{ one of each}) =$ 

$$(P(all \forall s and \bigstar s, with \ge one of each) =$$

(12)(20)

$$[P(1 \lor \text{and } 4 \blacklozenge) + P(2 \lor \text{and } 3 \blacklozenge) + P(3 \lor \text{and } 2 \blacklozenge) + P(4 \lor \text{and } 1 \blacklozenge)] = 0$$

$$6 \cdot \left[ 2 \cdot \frac{\binom{13}{4}\binom{13}{1}}{\binom{52}{5}} + 2 \cdot \frac{\binom{13}{3}\binom{13}{2}}{\binom{52}{5}} \right] = 6 \left[ \frac{18,590 + 44,616}{2,598,960} \right] = .14592 \, .$$

Finally, 
$$p(3) = 1 - [p(1) + p(2) + p(4)] = .58835$$
.

**b.** 
$$\mu = \sum_{x=1}^{4} x \cdot p(x) = 3.114$$
;  $\sigma^2 = \left[\sum_{x=1}^{4} x^2 \cdot p(x)\right] - (3.114)^2 = .405 \implies \sigma = .636.$ 

#### 97.

1

**a.** From the description,  $X \sim Bin(15, .75)$ . So, the pmf of X is b(x; 15, .75).

**b.** 
$$P(X > 10) = 1 - P(X \le 10) = 1 - B(10;15, .75) = 1 - .314 = .686.$$

c.  $P(6 \le X \le 10) = B(10; 15, .75) - B(5; 15, .75) = .314 - .001 = .313.$ 

**d.** 
$$\mu = (15)(.75) = 11.75, \sigma^2 = (15)(.75)(.25) = 2.81.$$

- e. Requests can all be met if and only if  $X \le 10$ , and  $15 X \le 8$ , i.e. iff  $7 \le X \le 10$ . So,  $P(\text{all requests met}) = P(7 \le X \le 10) = B(10; 15, .75) - B(6; 15, .75) = .310.$
- 99. Let X = the number of components out of 5 that function, so  $X \sim Bin(5, .9)$ . Then a 3-out-of 5 system works when X is at least 3, and  $P(X \ge 3) = 1 P(X \le 2) = 1 B(2; 5, .9) = .991$ .

#### 101.

**a.**  $X \sim \text{Bin}(n = 500, p = .005)$ . Since *n* is large and *p* is small, *X* can be approximated by a Poisson distribution with  $\mu = np = 2.5$ . The approximate pmf of *X* is  $p(x; 2.5) = \frac{e^{-2.5} 2.5^x}{x!}$ .

**b.** 
$$P(X=5) = \frac{e^{-2.5} 2.5^5}{5!} = .0668.$$

$$P(X ≥ 5) = 1 - P(X ≤ 4) = 1 - p(4; 2.5) = 1 - .8912 = .1088.$$

103. Let *Y* denote the number of tests carried out. For n = 3, possible *Y* values are 1 and 4.  $P(Y = 1) = P(\text{no one has the disease}) = (.9)^3 = .729$  and P(Y = 4) = 1 - .729 = .271, so E(Y) = (1)(.729) + (4)(.271) = 1.813, as contrasted with the 3 tests necessary without group testing. For n = 5, possible values of *Y* are 1 and 6.  $P(Y = 1) = P(\text{no one has the disease}) = (.9)^5 = .5905$ , so P(Y = 6) = 1 - .5905 = .4095 and E(Y) = (1)(.5905) + (6)(.4095) = 3.0475, less than the 5 tests necessary

without group testing.

**105.** 
$$p(2) = P(X = 2) = P(SS) = p^2$$
, and  $p(3) = P(FSS) = (1 - p)p^2$ .

For  $x \ge 4$ , consider the first x - 3 trials and the last 3 trials separately. To have X = x, it must be the case that the last three trials were *FSS*, and that two-successes-in-a-row was <u>not</u> already seen in the first x - 3 tries.

The probability of the first event is simply  $(1 - p)p^2$ .

The second event occurs if two-in-a-row hadn't occurred after 2 or 3 or ... or x - 3 tries. The probability of this second event equals 1 - [p(2) + p(3) + ... + p(x - 3)]. (For x = 4, the probability in brackets is empty; for x = 5, it's p(2); for x = 6, it's p(2) + p(3); and so on.)

Finally, since trials are independent,  $P(X = x) = (1 - [p(2) + ... + p(x - 3)]) \cdot (1 - p)p^2$ .

For p = .9, the pmf of X up to x = 8 is shown below.



So, 
$$P(X \le 8) = p(2) + \dots + p(8) = .9995$$
.

107.

**a.** Let event A = seed carries single spikelets, and event B = seed produces ears with single spikelets. Then  $P(A \cap B) = P(A) \cdot P(B \mid A) = (.40)(.29) = .116$ .

Next, let X = the number of seeds out of the 10 selected that meet the condition  $A \cap B$ . Then X~

Bin(10, .116). So, 
$$P(X = 5) = {10 \choose 5} (.116)^5 (.884)^5 = .002857$$
.

**b.** For any one seed, the event of interest is B = seed produces ears with single spikelets. Using the law of total probability,  $P(B) = P(A \cap B) + P(A' \cap B) = (.40)(.29) + (.60)(.26) = .272$ . Next, let Y = the number out of the 10 seeds that meet condition B. Then  $Y \sim Bin(10, .272)$ . P(Y = 5) =

$$\binom{10}{5}(.272)^{5}(1-.272)^{5} = .0767, \text{ while}$$

$$P(Y \le 5) = \sum_{y=0}^{5} \binom{10}{y} (.272)^{y} (1-.272)^{10-y} = .041813 + \dots + .076719 = .97024$$

109.

**a.** 
$$P(X=0) = F(0; 2)$$
 or  $\frac{e^{-2}2^0}{0!} = 0.135$ .

- **b.** Let S = an operator who receives no requests. Then the number of operators that receive no requests follows a Bin(n = 5, p = .135) distribution. So,  $P(4 \ S' \text{s in 5 trials}) = b(4; 5, .135) = {\binom{5}{4}}(.135)^4(.865)^1 = .00144$ .
- c. For any non-negative integer x, P(all operators receive exactly x requests) =

 $P(\text{first operator receives } x) \cdot \ldots \cdot P(\text{fifth operator receives } x) = [p(x; 2)]^5 = \left[\frac{e^{-2}2^x}{x!}\right]^5 = \frac{e^{-10}2^{5x}}{(x!)^5}.$ Then, P(all receive the same number) = P(all receive 0 requests) + P(all receive 1 request) +

111.

The number of magazine copies sold is X so long as X is no more than five; otherwise, all five copies are sold. So, mathematically, the number sold is  $\min(X, 5)$ , and  $E[\min(x, 5)] = \sum_{i=1}^{\infty} \min(x, 5) p(x; 4) = 0p(0; 4) + 0p(0; 4)$ 

$$1p(1; 4) + 2p(2; 4) + 3p(3; 4) + 4p(4; 4) + \sum_{x=5}^{\infty} 5p(x; 4) =$$
  
$$1.735 + 5\sum_{x=5}^{\infty} p(x; 4) = 1.735 + 5\left[1 - \sum_{x=0}^{4} p(x; 4)\right] = 1.735 + 5[1 - F(4; 4)] = 3.594$$

113.

- a. No, since the probability of a "success" is not the same for all tests.
- **b.** There are four ways exactly three could have positive results. Let D represent those with the disease and D' represent those without the disease.

#### Probability

CombinationDD'03

1 2

$$\begin{bmatrix} \binom{5}{0}(.2)^{9}(.8)^{5} \\ = (.32768)(.0729) = .02389 \\ \begin{bmatrix} \binom{5}{1}(.2)^{1}(.8)^{4} \\ = (.4096)(.0081) = .00332 \\ \end{bmatrix} \begin{bmatrix} \binom{5}{2}(.2)^{2}(.8)^{3} \\ \end{bmatrix} \cdot \begin{bmatrix} \binom{5}{1}(.9)^{1}(.1)^{4} \\ \end{bmatrix}$$

3

0

2 1

)5	(.9) <sup>0</sup> (.1	$\binom{5}{0}$	].[	$(.8)^{2}$	(.2)	(5)
7		(0)	ΙL		/	(3)
	00000	= 0	1)	)( 0000	512)	=( 0

=(.2048)(.00045) = .00009216

Adding up the probabilities associated with the four combinations yields 0.0273.

115.

**a.** Notice that  $p(x; \mu_1, \mu_2) = .5 \ p(x; \mu_1) + .5 \ p(x; \mu_2)$ , where both terms  $p(x; \mu_i)$  are Poisson pmfs. Since both pmfs are  $\ge 0$ , so is  $p(x; \mu_1, \mu_2)$ . That verifies the first requirement.

Next,  $\sum_{x=0}^{\infty} p(x; \mu_1, \mu_2) = .5 \sum_{x=0}^{\infty} p(x; \mu_1) + .5 \sum_{x=0}^{\infty} p(x; \mu_2) = .5 + .5 = 1$ , so the second requirement for a pmf is met. Therefore,  $p(x; \mu_1, \mu_2)$  is a valid pmf.

**b.** 
$$E(X) = \sum_{x=0}^{\infty} x \cdot p(x; \mu_1, \mu_2) = \sum_{x=0}^{\infty} x[.5p(x; \mu_1) + .5p(x; \mu_2)] = .5\sum_{x=0}^{\infty} x \cdot p(x; \mu_1) + .5\sum_{x=0}^{\infty} x \cdot p(x; \mu_2) = .5E(X_1) + .5E(X_2), \text{ where } X_i \sim \text{Poisson}(\mu_i). \text{ Therefore, } E(X) = .5\mu_1 + .5\mu_2.$$

c. This requires using the variance shortcut. Using the same method as in b,

 $E(X^2) = .5\sum_{x=0}^{\infty} x^2 \cdot p(x;\mu_1) + .5\sum_{x=0}^{\infty} x^2 \cdot p(x;\mu_2) = .5E(X_1^2) + .5E(X_2^2).$  For any Poisson rv,  $E(X^2) = V(X) + [E(X)]^2 = \mu + \mu^2, \text{ so } E(X^2) = .5(\mu_1 + \mu_1^2) + .5(\mu_2 + \mu_2^2).$ Finally,  $V(X) = .5(\mu_1 + \mu_1^2) + .5(\mu_2 + \mu_2^2) - [.5\mu_1 + .5\mu_2]^2$ , which can be simplified to equal  $.5\mu_1 + .5\mu_2 + .25(\mu_1 - \mu_2)^2$ .

**d.** Simply replace the weights .5 and .5 with .6 and .4, so  $p(x; \mu_1, \mu_2) = .6 p(x; \mu_1) + .4 p(x; \mu_2)$ .

117. 
$$P(X=j) = \sum_{i=1}^{10} P(\text{arm on track } i \cap X=j) = \sum_{i=1}^{10} P(X=j \mid \text{arm on } i) \cdot p_i = \sum_{i=1}^{10} P(\text{next seek at } i+j+1 \text{ or } i-j-1) \cdot p_i = \sum_{i=1}^{10} (p_{i+j+1}+p_{i-j-1})p_i$$
, where in the summation we take  $p_k = 0$  if  $k < 0$  or  $k > 10$ .

119. Using the hint,  $\sum_{\text{all }x} (x - \mu)^2 p(x) \ge \sum_{x:|x - \mu| \ge k\sigma} (x - \mu)^2 p(x) \ge \sum_{x:|x - \mu| \ge k\sigma} (k\sigma)^2 p(x) = k^2 \sigma^2 \sum_{x:|x - \mu| \ge k\sigma} p(x)$ . The left-hand side is, by definition,  $\sigma^2$ . On the other hand, the summation on the right-hand side represents  $P(|X - \mu| \ge k\sigma)$ . So  $\sigma^2 \ge k^2 \sigma^2 \cdot P(|X - \mu| \ge k\sigma)$ , whence  $P(|X - \mu| \ge k\sigma) \le 1/k^2$ .

- 121.
- a. Let  $A_1 = \{\text{voice}\}, A_2 = \{\text{data}\}, \text{ and } X = \text{duration of a call. Then } E(X) = E(X|A_1)P(A_1) + E(X|A_2)P(A_2) = 3(.75) + 1(.25) = 2.5 \text{ minutes.}$
- **b.** Let X = the number of chips in a cookie. Then E(X) = E(X|i=1)P(i=1) + E(X|i=2)P(i=2) + E(X|i=3)P(i=3). If X is Poisson, then its mean is the specified  $\mu$  that is, E(X|i) = i+1. Therefore, E(X) = 2(.20) + 3(.50) + 4(.30) = 3.1 chips.

**CHAPTER 4** 

# Section 4.1

1.

**a.** The pdf is the straight-line function graphed below on [3, 5]. The function is clearly non-negative; to verify its integral equals 1, compute:

 $\int_{3}^{5} (.075x + .2) dx = .0375x^{2} + .2x \Big]_{3}^{5} = (.0375(5)^{2} + .2(5)) - (.0375(3)^{2} + .2(3))$ = 1.9375 - .9375 = 1



**b.**  $P(X \le 4) = \int_{3}^{4} (.075x + .2) dx = .0375x^{2} + .2x \Big]_{3}^{4} = (.0375(4)^{2} + .2(4)) - (.0375(3)^{2} + .2(3))$ = 1.4 - .9375 = .4625. Since X is a continuous rv,  $P(X \le 4) = P(X \le 4) = .4625$  as well.

c. 
$$P(3.5 \le X \le 4.5) = \int_{3.5}^{4.5} (.075x + .2) dx = .0375x^2 + .2x \Big]_{3.5}^{4.5} = \dots = .5$$
.  
 $P(4.5 < X) = P(4.5 \le X) = \int_{4.5}^{5} (.075x + .2) dx = .0375x^2 + .2x \Big]_{4.5}^{5} = \dots = .278125$ .



57

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

3.

a.

**b.** 
$$P(X > 0) = \int_0^2 .09375(4 - x^2) dx = .09375\left(4x - \frac{x^3}{3}\right) \Big]_0^2 = .5$$
.

This matches the symmetry of the pdf about x = 0.

c. 
$$P(-1 < X < 1) = \int_{-1}^{1} .09375(4 - x^2) dx = .6875$$

**d.** 
$$P(X < -.5 \text{ or } X > .5) = 1 - P(-.5 \le X \le .5) = 1 - \int_{-.5}^{.5} .09375(4 - x^2) dx = 1 - .3672 = .6328.$$

5.



**d.** 
$$P(X \ge 1.5) = 1 - \int_{1.5}^{2} \frac{3}{8} x^2 dx = \frac{1}{8} x^3 \Big]_{1.5}^{2} = \frac{1}{8} (2)^3 - \frac{1}{8} (1.5)^3 = .578125$$

7.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

c.  $P(\mu - 1 \le X \le \mu + 1) = \int_{\mu - 1}^{\mu + 1} \frac{1}{4.05} dx = \frac{2}{4.05} = .494$ . (We don't actually need to know  $\mu$  here, but it's clearly the midpoint of 2.225 mm by symmetry.)

**d.** 
$$P(a \le X \le a+1) = \int_{a}^{a+1} \frac{1}{4.05} dx = \frac{1}{4.05} = .247$$

9.

**a.** 
$$P(X \le 5) = \int_{1}^{5} .15e^{-.15(x-1)} dx = .15\int_{0}^{4} e^{-.15u} du$$
 (after the substitution  $u = x - 1$ )  
=  $-e^{-.15u} \int_{0}^{4} = 1 - e^{-.6} \approx .451$ .  $P(X > 5) = 1 - P(X \le 5) = 1 - .451 = .549$ .

**b.** 
$$P(2 \le X \le 5) = \int_{2}^{5} .15e^{-.15(x-1)} dx = \int_{1}^{4} .15e^{-.15u} du = -e^{-.15u} \Big]_{1}^{4} = .312.$$

# Section 4.2

11.

**a.** 
$$P(X \le 1) = F(1) = \frac{1^2}{4} = .25$$
.

**b.** 
$$P(.5 \le X \le 1) = F(1) - F(.5) = \frac{1^2}{4} - \frac{.5^2}{4} = .1875.$$

c. 
$$P(X > 1.5) = 1 - P(X \le 1.5) = 1 - F(1.5) = 1 - \frac{1.5^2}{4} = .4375.$$

**d.** 
$$.5 = F(\tilde{\mu}) = \frac{\tilde{\mu}^2}{4} \Rightarrow \tilde{\mu}^2 = 2 \Rightarrow \tilde{\mu} = \sqrt{2} \approx 1.414$$

e. 
$$f(x) = F'(x) = \frac{x}{2}$$
 for  $0 \le x < 2$ , and  $= 0$  otherwise.  
f.  $E(X) = \int_{-\infty}^{\infty} x \cdot f(x) dx = \int_{0}^{2} x \cdot \frac{x}{2} dx = \frac{1}{2} \int_{0}^{2} x^{2} dx = \frac{x^{3}}{6} \Big]_{0}^{2} = \frac{8}{6} \approx 1.333$ .

g. 
$$E(X^2) = \int_{-\infty}^{\infty} x^2 f(x) dx = \int_0^2 x^2 \frac{x}{2} dx = \frac{1}{2} \int_0^2 x^3 dx = \frac{x^4}{8} \Big]_0^2 = 2$$
, so  $V(X) = E(X^2) - [E(X)]^2 = 2 - \left(\frac{8}{6}\right)^2 = \frac{8}{36} \approx .222$ , and  $\sigma_X = \sqrt{.222} = .471$ .

h. From g,  $E(X^2) = 2$ .

**a.** 
$$1 = \int_{1}^{\infty} \frac{k}{x^4} dx = k \int_{1}^{\infty} x^{-4} dx = \frac{k}{-3} x^{-3} \Big]_{1}^{\infty} = 0 - \left(\frac{k}{-3}\right) (1)^{-3} = \frac{k}{3} \Longrightarrow k = 3.$$

**b.** For  $x \ge 1$ ,  $F(x) = \int_{-\infty}^{x} f(y) dy = \int_{1}^{x} \frac{3}{y^{4}} dy = -y^{-3} \Big|_{1}^{x} = -x^{-3} + 1 = 1 - \frac{1}{x^{3}}$ . For x < 1, F(x) = 0 since the distribution begins at 1. Put together,  $F(x) = \begin{cases} 0 & x < 1 \\ 1 - \frac{1}{x^{3}} & 1 \le x \end{cases}$ .

c. 
$$P(X>2) = 1 - F(2) = 1 - \frac{7}{8} = \frac{1}{8}$$
 or .125;  
 $P(2 < X < 3) = F(3) - F(2) = (1 - \frac{1}{27}) - (1 - \frac{1}{8}) = .963 - .875 = .088$ .

- **d.** The mean is  $E(X) = \int_{1}^{\infty} x \left(\frac{3}{x^{4}}\right) dx = \int_{1}^{\infty} \left(\frac{3}{x^{3}}\right) dx = -\frac{3}{2} x^{-2} \Big|_{1}^{\infty} = 0 + \frac{3}{2} = \frac{3}{2} = 1.5$ . Next,  $E(X^{2}) = \int_{1}^{\infty} x^{2} \left(\frac{3}{x^{4}}\right) dx = \int_{1}^{\infty} \left(\frac{3}{x^{2}}\right) dx = -3x^{-1} \Big|_{1}^{\infty} = 0 + 3 = 3$ , so  $V(X) = 3 - (1.5)^{2} = .75$ . Finally, the standard deviation of X is  $\sigma = \sqrt{.75} = .866$ .
- e. P(1.5 .866 < X < 1.5 + .866) = P(.634 < X < 2.366) = F(2.366) F(.634) = .9245 0 = .9245.

15.

13.

a. Since X is limited to the interval (0, 1), F(x) = 0 for  $x \le 0$  and F(x) = 1 for  $x \ge 1$ . For  $0 \le x \le 1$ ,

$$F(x) = \int_{-\infty}^{x} f(y) dy = \int_{0}^{x} 90y^{8}(1-y) dy = \int_{0}^{x} (90y^{8} - 90y^{9}) dy = 10y^{9} - 9y^{10} \int_{0}^{10} = 10x^{9} - 9x^{10}$$

The graphs of the pdf and cdf of X appear below.



- **b.**  $F(.5) = 10(.5)^9 9(.5)^{10} = .0107.$
- c.  $P(.25 < X \le .5) = F(.5) F(.25) = .0107 [10(.25)^9 9(.25)^{10}] = .0107 .0000 = .0107.$ Since X is continuous,  $P(.25 \le X \le .5) = P(.25 < X \le .5) = .0107.$
- **d.** The 75<sup>th</sup> percentile is the value of x for which F(x) = .75:  $10x^9 9x^{10} = .75 \Rightarrow x = .9036$  using software.

e. 
$$E(X) = \int_{-\infty}^{\infty} x \cdot f(x) dx = \int_{0}^{1} x \cdot 90x^{8}(1-x) dx = \int_{0}^{1} (90x^{9} - 90x^{10}) dx = 9x^{10} - \frac{90}{11}x^{11} \Big]_{0}^{1} = 9 - \frac{90}{11} = \frac{9}{11} = .8182.$$

Similarly,  $E(X^2) = \int_{-\infty}^{\infty} x^2 \cdot f(x) dx = \int_{0}^{1} x^2 \cdot 90x^8 (1-x) dx = \dots = .6818$ , from which  $V(X) = .6818 - (.8182)^2 = .0124$  and  $\sigma_X = .11134$ .

f.  $\mu \pm \sigma = (.7068, .9295)$ . Thus,  $P(\mu - \sigma \le X \le \mu + \sigma) = F(.9295) - F(.7068) = .8465 - .1602 = .6863$ , and the probability X is more than 1 standard deviation from its mean value equals 1 - .6863 = 3137.

17.

**a.** To find the (100*p*)th percentile, set F(x) = p and solve for *x*:  $\frac{x-A}{B-A} = p \Rightarrow x = A + (B-A)p.$ 

b. 
$$E(X) = \int_{A}^{B} x \cdot \frac{1}{B-A} dx = \frac{A+B}{2}$$
, the midpoint of the interval. Also,  
 $E(X^{2}) = \frac{A^{2} + AB + B^{2}}{3}$ , from which  $V(X) = E(X^{2}) - [E(X)]^{2} = \dots = \frac{(B-A)^{2}}{12}$ . Finally,  
 $\sigma_{X} = \sqrt{V(X)} = \frac{B-A}{\sqrt{12}}$ .

c. 
$$E(X^n) = \int_A^B x^n \cdot \frac{1}{B-A} dx = \frac{1}{B-A} \frac{x^{n+1}}{n+1} \bigg]_A^B = \frac{B^{n+1} - A^{n+1}}{(n+1)(B-A)}.$$

19.

2

$$P(X \le 1) = F(1) = .25[1 + \ln(4)] = .597$$

- **b.**  $P(1 \le X \le 3) = F(3) F(1) = .966 .597 = .369.$
- c. For x < 0 or x > 4, the pdf is f(x) = 0 since X is restricted to (0, 4). For 0 < x < 4, take the first derivative of the cdf:

$$F(x) = \frac{x}{4} \left[ 1 + \ln\left(\frac{4}{x}\right) \right] = \frac{1}{4}x + \frac{\ln(4)}{4}x - \frac{1}{4}x\ln(x) \Rightarrow$$
  
$$f(x) = F'(x) = \frac{1}{4} + \frac{\ln(4)}{4} - \frac{1}{4}\ln(x) - \frac{1}{4}x\frac{1}{x} = \frac{\ln(4)}{4} - \frac{1}{4}\ln(x) = .3466 - .25\ln(x)$$

21. 
$$E(\text{area}) = E(\pi R^2) = \int_{-\infty}^{\infty} \pi r^2 f(r) dr = \int_{9}^{11} \pi r^2 \frac{3}{4} (1 - (10 - r)^2) dr = \dots = \frac{501}{5} \pi = 314.79 \text{ m}^2.$$

23. With X = temperature in °C, the temperature in °F equals 1.8X + 32, so the mean and standard deviation in °F are  $1.8\mu_X + 32 = 1.8(120) + 32 = 248$ °F and  $|1.8|\sigma_X = 1.8(2) = 3.6$ °F. Notice that the additive constant, 32, affects the mean but does <u>not</u> affect the standard deviation.

25.

- **a.**  $P(Y \le 1.8 \,\tilde{\mu} + 32) = P(1.8X + 32 \le 1.8 \,\tilde{\mu} + 32) = P(X \le \tilde{\mu}) = .5$  since  $\tilde{\mu}$  is the median of X. This shows that  $1.8 \,\tilde{\mu} + 32$  is the median of Y.
- **b.** The 90<sup>th</sup> percentile for Y equals  $1.8\eta(.9) + 32$ , where  $\eta(.9)$  is the 90<sup>th</sup> percentile for X. To see this,  $P(Y \le 1.8\eta(.9) + 32) = P(1.8X + 32 \le 1.8\eta(.9) + 32) = P(X \le \eta(.9)) = .9$ , since  $\eta(.9)$  is the 90<sup>th</sup> percentile of X. This shows that  $1.8\eta(.9) + 32$  is the 90<sup>th</sup> percentile of Y.
- c. When Y = aX + b (i.e. a linear transformation of X) and the (100*p*)th percentile of the X distribution is  $\eta(p)$ , then the corresponding (100*p*)th percentile of the Y distribution is  $a \cdot \eta(p) + b$ . This can be demonstrated using the same technique as in **a** and **b** above.
- 27. Since X is uniform on [0, 360],  $E(X) = \frac{0+360}{2} = 180^{\circ}$  and  $\sigma_X = \frac{360-0}{\sqrt{12}} = 103.82^{\circ}$ . Using the suggested linear representation of Y,  $E(Y) = (2\pi/360)\mu_X \pi = (2\pi/360)(180) \pi = 0$  radians, and  $\sigma_Y = (2\pi/360)\sigma_X = 1.814$  radians. (In fact, Y is uniform on  $[-\pi, \pi]$ .)

## Section 4.3

29.

- a. .9838 is found in the 2.1 row and the .04 column of the standard normal table so c = 2.14.
- **b.**  $P(0 \le Z \le c) = .291 \Rightarrow \Phi(c) \Phi(0) = .2910 \Rightarrow \Phi(c) .5 = .2910 \Rightarrow \Phi(c) = .7910 \Rightarrow$  from the standard normal table, c = .81.
- c.  $P(c \le Z) = .121 \implies 1 P(Z \le c) = .121 \implies 1 \Phi(c) = .121 \implies \Phi(c) = .879 \implies c = 1.17.$
- **d.**  $P(-c \le Z \le c) = \Phi(c) \Phi(-c) = \Phi(c) (1 \Phi(c)) = 2\Phi(c) 1 = .668 \implies \Phi(c) = .834 \implies c = 0.97.$
- e.  $P(c \le |Z|) = 1 P(|Z| \le c) = 1 [\Phi(c) \Phi(-c)] = 1 [2\Phi(c) 1] = 2 2\Phi(c) = .016 \implies \Phi(c) = .992 \implies c = 2.41.$
- 31. By definition,  $z_a$  satisfies  $\alpha = P(Z \ge z_a) = 1 P(Z < z_a) = 1 \Phi(z_a)$ , or  $\Phi(z_a) = 1 \alpha$ . a.  $\Phi(z_{.0055}) = 1 - .0055 = .9945 \Rightarrow z_{.0055} = 2.54$ .
  - **b.**  $\Phi(z_{.09}) = .91 \implies z_{.09} \approx 1.34.$
  - c.  $\Phi(z_{.663}) = .337 \implies z_{.633} \approx -.42$ .
- 33.
- a.  $P(X \le 50) = P\left(Z \le \frac{50 46.8}{1.75}\right) = P(Z \le 1.83) = \Phi(1.83) = .9664.$ b.  $P(X \ge 48) = P\left(Z \ge \frac{48 - 46.8}{1.75}\right) = P(Z \ge 0.69) = 1 - \Phi(0.69) = 1 - .7549 = .2451.$
- c. The mean and standard deviation aren't important here. The probability a normal random variable is within 1.5 standard deviations of its mean equals  $P(-1.5 \le Z \le 1.5) = \Phi(1.5) \Phi(-1.5) = .9332 .0668 = .8664$ .

35.

- a.  $P(X \ge 10) = P(Z \ge .43) = 1 \Phi(.43) = 1 .6664 = .3336$ . Since X is continuous,  $P(X > 10) = P(X \ge 10) = .3336$ .
- **b.**  $P(X > 20) = P(Z > 4) \approx 0.$
- c.  $P(5 \le X \le 10) = P(-1.36 \le Z \le .43) = \Phi(.43) \Phi(-1.36) = .6664 .0869 = .5795.$
- d. P(8.8 c ≤ X ≤ 8.8 + c) = .98, so 8.8 c and 8.8 + c are at the 1<sup>st</sup> and the 99<sup>th</sup> percentile of the given distribution, respectively. The 99<sup>th</sup> percentile of the standard normal distribution satisfies Φ(z) = .99, which corresponds to z = 2.33.
   So, 8.8 + c = μ + 2.33 σ = 8.8 + 2.33(2.8) ⇒ c = 2.33(2.8) = 6.524.
- e. From a, P(X > 10) = .3336, so  $P(X \le 10) = 1 .3336 = .6664$ . For four independent selections,  $P(\text{at least one diameter exceeds } 10) = 1 - P(\text{none of the four exceeds } 10) = 1 - P(\text{first doesn't} \cap ... \text{ fourth doesn't}) = 1 - (.6664)(.6664)(.6664)(.6664) \text{ by independence} = 1 - (.6664)^4 = .8028.$

37.

- P(X = 105) = 0, since the normal distribution is continuous;  $P(X < 105) = P(Z < 0.2) = P(Z \le 0.2) = \Phi(0.2) = .5793$ ;  $P(X \le 105) = .5793$  as well, since X is continuous.
- **b.** No, the answer does not depend on  $\mu$  or  $\sigma$ . For any normal rv,  $P(|X \mu| > \sigma) = P(|Z| > 1) = P(Z < -1 \text{ or } Z > 1) = 2P(Z < -1)$  by symmetry  $= 2\Phi(-1) = 2(.1587) = .3174$ .
- c. From the table,  $\Phi(z) = .1\% = .001 \Rightarrow z = -3.09 \Rightarrow x = 104 3.09(5) = 88.55$  mmol/L. The smallest .1% of chloride concentration values are those less than 88.55 mmol/L

### **39.** $\mu = 30 \text{ mm}, \sigma = 7.8 \text{ mm}$

- a.  $P(X \le 20) = P(Z \le -1.28) = .1003$ . Since X is continuous,  $P(X \le 20) = .1003$  as well.
- **b.** Set  $\Phi(z) = .75$  to find  $z \approx 0.67$ . That is, 0.67 is roughly the 75<sup>th</sup> percentile of a <u>standard</u> normal distribution. Thus, the 75<sup>th</sup> percentile of X's distribution is  $\mu + 0.67\sigma = 30 + 0.67(7.8) = 35.226$  mm.
- c. Similarly,  $\Phi(z) = .15 \Rightarrow z \approx -1.04 \Rightarrow \eta(.15) = 30 1.04(7.8) = 21.888$  mm.
- **d.** The values in question are the 10<sup>th</sup> and 90<sup>th</sup> percentiles of the distribution (in order to have 80% in the middle). Mimicking **b** and **c**,  $\Phi(z) = .1 \Rightarrow z \approx -1.28 \& \Phi(z) = .9 \Rightarrow z \approx +1.28$ , so the 10<sup>th</sup> and 90<sup>th</sup> percentiles are  $30 \pm 1.28(7.8) = 20.016$  mm and 39.984 mm.
- 41. For a single drop,  $P(\text{damage}) = P(X < 100) = P\left(Z < \frac{100 200}{30}\right) = P(Z < -3.33) = .0004$ . So, the

probability of <u>no</u> damage on any single drop is 1 - .0004 = .9996, and  $P(\text{at least one among five is damaged}) = 1 - P(\text{none damaged}) = 1 - (.9996)^5 = 1 - .998 = .002.$ 

43.

**a.** Let  $\mu$  and  $\sigma$  denote the unknown mean and standard deviation. The given information provides  $.05 = P(X < 39.12) = \Phi\left(\frac{39.12 - \mu}{\sigma}\right) \Rightarrow \frac{39.12 - \mu}{\sigma} \approx -1.645 \Rightarrow 39.12 - \mu = -1.645\sigma$  and  $.10 = P(X > 73.24) = 1 - \Phi\left(\frac{73.24 - \mu}{\sigma}\right) \Rightarrow \frac{73.24 - \mu}{\sigma} = \Phi^{-1}(.9) \approx 1.28 \Rightarrow 73.24 - \mu = 1.28\sigma$ .

Subtract the top equation from the bottom one to get  $34.12 = 2.925\sigma$ , or  $\sigma \approx 11.665$  mph. Then, substitute back into either equation to get  $\mu \approx 58.309$  mph.

- **b.**  $P(50 \le X \le 65) = \Phi(.57) \Phi(-.72) = .7157 .2358 = .4799.$
- c.  $P(X > 70) = 1 \Phi(1.00) = 1 .8413 = .1587.$
- With  $\mu = .500$  inches, the acceptable range for the diameter is between .496 and .504 inches, so unacceptable bearings will have diameters smaller than .496 or larger than .504. The new distribution has  $\mu = .499$  and  $\sigma = .002$ .

$$P(X < .496 \text{ or } X > .504) = P\left(Z < \frac{.496 - .499}{.002}\right) + P\left(Z > \frac{.504 - .499}{.002}\right) = P(Z < -1.5) + P(Z > 2.5) =$$

 $\Phi(-1.5) + [1 - \Phi(2.5)] = .073.7.3\%$  of the bearings will be unacceptable.

- 47. The stated condition implies that 99% of the area under the normal curve with  $\mu = 12$  and  $\sigma = 3.5$  is to the left of c 1, so c 1 is the 99<sup>th</sup> percentile of the distribution. Since the 99<sup>th</sup> percentile of the standard normal distribution is z = 2.33,  $c 1 = \mu + 2.33\sigma = 20.155$ , and c = 21.155.
- 49.

45.

a. 
$$P(X > 4000) = P\left(Z > \frac{4000 - 3432}{482}\right) = P(Z > 1.18) = 1 - \Phi(1.18) = 1 - .8810 = .1190;$$
  
 $P(3000 < X < 4000) = P\left(\frac{3000 - 3432}{482} < Z < \frac{4000 - 3432}{482}\right) = \Phi(1.18) - \Phi(-.90) = .8810 - .1841 = .6969.$ 

- **b.**  $P(X < 2000 \text{ or } X > 5000) = P\left(Z < \frac{2000 3432}{482}\right) + P\left(Z > \frac{5000 3432}{482}\right)$ =  $\Phi(-2.97) + [1 - \Phi(3.25)] = .0015 + .0006 = .0021$ .
- c. We will use the conversion 1 lb = 454 g, then 7 lbs = 3178 grams, and we wish to find  $P(X > 3178) = P\left(Z > \frac{3178 3432}{482}\right) = 1 \Phi(-.53) = .7019.$
- d. We need the top .0005 and the bottom .0005 of the distribution. Using the z table, both .9995 and .0005 have multiple z values, so we will use a middle value, ±3.295. Then 3432 ± 3.295(482) = 1844 and 5020. The most extreme .1% of all birth weights are less than 1844 g and more than 5020 g.
- e. Converting to pounds yields a mean of 7.5595 lbs and a standard deviation of 1.0608 lbs. Then

 $P(X > 7) = P\left(Z > \frac{7 - 7.5595}{1.0608}\right) = 1 - \Phi(-.53) = .7019$ . This yields the same answer as in part c.

51.  $P(|X - \mu| \ge \sigma) = 1 - P(|X - \mu| < \sigma) = 1 - P(\mu - \sigma < X < \mu + \sigma) = 1 - P(-1 \le Z \le 1) = .3174.$ Similarly,  $P(|X - \mu| \ge 2\sigma) = 1 - P(-2 \le Z \le 2) = .0456$  and  $P(|X - \mu| \ge 3\sigma) = .0026.$ These are considerably less than the bounds 1, .25, and .11 given by Chebyshev.

a.

Chapter 4: Continuous Random Variables and Probability Distributions

53.  $p = .5 \Rightarrow \mu = 12.5 \& \sigma^2 = 6.25; p = .6 \Rightarrow \mu = 15 \& \sigma^2 = 6; p = .8 \Rightarrow \mu = 20 \text{ and } \sigma^2 = 4$ . These mean and standard deviation values are used for the normal calculations below.

For the	binomial calculat	ion, $P(15 \le X \le 20) = B(20; 25, p) - B(14; 25, p)$ .	
р	$P(15 \le X \le 20)$	$P(14.5 \le \text{Normal} \le 20.5)$	
.5	= .212	$= P(.80 \le Z \le 3.20) = .2112$	
.6	= .577	$= P(20 \le Z \le 2.24) = .5668$	
.8	= .573	$= P(-2.75 \le Z \le .25) = .5957$	

**b.** For the binomial calculation,  $P(X \le 15) = B(15; 25, p)$ .

p	$P(X \le 15)$	$P(\text{Normal} \le 15.5)$
.5	= .885	$= P(Z \le 1.20) = .8849$
.6	= .575	$= P(Z \le .20) = .5793$
.8	= .017	$= P(Z \le -2.25) = .0122$

c. For the binomial calculation,  $P(X \ge 20) = 1 - B(19; 25, p)$ .

 $\begin{array}{c|cccc} p & P(X \ge 20) & P(\text{Normal} \ge 19.5) \\ \hline .5 & = .002 & = P(Z \ge 2.80) = .0026 \\ .6 & = .029 & = P(Z \ge 1.84) = .0329 \\ .8 & = .617 & = P(Z \ge -0.25) = .5987 \end{array}$ 

- 55. Use the normal approximation to the binomial, with a continuity correction. With p = .75 and n = 500,  $\mu = np = 375$ , and  $\sigma = 9.68$ . So, Bin(500, .75)  $\approx N(375, 9.68)$ .
  - a.  $P(360 \le X \le 400) = P(359.5 \le X \le 400.5) = P(-1.60 \le Z \le 2.58) = \Phi(2.58) \Phi(-1.60) = .9409.$

$$P(X < 400) = P(X \le 399.5) = P(Z \le 2.53) = \Phi(2.53) = .9943$$

57.

sist et mi

**a.** For any 
$$a > 0$$
,  $F_Y(y) = P(Y \le y) = P(aX + b \le y) = P\left(X \le \frac{y-b}{a}\right) = F_X\left(\frac{y-b}{a}\right)$ . This, in turn, implies  
 $f_Y(y) = \frac{d}{dy}F_Y(y) = \frac{d}{dy}F_X\left(\frac{y-b}{a}\right) = \frac{1}{a}f_X\left(\frac{y-b}{a}\right)$ .

Now let X have a normal distribution. Applying this rule,

$$f_{\gamma}(y) = \frac{1}{a} \frac{1}{\sqrt{2\pi\sigma}} \exp\left(-\frac{\left((y-b)/a-\mu\right)^2}{2\sigma^2}\right) = \frac{1}{\sqrt{2\pi}a\sigma} \exp\left(-\frac{\left(y-b-a\mu\right)^2}{2a^2\sigma^2}\right).$$
 This is the pdf of a normal

distribution. In particular, from the exponent we can read that the mean of Y is  $E(Y) = a\mu + b$  and the variance of Y is  $V(Y) = a^2 \sigma^2$ . These match the usual rescaling formulas for mean and variance. (The same result holds when a < 0.)

**b.** Temperature in °F would also be normal, with a mean of 1.8(115) + 32 = 239°F and a variance of  $1.8^22^2 = 12.96$  (i.e., a standard deviation of 3.6°F).

www.konkur.in

Chapter 4: Continuous Random Variables and Probability Distributions

# Section 4.4

59.

**a.**  $E(X) = \frac{1}{\lambda} = 1$ .

**b.**  $\sigma = \frac{1}{\lambda} = 1$ .

- c.  $P(X \le 4) = 1 e^{-(1)(4)} = 1 e^{-4} = .982$ .
- **d.**  $P(2 \le X \le 5) = (1 e^{-(1)(5)}) (1 e^{-(1)(2)}) = e^{-2} e^{-5} = .129$ .

Note that a mean value of 2.725 for the exponential distribution implies  $\lambda = \frac{1}{2.725}$ . Let X denote the 61.

- **a.**  $P(X \ge 2) = 1 P(X \le 2) = 1 P(X \le 2) = 1 F(2; \lambda) = 1 [1 e^{-(1/2.725)(2)}] = e^{-2/2.725} = .4800;$  $P(X \le 3) = F(3; \lambda) = 1 e^{-(1/2.725)(3)} = .6674; P(2 \le X \le 3) = .6674 .4800 = .1874.$
- **b.** For this exponential distribution,  $\sigma = \mu = 2.725$ , so  $P(X > \mu + 2\sigma) = P(X > 2.725 + 2(2.725)) = P(X > 8.175) = 1 F(8.175; \lambda) = e^{-(1/2.725)(8.175)} = e^{-3} = .0498.$ On the other hand,  $P(X < \mu - \sigma) = P(X < 2.725 - 2.725) = P(X < 0) = 0$ , since an exponential random variable is non-negative.

63.

- a. If a customer's calls are typically short, the first calling plan makes more sense. If a customer's calls are somewhat longer, then the second plan makes more sense, viz. 99¢ is less than 20min(10¢/min) = \$2 for the first 20 minutes under the first (flat-rate) plan.
- **b.**  $h_1(X) = 10X$ , while  $h_2(X) = 99$  for  $X \le 20$  and 99 + 10(X 20) for X > 20. With  $\mu = 1/\lambda$  for the exponential distribution, it's obvious that  $E[h_1(X)] = 10E[X] = 10\mu$ . On the other hand,

 $E[h_2(X)] = 99 + 10 \int_{20}^{\infty} (x - 20)\lambda e^{-\lambda x} dx = 99 + \frac{10}{\lambda} e^{-20\lambda} = 99 + 10\mu e^{-20/\mu}.$ When  $\mu = 10$ ,  $E[h_1(X)] = 100 \neq \$1.00$  while  $E[h_2(X)] = 99 + 100e^{-2} \approx \$1.13$ . When  $\mu = 15$ ,  $E[h_1(X)] = 150 \notin = \$1.50$  while  $E[h_2(X)] = 99 + 150 e^{-4/3} \approx \$1.39$ . As predicted, the first plan is better when expected call length is lower, and the second plan is better when expected call length is somewhat higher.

- 65.
- From the mean and sd equations for the gamma distribution,  $\alpha\beta = 37.5$  and  $\alpha\beta^2 = (21.6)2 = 466.56$ . Take the quotient to get  $\beta = 466.56/37.5 = 12.4416$ . Then,  $\alpha = 37.5/\beta = 37.5/12.4416 = 3.01408...$ a.
- **b.**  $P(X > 50) = 1 P(X \le 50) = 1 F(50/12.4416; 3.014) = 1 F(4.0187; 3.014)$ . If we approximate this by 1 - F(4; 3), Table A.4 gives 1 - .762 = .238. Software gives the more precise answer of .237.
- $P(50 \le X \le 75) = F(75/12.4416; 3.014) F(50/12.4416; 3.014) = F(6.026; 3.014) F(4.0187; 3.014) = F(50/12.4416; 3.0416; 3.0416; 3.0416; 3.0416; 3.0416; 3.04$ c. F(6; 3) - F(4; 3) = .938 - .762 = .176.
- Notice that  $\mu = 24$  and  $\sigma^2 = 144 \implies \alpha\beta = 24$  and  $\alpha\beta^2 = 144 \implies \beta = \frac{144}{24} = 6$  and  $\alpha = \frac{24}{\beta} = 4$ . 67.
  - $P(12 \le X \le 24) = F(4; 4) F(2; 4) = .424.$
  - $P(X \le 24) = F(4; 4) = .567$ , so while the mean is 24, the median is less than 24, since  $P(X \le \tilde{\mu}) = .5$ . b. This is a result of the positive skew of the gamma distribution.
  - We want a value x for which  $F\left(\frac{x}{\beta},\alpha\right) = F\left(\frac{x}{6},4\right) = .99$ . In Table A.4, we see F(10; 4) = .990. So x/6 =c. 10, and the 99<sup>th</sup> percentile is 6(10) = 60.
  - We want a value t for which P(X > t) = .005, i.e.  $P(X \le t) = .005$ . The left-hand side is the cdf of X, so d. we really want  $F\left(\frac{t}{6}, 4\right) = .995$ . In Table A.4, F(11; 4) = .995, so t/6 = 11 and t = 6(11) = 66. At 66 weeks, only .5% of all transistors would still be operating.
  - 69.
- $\{X \ge t\} = \{\text{the lifetime of the system is at least } t\}$ . Since the components are connected in series, this a. equals {all 5 lifetimes are at least t} =  $A_1 \cap A_2 \cap A_3 \cap A_4 \cap A_5$ .
- Since the events  $A_i$  are assumed to be independent,  $P(X \ge t) = P(A_1 \cap A_2 \cap A_3 \cap A_4 \cap A_5) = P(A_1)$ b.  $P(A_2) \cdot P(A_3) \cdot P(A_4) \cdot P(A_5)$ . Using the exponential cdf, for any i we have  $P(A_i) = P(\text{component lifetime is } \geq t) = 1 - F(t) = 1 - [1 - e^{-01t}] = e^{-01t}.$ Therefore,  $P(X \ge t) = (e^{-01t}) \cdots (e^{-01t}) = e^{-.05t}$ , and  $F_X(t) = P(X \le t) = 1 - e^{-.05t}.$ Taking the derivative, the pdf of X is  $f_X(t) = .05e^{-.05t}$  for  $t \ge 0$ . Thus X also has an exponential distribution, but with parameter  $\lambda = .05$ .
- By the same reasoning,  $P(X \le t) = 1 e^{-n\lambda t}$ , so X has an exponential distribution with parameter  $n\lambda$ . c.

71.

a.

$$\{X^2 \leq y\} = \{-\sqrt{y} \leq X \leq \sqrt{y}\}.$$

**b.**  $F_Y(y) = P(Y \le y) = P(X^2 \le y) = P(-\sqrt{y} \le X \le \sqrt{y}) = \int_{-\sqrt{y}}^{\sqrt{y}} \frac{1}{\sqrt{2\pi}} e^{-z^2/2} dz$ . To find the pdf of Y, use the identity (Leibniz's rule):

$$f_{Y}(y) = \frac{1}{\sqrt{2\pi}} e^{-(\sqrt{y})^{2}/2} \cdot \frac{d\sqrt{y}}{dy} - \frac{1}{\sqrt{2\pi}} e^{-(-\sqrt{y})^{2}/2} \cdot \frac{d(-\sqrt{y})}{dy}$$
$$= \frac{1}{\sqrt{2\pi}} e^{-y/2} \cdot \frac{1}{2\sqrt{y}} - \frac{1}{\sqrt{2\pi}} e^{-y/2} \cdot \frac{-1}{2\sqrt{y}} = \frac{1}{\sqrt{2\pi}} y^{-1/2} e^{-y/2}$$

This is valid for y > 0. We recognize this as the chi-squared pdf with y = 1.

### Section 4.5

73.

**a.**  $P(X \le 250) = F(250; 2.5, 200) = 1 - e^{-(250/200)^{23}} = 1 - e^{-1.75} = .8257.$  $P(X < 250) = P(X \le 250) = .8257.$  $P(X > 300) = 1 - F(300; 2.5, 200) = e^{-(1.5)^{25}} = .0636.$ 

- **b.**  $P(100 \le X \le 250) = F(250; 2.5, 200) F(100; 2.5, 200) = .8257 .162 = .6637.$
- The question is asking for the median,  $\tilde{\mu}$ . Solve  $F(\tilde{\mu}) = .5: .5 = 1 e^{-(\tilde{\mu}/200)^{2.5}} \Rightarrow$ c.  $e^{-(\tilde{\mu}/200)^{25}} = .5 \Rightarrow (\tilde{\mu}/200)^{2.5} = -\ln(.5) \Rightarrow \tilde{\mu} = 200(-\ln(.5))^{1/2.5} = 172.727$  hours.
- Using the substitution  $y = \left(\frac{x}{\beta}\right)^{\alpha} = \frac{x^{\alpha}}{\beta^{\alpha}}$ . Then  $dy = \frac{\alpha x^{\alpha-1}}{\beta^{\alpha}} dx$ , and  $\mu = \int_{0}^{\infty} x \cdot \frac{\alpha}{\beta^{\alpha}} x^{\alpha-1} e^{-\left(\frac{x}{\beta}\right)^{\alpha}} dx = \int_{0}^{\infty} x^{\alpha-1} e^{-\left(\frac{x}{\beta}\right)^{\alpha}} dx$ 75.  $\int_{0}^{\infty} (\beta^{\alpha} y)^{1/\alpha} \cdot e^{-y} dy = \beta \int_{0}^{\infty} y^{1/\alpha} e^{-y} dy = \beta \cdot \Gamma\left(1 + \frac{1}{\alpha}\right)$  by definition of the gamma function.
- 77.
- **a.**  $E(X) = e^{\mu + \sigma^2/2} = e^{4.82} = 123.97.$  $V(X) = \left(e^{2(4.5)+.8^2}\right) \cdot \left(e^{-.8}-1\right) = 13,776.53 \implies \sigma = 117.373.$

$$P(X \le 100) = \Phi\left(\frac{\ln(100) - 4.5}{.8}\right) = \Phi(0.13) = .5517.$$
  

$$P(X \ge 200) = 1 - P(X < 200) = 1 - \Phi\left(\frac{\ln(200) - 4.5}{.8}\right) = 1 - \Phi(1.00) = 1 - .8413 = .1587.$$
 Since X is continuous,  

$$P(X > 200) = .1587 \text{ as well.}$$

79.

Notice that  $\mu_X$  and  $\sigma_X$  are the mean and standard deviation of the lognormal variable X in this example; they are not the parameters  $\mu$  and  $\sigma$  which usually refer to the mean and standard deviation of ln(X). We're given  $\mu_X = 10,281$  and  $\sigma_X/\mu_X = .40$ , from which  $\sigma_X = .40\mu_X = 4112.4$ .

To find the mean and standard deviation of ln(X), set the lognormal mean and variance equal to the appropriate quantities:  $10,281 = E(X) = e^{\mu + \sigma^2/2}$  and  $(4112.4)^2 = V(X) = e^{2\mu + \sigma^2} (e^{\sigma^2} - 1)$ . Square the first a.

equation: 
$$(10.281)^2 = e^{2\mu+\sigma^2}$$
. Now divide the variance by this amount:

$$\frac{(4112.4)^2}{(10,281)^2} = \frac{e^{2\mu+\sigma^2}(e^{\sigma^2}-1)}{e^{2\mu+\sigma^2}} \Rightarrow e^{\sigma^2}-1 = (.40)^2 = .16 \Rightarrow \sigma = \sqrt{\ln(1.16)} = .38525$$

That's the standard deviation of ln(X). Use this in the formula for E(X) to solve for  $\mu$ :  $10,281 = e^{\mu + (.38525)^2/2} = e^{\mu + .07421} \Longrightarrow \mu = 9.164$ . That's  $E(\ln(X))$ .

**b.** 
$$P(X \le 15,000) = P\left(Z \le \frac{\ln(15,000) - 9.164}{.38525}\right) = P(Z \le 1.17) = \Phi(1.17) = .8790.$$

c. 
$$P(X \ge \mu_X) = P(X \ge 10,281) = P\left(Z \ge \frac{\ln(10,281) - 9.164}{.38525}\right) = P(Z \ge .19) = 1 - \Phi(0.19) = .4247$$
. Even

though the normal distribution is symmetric, the lognormal distribution is <u>not</u> a symmetric distribution. (See the lognormal graphs in the textbook.) So, the mean and the median of X aren't the same and, in particular, the probability X exceeds its own mean doesn't equal .5.

**d.** One way to check is to determine whether P(X < 17,000) = .95; this would mean 17,000 is indeed the 95<sup>th</sup> percentile. However, we find that  $P(X < 17,000) = \Phi\left(\frac{\ln(17,000) - 9.164}{.38525}\right) = \Phi(1.50) = .9332$ , so 17,000 is <u>not</u> the 95<sup>th</sup> percentile of this distribution (it's the 93.32%ile).

81.

**a.**  $V(X) = e^{2(2.05)+.06}(e^{.06}-1) = 3.96 \Rightarrow SD(X) = 1.99$  months.

**b.** 
$$P(X > 12) = 1 - P(X \le 12) = 1 - P\left(Z \le \frac{\ln(12) - 2.05}{\sqrt{.06}}\right) = 1 - \Phi(1.78) = .0375.$$

- c. The mean of X is  $E(X) = e^{2.05 + .06/2} = 8.00$  months, so  $P(\mu_X \sigma_X < X < \mu_X + \sigma_X) = P(6.01 < X < 9.99) = \Phi\left(\frac{\ln(9.99) 2.05}{\sqrt{.06}}\right) \Phi\left(\frac{\ln(6.01) 2.05}{\sqrt{.06}}\right) = \Phi(1.03) \Phi(-1.05) = .8485 .1469 = .7016.$
- **d.**  $.5 = F(x) = \Phi\left(\frac{\ln(x) 2.05}{\sqrt{.06}}\right) \Rightarrow \frac{\ln(x) 2.05}{\sqrt{.06}} = \Phi^{-1}(.5) = 0 \Rightarrow \ln(x) 2.05 = 0 \Rightarrow \text{the median is given}$ by  $x = e^{2.05} = 7.77$  months.
- e. Similarly,  $\frac{\ln(\eta_{99}) 2.05}{\sqrt{.06}} = \Phi^{-1}(.99) = 2.33 \Rightarrow \eta_{.99} = e^{2.62} = 13.75$  months.
- **f.** The probability of exceeding 8 months is  $P(X > 8) = 1 \Phi\left(\frac{\ln(8) 2.05}{\sqrt{.06}}\right) = 1 \Phi(.12) = .4522$ , so the expected number that will exceed 8 months out of n = 10 is just 10(.4522) = 4.522.
- 83. Since the standard beta distribution lies on (0, 1), the point of symmetry must be  $\frac{1}{2}$ , so we require that  $f(\frac{1}{2}-\mu) = f(\frac{1}{2}+\mu)$ . Cancelling out the constants, this implies

 $\left(\frac{1}{2}-\mu\right)^{\alpha-1}\left(\frac{1}{2}+\mu\right)^{\beta-1} = \left(\frac{1}{2}+\mu\right)^{\alpha-1}\left(\frac{1}{2}-\mu\right)^{\beta-1}$ , which (by matching exponents on both sides) in turn implies that  $\alpha = \beta$ .

Alternatively, symmetry about  $\frac{1}{2}$  requires  $\mu = \frac{1}{2}$ , so  $\frac{\alpha}{\alpha + \beta} = .5$ . Solving for  $\alpha$  gives  $\alpha = \beta$ .

85.

8.

Notice from the definition of the standard beta pdf that, since a pdf must integrate to 1,  $\int_{1}^{1} \Gamma(\alpha + \beta) x^{\alpha-1} (1-x)^{\beta-1} dx \rightarrow \int_{1}^{1} x^{\alpha-1} (1-x)^{\beta-1} dx = \frac{\Gamma(\alpha)\Gamma(\beta)}{1-x^{\alpha-1}}$ 

$$1 = \int_{0}^{1} \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} x^{\alpha + 1} (1 - x)^{\alpha} dx \Rightarrow \int_{0}^{1} x \cdot (1 - x)^{\alpha} dx - \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha + \beta)}$$
Using this,  $E(X) = \int_{0}^{1} x \cdot \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} x^{\alpha - 1} (1 - x)^{\beta - 1} dx = \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \int_{0}^{1} x^{\alpha} (1 - x)^{\beta - 1} dx = \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \cdot \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} = \frac{\alpha\Gamma(\alpha)}{\Gamma(\alpha)\Gamma(\beta)} \cdot \frac{\Gamma(\alpha + \beta)}{(\alpha + \beta)\Gamma(\alpha + \beta)} = \frac{\alpha}{\alpha + \beta}.$ 

**b.** Similarly,  $E[(1-X)^m] = \int_0^1 (1-x)^m \cdot \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)} x^{\alpha-1} (1-x)^{\beta-1} dx =$  $\Gamma(\alpha+\beta) = \Gamma(\alpha+\beta) \Gamma(\alpha)\Gamma(m+\beta) - \frac{\Gamma(\alpha+\beta)\cdot\Gamma(m+\beta)}{\Gamma(\alpha+\beta)} = \Gamma(\alpha+\beta)\cdot\Gamma(m+\beta) - \Gamma(\alpha+\beta)\cdot\Gamma(m+\beta) = \Gamma(\alpha+\beta)\cdot\Gamma(m+\beta)$ 

$$=\frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)}\int_{0}^{1}x^{\alpha-1}(1-x)^{m+\beta-1}dx=\frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)}\frac{\Gamma(\alpha)\Gamma(m+\beta)}{\Gamma(\alpha+m+\beta)}=\frac{\Gamma(\alpha+\beta)\Gamma(m+\beta)}{\Gamma(\alpha+m+\beta)\Gamma(\beta)}$$

If X represents the proportion of a substance consisting of an ingredient, then 1 - X represents the proportion <u>not</u> consisting of this ingredient. For m = 1 above,

$$E(1-X) = \frac{\Gamma(\alpha+\beta)\cdot\Gamma(1+\beta)}{\Gamma(\alpha+1+\beta)\Gamma(\beta)} = \frac{\Gamma(\alpha+\beta)\cdot\beta\Gamma(\beta)}{(\alpha+\beta)\Gamma(\alpha+\beta)\Gamma(\beta)} = \frac{\beta}{\alpha+\beta}$$

# Section 4.6

E

- 87. The given probability plot is quite linear, and thus it is quite plausible that the tension distribution is normal.
- **89.** The plot below shows the (observation, *z* percentile) pairs provided. Yes, we would feel comfortable using a normal probability model for this variable, because the normal probability plot exhibits a strong, linear pattern.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

**91.** The (*z* percentile, observation) pairs are (-1.66, .736), (-1.32, .863), (-1.01, .865), (-.78, .913), (-.58, .915), (-.40, .937), (-.24, .983), (-.08, 1.007), (.08, 1.011), (.24, 1.064), (.40, 1.109), (.58, 1.132), (.78, 1.140), (1.01, 1.153), (1.32, 1.253), (1.86, 1.394).

The accompanying probability plot is straight, suggesting that an assumption of population normality is plausible.



**93.** To check for plausibility of a lognormal population distribution for the rainfall data of Exercise 81 in Chapter 1, take the natural logs and construct a normal probability plot. This plot and a normal probability plot for the original data appear below. Clearly the log transformation gives quite a straight plot, so lognormality is plausible. The curvature in the plot for the original data implies a positively skewed population distribution — like the lognormal distribution.



Beennie A is a scale peramone (is in other the normal inicity), A = 1 circl to med to access the planarbilly of the units express that its airy if we used a different value of A to find the percentiles, the alope of the graph would charge, but not its linearly (or task thereofy

71

The pattern in the plot (below, generated by Minitab) is reasonably linear. By visual inspection alone, it is plausible that strength is normally distributed.



97. The  $(100p)^{\text{th}}$  percentile  $\eta(p)$  for the exponential distribution with  $\lambda = 1$  is given by the formula  $\eta(p) = -\ln(1-p)$ . With n = 16, we need  $\eta(p)$  for  $p = \frac{0.5}{16}, \frac{1.5}{16}, \dots, \frac{15.5}{16}$ . These are .032, .398, .170, .247, .330, .421, .521, .633, .758, .901, 1.068, 1.269, 1.520, 1.856, 2.367, 3.466.

The accompanying plot of (percentile, failure time value) pairs exhibits substantial curvature, casting doubt on the assumption of an exponential population distribution.





# 5 34

95.

# Supplementary Exercises

99.

**a.** For 
$$0 \le y \le 25$$
,  $F(y) = \frac{1}{24} \int_0^y \left( u - \frac{u^2}{12} \right) du = \frac{1}{24} \left( \frac{u^2}{2} - \frac{u^3}{36} \right) \Big]_0^y = \frac{y^2}{48} - \frac{y^3}{864}$ . Thus  

$$F(y) = \begin{cases} 0 & y < 0 \\ \frac{y^2}{48} - \frac{y^3}{864} & 0 \le y \le 12 \\ 1 & y > 12 \end{cases}$$

**b.**  $P(Y \le 4) = F(4) = .259$ . P(Y > 6) = 1 - F(6) = .5.  $P(4 \le X \le 6) = F(6) - F(4) = .5 - .259 = .241$ .

c. 
$$E(Y) = \int_0^{12} y \frac{1}{24} y \left(1 - \frac{y}{12}\right) dy = \frac{1}{24} \int_0^{12} \left(y^2 - \frac{y^3}{12}\right) dy = \frac{1}{24} \left[\frac{y^3}{3} - \frac{y^4}{48}\right]_0^{12} = 6$$
 inches.  
 $E(Y^2) = \frac{1}{24} \int_0^{12} \left(y^3 - \frac{y^4}{12}\right) dy = 43.2$ , so  $V(Y) = 43.2 - 36 = 7.2$ .

**d.** 
$$P(Y < 4 \text{ or } Y > 8) = 1 - P(4 \le Y \le 8) = 1 - [F(8) - F(4)] = .518.$$

e. The shorter segment has length equal to 
$$\min(Y, 12 - Y)$$
, and  

$$E[\min(Y, 12 - Y)] = \int_0^{12} \min(y, 12 - y) \cdot f(y) dy = \int_0^6 \min(y, 12 - y) \cdot f(y) dy$$

$$+ \int_6^{12} \min(y, 12 - y) \cdot f(y) dy = \int_0^6 y \cdot f(y) dy + \int_6^{12} (12 - y) \cdot f(y) dy = \frac{90}{24} = 3.75 \text{ inches.}$$

101.

**a.** By differentiation, 
$$f(x) = \begin{cases} x^2 & 0 \le x < 1\\ \frac{7}{4} - \frac{3}{4}x & 1 \le x < \frac{7}{3} \end{cases}$$



Using the pdf from **a**, 
$$E(X) = \int_0^1 x \cdot x^2 dx + \int_1^{\frac{1}{2}} x \cdot \left(\frac{7}{4} - \frac{3}{4}x\right) dx = \frac{131}{108} = 1.213.$$

103.

6

**a.** 
$$P(X > 135) = 1 - \Phi\left(\frac{135 - 137.2}{1.6}\right) = 1 - \Phi(-1.38) = 1 - .0838 = .9162.$$

- **b.** With Y = the number among ten that contain more than 135 oz,  $Y \sim Bin(10, .9162)$ . So,  $P(Y \ge 8) = b(8; 10, .9162) + b(9; 10, .9162) + b(10; 10, .9162) = .9549$ .
- c. We want P(X > 135) = .95, i.e.  $1 \Phi\left(\frac{135 137.2}{\sigma}\right) = .95$  or  $\Phi\left(\frac{135 137.2}{\sigma}\right) = .05$ . From the standard normal table,  $\frac{135 137.2}{\sigma} = -1.65 \Rightarrow \sigma = 1.33$ .
- 105. Let A = the cork is acceptable and B = the first machine was used. The goal is to find P(B | A), which can be obtained from Bayes' rule:

$$P(B \mid A) = \frac{P(B)P(A \mid B)}{P(B)P(A \mid B) + P(B')P(A \mid B')} = \frac{.6P(A \mid B)}{.6P(A \mid B) + .4P(A \mid B')} \cdot$$

From Exercise 38, P(A | B) = P(machine 1 produces an acceptable cork) = .6826 and P(A | B') = P(machine 2 produces an acceptable cork) = .9987. Therefore,

$$P(B \mid A) = \frac{.6(.6826)}{.6(.6826) + .4(.9987)} = .5062.$$

107.



**b.** F(x) = 0 for x < -1, and F(x) = 1 for x > 2. For  $-1 \le x \le 2$ ,  $F(x) = \int_{-1}^{x} \frac{1}{9} (4 - y^2) dy = \frac{11 + 12x - x^3}{27}$ . This is





© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

www.konkur.in

677

- c. The median is 0 iff F(0) = .5. Since  $F(0) = \frac{11}{27}$ , this is not the case. Because  $\frac{11}{27} < .5$ , the median must be greater than 0. (Looking at the pdf in a it's clear that the line x = 0 does not evenly divide the distribution, and that such a line must lie to the right of x = 0.)
- **d.** *Y* is a binomial rv, with n = 10 and  $p = P(X > 1) = 1 F(1) = \frac{5}{27}$ .
- 109. Below, exp(u) is alternative notation for  $e^{u}$ .

ß

**a.** 
$$P(X \le 150) = \exp\left[-\exp\left(\frac{-(150-150)}{90}\right)\right] = \exp[-\exp(0)] = \exp(-1) = .368$$
,  
 $P(X \le 300) = \exp[-\exp(-1.6667)] = .828$ , and  $P(150 \le X \le 300) = .828 - .368 = .460$ .

- **b.** The desired value c is the 90<sup>th</sup> percentile, so c satisfies  $.9 = \exp\left[-\exp\left(\frac{-(c-150)}{90}\right)\right].$  Taking the natural log of each side twice in succession yields  $\frac{-(c-150)}{90}$  $= \ln[-\ln(.9)] = -2.250367,$  so c = 90(2.250367) + 150 = 352.53.
- c. Use the chain rule:  $f(x) = F'(x) = \exp\left[-\exp\left(\frac{-(x-\alpha)}{\beta}\right)\right] \exp\left(\frac{-(x-\alpha)}{\beta}\right) \frac{1}{\beta} = \frac{1}{\beta} \exp\left[-\exp\left(\frac{-(x-\alpha)}{\beta}\right) \frac{(x-\alpha)}{\beta}\right].$
- **d.** We wish the value of x for which f(x) is a maximum; from calculus, this is the same as the value of x for which  $\ln[f(x)]$  is a maximum, and  $\ln[f(x)] = -\ln \beta e^{-(x-\alpha)/\beta} \frac{(x-\alpha)}{\beta}$ . The derivative of  $\ln[f(x)]$  is

$$\frac{d}{dx}\left[-\ln\beta - e^{-(x-\alpha)/\beta} - \frac{(x-\alpha)}{\beta}\right] = 0 + \frac{1}{\beta}e^{-(x-\alpha)/\beta} - \frac{1}{\beta}$$
; set this equal to 0 and we get  $e^{-(x-\alpha)/\beta} = 1$ , so  $-(x-\alpha) = 0$ , which implies that  $x = \alpha$ . Thus the mode is  $\alpha$ .

e.  $E(X) = .5772\beta + \alpha = 201.95$ , whereas the mode is  $\alpha = 150$  and the median is the solution to F(x) = .5. From b, this equals  $-90\ln[-\ln(.5)] + 150 = 182.99$ . Since mode < median < mean, the distribution is positively skewed. A plot of the pdf appears below.



75

111.

- a. From a graph of the normal pdf or by differentiation of the pdf,  $x^* = \mu$ .
- **b.** No; the density function has constant height for  $A \le x \le B$ .
- c.  $f(x; \lambda)$  is largest for x = 0 (the derivative at 0 does not exist since f is not continuous there), so  $x^* = 0$ .
- **d.**  $\ln[f(x;\alpha,\beta)] = -\ln(\beta^{\alpha}) \ln(\Gamma(\alpha)) + (\alpha 1)\ln(x) \frac{x}{\beta}$ , and  $\frac{d}{dx}\ln[f(x;\alpha,\beta)] = \frac{\alpha 1}{x} \frac{1}{\beta}$ . Setting this equal to 0 gives the mode:  $x^* = (\alpha 1)\beta$ .
- e. The chi-squared distribution is the gamma distribution with  $\alpha = v/2$  and  $\beta = 2$ . From d,  $x^* = \left(\frac{v}{2} - 1\right)(2) = v - 2$ .

113.

**a.**  $E(X) = \int_0^\infty x \cdot \left[ p \lambda_1 e^{-\lambda_1 x} + (1-p) \lambda_2 e^{-\lambda_2 x} \right] dx = p \int_0^\infty x \lambda_1 e^{-\lambda_1 x} dx + (1-p) \int_0^\infty x \lambda_2 e^{-\lambda_2 x} dx = \frac{p}{\lambda_1} + \frac{(1-p)}{\lambda_2} \cdot (\text{Each of } x + (1-p) \lambda_2 e^{-\lambda_2 x} dx)$ 

the two integrals represents the expected value of an exponential random variable, which is the reciprocal of  $\lambda$ .) Similarly, since the mean-square value of an exponential rv is  $E(Y^2) = V(Y) + [E(Y)]^2 = 2n - 2(1-n)$ 

$$\frac{1}{\lambda^2} + \frac{1}{\lambda^2} = \frac{2}{\lambda^2}, E(X^2) = \int_0^\infty x^2 f(x) dx = \dots = \frac{2p}{\lambda_1^2} + \frac{2(1-p)}{\lambda_2^2}.$$
 From this

$$V(X) = \frac{2p}{\lambda_1^2} + \frac{2(1-p)}{\lambda_2^2} - \left[\frac{p}{\lambda_1} + \frac{(1-p)}{\lambda_2}\right].$$

- **b.** For x > 0,  $F(x; \lambda_1, \lambda_2, p) = \int_0^x f(y; \lambda_1, \lambda_2, p) dy = \int_0^x \left[ p \lambda_1 e^{-\lambda_1 y} + (1-p) \lambda_2 e^{-\lambda_2 y} \right] dy = p \int_0^x \lambda_1 e^{-\lambda_2 y} dy + (1-p) \int_0^x \lambda_2 e^{-\lambda_2 y} dy = p(1-e^{-\lambda_1 x}) + (1-p)(1-e^{-\lambda_2 x})$ . For  $x \le 0$ , F(x) = 0.
- c.  $P(X > .01) = 1 F(.01) = 1 [.5(1 e^{-40(.01)}) + (1 .5)(1 e^{-200(.01)})] = .5e^{-0.4} + .5e^{-2} = .403.$
- **d.** Using the expressions in **a**,  $\mu = .015$  and  $\sigma^2 = .000425 \Rightarrow \sigma = .0206$ . Hence,  $P(\mu - \sigma < X < \mu + \sigma) = P(-.0056 < X < .0356) = P(X < .0356)$  because X can't be negative = F(.0356) = ... = .879.
- e. For an exponential rv,  $CV = \frac{\sigma}{\mu} = \frac{1/\lambda}{1/\lambda} = 1$ . For X hyperexponential,

 $CV = \frac{\sigma}{\mu} = \frac{\sqrt{E(X^2) - \mu^2}}{\mu} = \sqrt{\frac{E(X^2)}{\mu^2} - 1} = \sqrt{\frac{2p/\lambda_1^2 + 2(1-p)/\lambda_2^2}{[p/\lambda_1 + (1-p)/\lambda_2]^2} - 1} = \sqrt{\frac{2(p\lambda_2^2 + (1-p)\lambda_1^2)}{(p\lambda_2 + (1-p)\lambda_1)^2} - 1} = \sqrt{2r - 1}, \text{ where } r = \frac{p\lambda_2^2 + (1-p)\lambda_1^2}{(p\lambda_2 + (1-p)\lambda_1)^2}.$  But straightforward algebra shows that r > 1 when  $\lambda_1 \neq \lambda_2$ , so that CV > 1.

**f.** For the Erlang distribution,  $\mu = \frac{n}{\lambda}$  and  $\sigma = \frac{\sqrt{n}}{\lambda}$ , so  $CV = \frac{1}{\sqrt{n}} < 1$  for n > 1.

www.konkur.in

115.

## Chapter 4: Continuous Random Variables and Probability Distributions

**a.** Since  $\ln\left(\frac{I_o}{I_i}\right)$  has a normal distribution, by definition  $\frac{I_o}{I_i}$  has a lognormal distribution.

**b.** 
$$P(I_o > 2I_i) = P\left(\frac{I_o}{I_i} > 2\right) = P\left(\ln\left(\frac{I_o}{I_i}\right) > \ln 2\right) = 1 - P\left(\ln\left(\frac{I_o}{I_i}\right) \le \ln 2\right) = 1 - P(X \le \ln 2) = 1 - \Phi\left(\frac{\ln 2 - 1}{05}\right) = 1 - \Phi\left(-6.14\right) = 1.$$

c. 
$$E\left(\frac{I_o}{I_i}\right) = e^{1+.0025/2} = 2.72$$
 and  $V\left(\frac{I_o}{I_i}\right) = e^{2+.0025} \cdot \left(e^{.0025} - 1\right) = .0185$ 

117. 
$$F(y) = P(Y \le y) = P(\sigma Z + \mu \le y) = P\left(Z \le \frac{y - \mu}{\sigma}\right) = \int_{-\infty}^{\frac{y - \mu}{\sigma}} \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}z^2} dz \implies \text{by the fundamental theorem of calculus, } f(y) = F'(y) = \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{y - \mu}{\sigma}\right)^2} \cdot \frac{1}{\sigma} = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{1}{2}\left(\frac{y - \mu}{\sigma}\right)^2}, \text{ a normal pdf with parameters } \mu \text{ and } \sigma.$$

119.

- **a.**  $Y = -\ln(X) \Rightarrow x = e^{-y} = k(y)$ , so  $k'(y) = -e^{-y}$ . Thus since f(x) = 1,  $g(y) = 1 \cdot |-e^{-y}| = e^{-y}$  for  $0 < y < \infty$ . *Y* has an exponential distribution with parameter  $\lambda = 1$ .
- **b.**  $y = \sigma Z + \mu \Rightarrow z = k(y) = \frac{y \mu}{\sigma}$  and  $k'(y) = \frac{1}{\sigma}$ , from which the result follows easily.

c.  $y = h(x) = cx \Rightarrow x = k(y) = \frac{y}{c}$  and  $k'(y) = \frac{1}{c}$ , from which the result follows easily.

121.

**a.** Assuming the three birthdays are independent and that all 365 days of the calendar year are equally likely,  $P(\text{all 3 births occur on March 11}) = \left(\frac{1}{365}\right)^3$ .

**b.**  $P(\text{all 3 births on the same day}) = P(\text{all 3 on Jan. 1}) + P(\text{all 3 on Jan. 2}) + ... = \left(\frac{1}{365}\right)^3 + \left(\frac{1}{365}\right)^3 + ... = 365\left(\frac{1}{365}\right)^3 = \left(\frac{1}{365}\right)^2$ .

c. Let X = deviation from due date, so  $X \sim N(0, 19.88)$ . The baby due on March 15 was 4 days early, and  $P(X = -4) \approx P(-4.5 < X < -3.5) = \Phi\left(\frac{-3.5}{19.88}\right) - \Phi\left(\frac{-4.5}{19.88}\right) = \Phi(-.18) - \Phi(-.237) = .4286 - .4090 = .0196$ . Similarly, the baby due on April 1 was 21 days early, and  $P(X = -21) \approx \Phi\left(\frac{-20.5}{19.88}\right) - \Phi\left(\frac{-21.5}{19.88}\right) = \Phi(-1.03) - \Phi(-1.08) = .1515 - .1401 = .0114$ .

Finally, the baby due on April 4 was 24 days early, and  $P(X = -24) \approx .0097$ .

Again assuming independence, P(all 3 births occurred on March 11) = (.0196)(.0114)(.0097) = .0002145.

**d.** To calculate the probability of the three births happening on any day, we could make similar calculations as in part **c** for each possible day, and then add the probabilities.

123.

**a.**  $F(x) = P(X \le x) = P\left(-\frac{1}{\lambda}\ln(1-U) \le x\right) = P\left(\ln(1-U) \ge -\lambda x\right) = P\left(1-U \ge e^{-\lambda x}\right)$ 

 $= P(U \le 1 - e^{-\lambda x}) = 1 - e^{-\lambda x}$  since the cdf of a uniform rv on [0, 1] is simply F(u) = u. Thus X has an exponential distribution with parameter  $\lambda$ .

- **b.** By taking successive random numbers  $u_1, u_2, u_3, ...$  and computing  $x_i = -\frac{1}{10} \ln(1-u_i)$  for each one, we obtain a sequence of values generated from an exponential distribution with parameter  $\lambda = 10$ .
- 125. If g(x) is convex in a neighborhood of  $\mu$ , then  $g(\mu) + g'(\mu)(x \mu) \le g(x)$ . Replace x by X:  $E[g(\mu) + g'(\mu)(X - \mu)] \le E[g(X)] \Rightarrow E[g(X)] \ge g(\mu) + g'(\mu)E[(X - \mu)] = g(\mu) + g'(\mu) \cdot 0 = g(\mu).$ That is, if g(x) is convex,  $g(E(X)) \le E[g(X)]$ .
- 127.
- **a.**  $E(X) = 150 + (850 150)\frac{8}{8+2} = 710 \text{ and } V(X) = \frac{(850 150)^2(8)(2)}{(8+2)^2(8+2+1)} = 7127.27 \Rightarrow SD(X) \approx 84.423.$ Using software,  $P(|X - 710| \le 84.423) = P(625.577 \le X \le 794.423) = \int_{625.577}^{794.423} \frac{1}{700} \frac{\Gamma(10)}{\Gamma(8)\Gamma(2)} \left(\frac{x - 150}{700}\right)^7 \left(\frac{850 - x}{700}\right)^1 dx = .684.$
- **b.**  $P(X > 750) = \int_{750}^{850} \frac{1}{700} \frac{\Gamma(10)}{\Gamma(8)\Gamma(2)} \left(\frac{x-150}{700}\right)^7 \left(\frac{850-x}{700}\right)^1 dx = .376$ . Again, the computation of the requested integral requires a calculator or computer.

Name? In about 19 and an about 19 (1) which we don't hand and Sample

# **CHAPTER 5**

Section 5.1

1.

- **a.** P(X = 1, Y = 1) = p(1,1) = .20.
  - **b.**  $P(X \le 1 \text{ and } Y \le 1) = p(0,0) + p(0,1) + p(1,0) + p(1,1) = .42.$
  - c. At least one hose is in use at both islands.  $P(X \neq 0 \text{ and } Y \neq 0) = p(1,1) + p(1,2) + p(2,1) + p(2,2) = .70$ .
  - **d.** By summing row probabilities,  $p_X(x) = .16, .34, .50$  for x = 0, 1, 2, By summing column probabilities,  $p_X(y) = .24, .38, .38$  for y = 0, 1, 2.  $P(X \le 1) = p_X(0) + p_X(1) = .50$ .
  - e. p(0,0) = .10, but  $p_X(0) \cdot p_Y(0) = (.16)(.24) = .0384 \neq .10$ , so X and Y are not independent.
- 3.
- **a.** p(1,1) = .15, the entry in the 1<sup>st</sup> row and 1<sup>st</sup> column of the joint probability table.
- **b.**  $P(X_1 = X_2) = p(0,0) + p(1,1) + p(2,2) + p(3,3) = .08 + .15 + .10 + .07 = .40.$
- c.  $A = \{X_1 \ge 2 + X_2 \cup X_2 \ge 2 + X_1\}$ , so P(A) = p(2,0) + p(3,0) + p(4,0) + p(3,1) + p(4,1) + p(4,2) + p(0,2) + p(0,3) + p(1,3) = .22.
- **d.**  $P(X_1 + X_2 = 4) = p(1,3) + p(2,2) + p(3,1) + p(4,0) = .17.$  $P(X_1 + X_2 \ge 4) = P(X_1 + X_2 = 4) + p(4,1) + p(4,2) + p(4,3) + p(3,2) + p(3,3) + p(2,3) = .46.$
- a. p(3, 3) = P(X = 3, Y = 3) = P(3 customers, each with 1 package)=  $P(\text{ each has 1 package } | 3 \text{ customers}) \cdot P(3 \text{ customers}) = (.6)^3 \cdot (.25) = .054.$
- **b.**  $p(4, 11) = P(X=4, Y=11) = P(\text{total of 11 packages} | 4 \text{ customers}) \cdot P(4 \text{ customers}).$ Given that there are 4 customers, there are four different ways to have a total of 11 packages: 3, 3, 3, 2 or 3, 3, 2, 3 or 3, 2, 3, 3 or 2, 3, 3, 3. Each way has probability  $(.1)^3(.3)$ , so  $p(4, 11) = 4(.1)^3(.3)(.15) = .00018$ .
- 7.

5.

- **a.** p(1,1) = .030.
- **b.**  $P(X \le 1 \text{ and } Y \le 1) = p(0,0) + p(0,1) + p(1,0) + p(1,1) = .120.$
- c. P(X=1) = p(1,0) + p(1,1) + p(1,2) = .100; P(Y=1) = p(0,1) + ... + p(5,1) = .300.
- **d.**  $P(\text{overflow}) = P(X + 3Y > 5) = 1 P(X + 3Y \le 5) = 1 P((X,Y) = (0,0) \text{ or } ...\text{ or } (5,0) \text{ or } (0,1) \text{ or } (1,1) \text{ or } (2,1)) = 1 .620 = .380.$

- e. The marginal probabilities for X (row sums from the joint probability table) are  $p_X(0) = .05$ ,  $p_X(1) = .10$ ,  $p_X(2) = .25$ ,  $p_X(3) = .30$ ,  $p_X(4) = .20$ ,  $p_X(5) = .10$ ; those for Y (column sums) are  $p_Y(0) = .5$ ,  $p_Y(1) = .3$ ,  $p_Y(2) = .2$ . It is now easily verified that for every (x,y),  $p(x,y) = p_X(x) \cdot p_Y(y)$ , so X and Y are independent.
- 9.

1. 
$$1 = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y) dx dy = \int_{20}^{30} \int_{20}^{30} K(x^2 + y^2) dx dy = K \int_{20}^{30} \int_{20}^{30} x^2 dy dx + K \int_{20}^{30} \int_{20}^{30} y^2 dx dy$$
$$= 10K \int_{20}^{30} x^2 dx + 10K \int_{20}^{30} y^2 dy = 20K \cdot \left(\frac{19,000}{3}\right) \Longrightarrow K = \frac{3}{380,000}.$$

**b.**  $P(X < 26 \text{ and } Y < 26) = \int_{20}^{26} \int_{20}^{26} K(x^2 + y^2) dx dy = K \int_{20}^{26} \left[ x^2 y + \frac{y^3}{3} \right]_{20}^{26} dx = K \int_{20}^{26} (6x^2 + 3192) dx = K \int_{20}^{26} (6x^2 + 3192) dx$ 

K(38,304) = .3024.

c. The region of integration is labeled III below.



$$P(|X-Y| \le 2) = \iint_{III} f(x,y) dx dy = 1 - \iint_{I} f(x,y) dx dy - \iint_{II} f(x,y) dx dy = 1 - \int_{20}^{28} \int_{x+2}^{30} f(x,y) dy dx - \int_{22}^{30} \int_{20}^{x-2} f(x,y) dy dx = .3593 \text{ (after much algebra)}.$$

- **d.**  $f_x(x) = \int_{-\infty}^{\infty} f(x, y) dy = \int_{20}^{30} K(x^2 + y^2) dy = 10Kx^2 + K \frac{y^3}{3} \Big|_{20}^{30} = 10Kx^2 + .05, \text{ for } 20 \le x \le 30.$
- e.  $f_Y(y)$  can be obtained by substituting y for x in (d); clearly  $f(x,y) \neq f_X(x) \cdot f_Y(y)$ , so X and Y are not independent.
- 11.

**a.** Since X and Y are independent,  $p(x,y) = p_X(x) \cdot p_Y(y) = \frac{e^{-\mu_1} \mu_1^x}{x!} \cdot \frac{e^{-\mu_2} \mu_2^y}{y!} = \frac{e^{-\mu_1 - \mu_2} \mu_1^x \mu_2^y}{x! y!}$ for x = 0, 1, 2, ...; y = 0, 1, 2, ...

**b.**  $P(X+Y \le 1) = p(0,0) + p(0,1) + p(1,0) = \dots = e^{-\mu_1 - \mu_2} [1 + \mu_1 + \mu_2].$ 

c. 
$$P(X + Y = m) = \sum_{k=0}^{m} P(X = k, Y = m - k) = e^{-\mu_1 - \mu_2} \sum_{k=0}^{m} \frac{\mu_1^k}{k!} \frac{\mu_2^{m-k}}{(m-k)!} = \frac{e^{-\mu_1 - \mu_2}}{m!} \sum_{k=0}^{m} \frac{m!}{k!(m-k)!} \mu_1^k \mu_2^{m-k} = \frac{e^{-\mu_1 - \mu_2}}{m!} \sum_{k=0}^{m} \binom{m}{k!} \mu_1^{k} \mu_2^{m-k} = \frac{e^{-\mu_1 - \mu_2}}{m!} (\mu_1 + \mu_2)^m$$
 by the binomial theorem. We recognize this as the pmf of

www.konkur.in

F

Poisson random variable with parameter  $\mu_1 + \mu_2$ . Therefore, the total number of errors, X + Y, also has a Poisson distribution, with parameter  $\mu_1 + \mu_2$ .

13.

a.

$$f(x,y) = f_{X}(x) \cdot f_{Y}(y) = \begin{cases} e^{-x-y} & x \ge 0, y \ge 0\\ 0 & \text{otherwise} \end{cases}$$

**b.** By independence,  $P(X \le 1 \text{ and } Y \le 1) = P(X \le 1) \cdot P(Y \le 1) = (1 - e^{-1})(1 - e^{-1}) = .400.$ 

c. 
$$P(X + Y \le 2) = \int_0^2 \int_0^{2-x} e^{-x-y} dy dx = \int_0^2 e^{-x} \left[1 - e^{-(2-x)}\right] dx = \int_0^2 (e^{-x} - e^{-2}) dx = 1 - e^{-2} - 2e^{-2} = .594.$$

**d.** 
$$P(X + Y \le 1) = \int_0^1 e^{-x} \left[ 1 - e^{-(1-x)} \right] dx = 1 - 2e^{-1} = .264$$
,  
so  $P(1 \le X + Y \le 2) = P(X + Y \le 2) - P(X + Y \le 1) = .594 - .264 = .330$ 

15.

**a.** Each  $X_i$  has  $\operatorname{cdf} F(x) = P(X_i \le x) = 1 - e^{-\lambda x}$ . Using this, the cdf of Y is  $F(y) = P(Y \le y) = P(X_1 \le y \cup [X_2 \le y \cap X_3 \le y])$ =  $P(X_1 \le y) + P(X_2 \le y \cap X_3 \le y) - P(X_1 \le y \cap [X_2 \le y \cap X_3 \le y])$ =  $(1 - e^{-\lambda y}) + (1 - e^{-\lambda y})^2 - (1 - e^{-\lambda y})^3$  for y > 0.

The pdf of Y is  $f(y) = F'(y) = \lambda e^{-\lambda y} + 2(1 - e^{-\lambda y})(\lambda e^{-\lambda y}) - 3(1 - e^{-\lambda y})^2(\lambda e^{-\lambda y}) = 4\lambda e^{-2\lambda y} - 3\lambda e^{-3\lambda y}$ for y > 0.

**b.** 
$$E(Y) = \int_0^\infty y \cdot \left(4\lambda e^{-2\lambda y} - 3\lambda e^{-3\lambda y}\right) dy = 2\left(\frac{1}{2\lambda}\right) - \frac{1}{3\lambda} = \frac{2}{3\lambda}$$

17.

**a.** Let A denote the disk of radius R/2. Then  $P((X, Y) \text{ lies in } A) = \iint_{X \to Y} f(x, y) dx dy$ 

$$= \iint_{A} \frac{1}{\pi R^2} dx dy = \frac{1}{\pi R^2} \iint_{A} dx dy = \frac{\text{area of } A}{\pi R^2} = \frac{\pi (R/2)^2}{\pi R^2} = \frac{1}{4} = .25$$
. Notice that, since the joint pdf of X

and Y is a constant (i.e., (X, Y) is <u>uniform</u> over the disk), it will be the case for any subset A that P((X, Y)lies in  $A) = \frac{\text{area of } A}{\pi R^2}$ .

**b.** By the same ratio-of-areas idea,  $P\left(-\frac{R}{2} \le X \le \frac{R}{2}, -\frac{R}{2} \le Y \le \frac{R}{2}\right) = \frac{R^2}{\pi R^2} = \frac{1}{\pi}$ . This region is the square depicted in the graph below.



c. Similarly,  $P\left(-\frac{R}{\sqrt{2}} \le X \le \frac{R}{\sqrt{2}}, -\frac{R}{\sqrt{2}} \le Y \le \frac{R}{\sqrt{2}}\right) = \frac{2R^2}{\pi R^2} = \frac{2}{\pi}$ . This region is the slightly larger square depicted in the graph below, whose corners actually touch the circle.



**d.** 
$$f_{X}(x) = \int_{-\infty}^{\infty} f(x, y) dy = \int_{-\sqrt{R^{2} - x^{2}}}^{\sqrt{R^{2} - x^{2}}} \frac{1}{\pi R^{2}} dy = \frac{2\sqrt{R^{2} - x^{2}}}{\pi R^{2}} \text{ for } -R \le x \le R.$$

Similarly,  $f_Y(y) = \frac{2\sqrt{R^2 - y^2}}{\pi R^2}$  for  $-R \le y \le R$ . X and Y are <u>not</u> independent, since the joint pdf is not

the product of the marginal pdfs: 
$$\frac{1}{\pi R^2} \neq \frac{2\sqrt{R^2 - x^2}}{\pi R^2} \cdot \frac{2\sqrt{R^2 - y^2}}{\pi R^2}$$

19. Throughout these solutions, 
$$K = \frac{3}{380,000}$$
, as calculated in Exercise 9.

**a.** 
$$f_{Y|X}(y \mid x) = \frac{f(x, y)}{f_X(x)} = \frac{K(x^2 + y^2)}{10Kx^2 + .05}$$
 for  $20 \le y \le 30$ .  
 $f_{X|Y}(x \mid y) = \frac{f(x, y)}{f_Y(y)} = \frac{K(x^2 + y^2)}{10Ky^2 + .05}$  for  $20 \le x \le 30$ .

**b.**  $P(Y \ge 25 \mid X = 22) = \int_{25}^{30} f_{Y|X}(y \mid 22) dy = \int_{25}^{30} \frac{K((22)^2 + y^2)}{10K(22)^2 + .05} dy = .5559.$ 

 $P(Y \ge 25) = \int_{25}^{30} f_Y(y) dy = \int_{25}^{30} (10Ky^2 + .05) dy = .75$ . So, given that the right tire pressure is 22 psi, it's much less likely that the left tire pressure is at least 25 psi.

c.  $E(Y|X=22) = \int_{-\infty}^{\infty} y \cdot f_{Y|X}(y|22) dy = \int_{20}^{30} y \cdot \frac{K((22)^2 + y^2)}{10K(22)^2 + .05} dy = 25.373 \text{ psi.}$   $E(Y^2|X=22) = \int_{20}^{30} y^2 \cdot \frac{k((22)^2 + y^2)}{10k(22)^2 + .05} dy = 652.03 \Rightarrow$   $V(Y|X=22) = E(Y^2|X=22) - [E(Y|X=22)]^2 = 652.03 - (25.373)^2 = 8.24 \Rightarrow$ SD(Y|X=22) = 2.87 psi.

21.

**a.** 
$$f_{X_3|X_1,X_2}(x_3 \mid x_1, x_2) = \frac{f(x_1, x_2, x_3)}{f_{X_1,X_2}(x_1, x_2)}$$
, where  $f_{X_1,X_2}(x_1, x_2) =$  the marginal joint pdf of  $X_1$  and  $X_2$ , i.e.  $f_{X_1,X_2}(x_1, x_2) = \int_{-\infty}^{\infty} f(x_1, x_2, x_3) dx_3$ .

**b.** 
$$f_{X_2,X_3|X_1}(x_2,x_3|X_1) = \frac{f(x_1,x_2,x_3)}{f_{X_1}(x_1)}$$
, where  $f_{X_1}(x_1) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x_1,x_2,x_3) dx_2 dx_3$ , the marginal pdf of  $X_1$ .

## Section 5.2

23. 
$$E(X_1 - X_2) = \sum_{x_1=0}^{4} \sum_{x_2=0}^{3} (x_1 - x_2) \cdot p(x_1, x_2) = (0 - 0)(.08) + (0 - 1)(.07) + \dots + (4 - 3)(.06) = .15.$$

Note: It can be shown that  $E(X_1 - X_2)$  always equals  $E(X_1) - E(X_2)$ , so in this case we could also work out the means of  $X_1$  and  $X_2$  from their marginal distributions:  $E(X_1) = 1.70$  and  $E(X_2) = 1.55$ , so  $E(X_1 - X_2) = E(X_1) - E(X_2) = 1.70 - 1.55 = .15$ .

- 25. The expected value of X, being uniform on [L A, L + A], is simply the midpoint of the interval, L. Since Y has the same distribution, E(Y) = L as well. Finally, since X and Y are independent,  $E(\text{area}) = E(XY) = E(X) \cdot E(Y) = L \cdot L = L^2$ .
- 27. The amount of time Annie waits for Alvie, if Annie arrives first, is Y X; similarly, the time Alvie waits for Annie is X Y. Either way, the amount of time the first person waits for the second person is h(X, Y) = |X Y|. Since X and Y are independent, their joint pdf is given by  $f_X(x) \cdot f_Y(y) = (3x^2)(2y) = 6x^2y$ . From these, the expected waiting time is  $E[h(X, Y)] = \int_{0}^{1} \int_{0}^{1} |x y| \cdot f(x y) dx dy = \int_{0}^{1} \int_{0}^{1} |x y| \cdot f(x y) dx dy = \int_{0}^{1} \int_{0}^{1} |x y| dx dy$

$$E[h(X,Y)] = \int_{0}^{1} \int_{0}^{1} |x-y| \cdot f(x,y) dx dy = \int_{0}^{1} \int_{0}^{1} |x-y| \cdot 6x^{2}y dx dy$$
  
=  $\int_{0}^{1} \int_{0}^{x} (x-y) \cdot 6x^{2}y dy dx + \int_{0}^{1} \int_{x}^{1} (x-y) \cdot 6x^{2}y dy dx = \frac{1}{6} + \frac{1}{12} = \frac{1}{4}$  hour, or 15 minutes.

29.

$$Cov(X,Y) = -\frac{1}{75} \text{ and } \mu_X = \mu_Y = \frac{1}{5}.$$
  

$$E(X^2) = \int_0^1 x^2 \cdot f_X(x) dx = 12 \int_0^1 x^3 (1 - x^2 dx) = \frac{12}{60} = \frac{1}{5}, \text{ so } V(X) = \frac{1}{5} - \left(\frac{2}{5}\right)^2 = \frac{1}{25}.$$
  
Similarly,  $V(Y) = \frac{1}{5}$ , so  $\rho_{X,Y} = \frac{-\frac{2}{75}}{5} = -\frac{50}{54} = -\frac{2}{5}.$ 

V25 V25

2

31.

$$E(X) = \int_{20}^{30} x f_X(x) dx = \int_{20}^{30} x \Big[ 10Kx^2 + .05 \Big] dx = \frac{1925}{76} = 25.329 = E(Y),$$

$$E(XY) = \int_{20}^{30} \int_{20}^{30} xy \cdot K(x^2 + y^2) dx dy = \frac{24375}{38} = 641.447 \Rightarrow$$

$$Cov(X, Y) = 641.447 - (25.329)^2 = -.1082.$$

**b.** 
$$E(X^2) = \int_{20}^{30} x^2 \Big[ 10Kx^2 + .05 \Big] dx = \frac{37040}{57} = 649.8246 = E(Y^2) \Rightarrow$$
  
 $V(X) = V(Y) = 649.8246 - (25.329)^2 = 8.2664 \Rightarrow \rho = \frac{-.1082}{\sqrt{(8.2664)(8.2664)}} = -.0131.$ 

33. Since  $E(XY) = E(X) \cdot E(Y)$ ,  $Cov(X, Y) = E(XY) - E(X) \cdot E(Y) = E(X) \cdot E(Y) - E(X) \cdot E(Y) = 0$ , and since  $Corr(X, Y) = \frac{Cov(X, Y)}{\sigma_x \sigma_y}$ , then Corr(X, Y) = 0.

35.

- **a.**  $\operatorname{Cov}(aX + b, cY + d) = E[(aX + b)(cY + d)] E(aX + b) \cdot E(cY + d)$ = E[acXY + adX + bcY + bd] - (aE(X) + b)(cE(Y) + d)= acE(XY) + adE(X) + bcE(Y) + bd - [acE(X)E(Y) + adE(X) + bcE(Y) + bd]=  $acE(XY) - acE(X)E(Y) = ac[E(XY) - E(X)E(Y)] = ac\operatorname{Cov}(X, Y).$
- **b.**  $\operatorname{Corr}(aX+b, cY+d) = \frac{\operatorname{Cov}(aX+b, cY+d)}{SD(aX+b)SD(cY+d)} = \frac{ac\operatorname{Cov}(X,Y)}{|a| \cdot |c|SD(X)SD(Y)} = \frac{ac}{|ac|}\operatorname{Corr}(X,Y)$ . When a and c have the same signs, ac = |ac|, and we have  $\operatorname{Corr}(aX+b, cY+d) = \operatorname{Corr}(X,Y)$
- c. When a and c differ in sign, |ac| = -ac, and we have Corr(aX + b, cY + d) = -Corr(X, Y).

# Section 5.3

**37.** The joint pmf of  $X_1$  and  $X_2$  is presented below. Each joint probability is calculated using the independence of  $X_1$  and  $X_2$ ; e.g.,  $p(25, 25) = P(X_1 = 25) \cdot P(X_2 = 25) = (.2)(.2) = .04$ .

$p(x_1, x_2)$	25	40	65	
25	.04	.10	.06	
40	.10	.25	.15	
65	.06	.15	.09	
	.2	.5	.3	

**a.** For each coordinate in the table above, calculate  $\overline{x}$ . The six possible resulting  $\overline{x}$  values and their corresponding probabilities appear in the accompanying pmf table.

<b>v</b>	25	32.5	40	45	52.5	65	
$p(\overline{x})$	.04	.20	.25	.12	.30	.09	

From the table,  $E(\overline{X}) = (25)(.04) + 32.5(.20) + ... + 65(.09) = 44.5$ . From the original pmf,  $\mu = 25(.2) + 40(.5) + 65(.3) = 44.5$ . So,  $E(\overline{X}) = \mu$ .

**b.** For each coordinate in the joint pmf table above, calculate  $s^2 = \frac{1}{2-1} \sum_{i=1}^{2} (x_i - \overline{x})^2$ . The four possible

resulting  $s^2$  values and their corresponding probabilities appear in the accompanying pmf table.

e <sup>2</sup>	0	112.5	312.5	800
$p(s^2)$	.38	.20	.30	.12

From the table,  $E(S^2) = 0(.38) + ... + 800(.12) = 212.25$ . From the original pmf,  $\sigma^2 = (25 - 44.5)^2(.2) + (40 - 44.5)^2(.5) + (65 - 44.5)^2(.3) = 212.25$ . So,  $E(S^2) = \sigma^2$ .

39.

X is a binomial random variable with n = 15 and p = .8. The values of X, then X/n = X/15 along with the corresponding probabilities b(x; 15, .8) are displayed in the accompanying pmf table.

x	0	1	2	3	4	5	6	7	8	9	10
x/15	0	.067	.133	.2	.267	.333	.4	.467	.533	.6	.667
p(x/15)	.000	.000	.000	.000	.000	.000	.001	.003	.014	.043	.103
x	11	12	13	14	15	APP 1		E.			
x/15	.733	.8	.867	.933	1						
p(x/15)	.188	.250	.231	.132	.035	1.1					

41. The tables below delineate all 16 possible  $(x_1, x_2)$  pairs, their probabilities, the value of  $\overline{x}$  for that pair, and the value of r for that pair. Probabilities are calculated using the independence of  $X_1$  and  $X_2$ .

$(x_1, x_2)$	1,1	1,2	1,3	1,4	2,1	2,2	2,3	2,4
probability	.16	.12	.08	.04	.12	.09	.06	.03
x	1	1.5	2	2.5	1.5	2	2.5	3
r	0	1	2	3	1	0	1	2
$(x_1, x_2)$	3,1	3,2	3,3	3,4	4,1	4,2	4,3	4,4
probability	.08	.06	.04	.02	.04	.03	.02	.01
x	2	2.5	3	3.5	2.5	3	3.5	4
r	2	1	0	1	3	2	1	2

**a.** Collecting the  $\overline{x}$  values from the table above yields the pmf table below.

$\overline{x}$	1	1.5	2	2.5	3	3.5	4
$p(\overline{x})$	.16	.24	.25	.20	.10	.04	.01

**b.** 
$$P(\overline{X} \le 2.5) = .16 + .24 + .25 + .20 = .85.$$

E discourses an

c. Collecting the r values from the table above yields the pmf table below.

r	0	1	2	3
<i>p</i> ( <i>r</i> )	.30	.40	.22	.08

- **d.** With n = 4, there are numerous ways to get a sample average of at most 1.5, since  $\overline{X} \le 1.5$  iff the sum of the  $X_i$  is at most 6. Listing out all options,  $P(\overline{X} \le 1.5) = P(1,1,1,1) + P(2,1,1,1) + ... + P(1,1,1,2) + P(1,1,2,2) + ... + P(2,2,1,1) + P(3,1,1,1) + ... + P(1,1,1,3) = (.4)^4 + 4(.4)^3(.3) + 6(.4)^2(.3)^2 + 4(.4)^2(.2)^2 = .2400.$
- 43. The statistic of interest is the fourth spread, or the difference between the medians of the upper and lower halves of the data. The population distribution is uniform with A = 8 and B = 10. Use a computer to generate samples of sizes n = 5, 10, 20, and 30 from a uniform distribution with A = 8 and B = 10. Keep the number of replications the same (say 500, for example). For each replication, compute the upper and lower fourth, then compute the difference. Plot the sampling distributions on separate histograms for n = 5, 10, 20, and 30.

45.

Using Minitab to generate the necessary sampling distribution, we can see that as n increases, the distribution slowly moves toward normality. However, even the sampling distribution for n = 50 is not yet approximately normal.



# Section 5.4

**a.** In the previous exercise, we found  $E(\overline{X}) = 70$  and  $SD(\overline{X}) = 0.4$  when n = 16. If the diameter distribution is normal, then  $\overline{X}$  is also normal, so

 $P(69 \le \overline{X} \le 71) = P\left(\frac{69 - 70}{0.4} \le Z \le \frac{71 - 70}{0.4}\right) = P(-2.5 \le Z \le 2.5) = \Phi(2.5) - \Phi(-2.5) = .9938 - .0062 = .9876.$ 

**b.** With 
$$n = 25$$
,  $E(\overline{X}) = 70$  but  $SD(\overline{X}) = \frac{1.6}{\sqrt{25}} = 0.32$  GPa. So,  $P(\overline{X} > 71) = P\left(Z > \frac{71-70}{0.32}\right) = 1 - \Phi(3.125) = 1 - .9991 = .0009.$ 

49.

**a.** 11 P.M. - 6:50 P.M. = 250 minutes. With  $T_o = X_1 + ... + X_{40}$  = total grading time,  $\mu_{T_o} = n\mu = (40)(6) = 240$  and  $\sigma_{T_o} = \sigma \cdot \sqrt{n} = 37.95$ , so  $P(T_o \le 250) \approx$ 

$$P\left(Z \le \frac{250 - 240}{37.95}\right) = P\left(Z \le .26\right) = .6026.$$

www.konkur.in

6

<sup>47.</sup> 

b. The sports report begins 260 minutes after he begins grading papers.

$$P(T_0 > 260) = P(Z > \frac{260 - 240}{37.95}) = P(Z > .53) = .2981.$$

51. Individual times are given by  $X \sim N(10, 2)$ . For day 1, n = 5, and so

$$P(\overline{X} \le 11) = P\left(Z \le \frac{11-10}{2/\sqrt{5}}\right) = P(Z \le 1.12) = .8686$$
.

For day 2, n = 6, and so

$$P(\overline{X} \le 11) = P(\overline{X} \le 11) = P\left(Z \le \frac{11-10}{2/\sqrt{6}}\right) = P(Z \le 1.22) = .8888$$

Finally, assuming the results of the two days are independent (which seems reasonable), the probability the sample average is at most 11 min on both days is (.8686)(.8888) = .7720.

53.

a. With the values provided,

$$P(\overline{X} \ge 51) = P\left(Z \ge \frac{51-50}{1.2/\sqrt{9}}\right) = P(Z \ge 2.5) = 1-.9938 = .0062.$$

**b.** Replace n = 9 by n = 40, and  $P(\overline{X} \ge 51) = P\left(Z \ge \frac{51 - 50}{1.2 / \sqrt{40}}\right) = P(Z \ge 5.27) \approx 0$ .

55.

**a.** With Y = # of tickets, Y has approximately a normal distribution with  $\mu = 50$  and  $\sigma = \sqrt{\mu} = 7.071$ . So, using a continuity correction from [35, 70] to [34.5, 70.5],

$$P(35 \le Y \le 70) \approx P\left(\frac{34.5 - 50}{7.071} \le Z \le \frac{70.5 - 50}{7.071}\right) = P(-2.19 \le Z \le 2.90) = .9838.$$

- **b.** Now  $\mu = 5(50) = 250$ , so  $\sigma = \sqrt{250} = 15.811$ . Using a continuity correction from [225, 275] to [224.5, 275.5],  $P(225 \le Y \le 275) \approx P\left(\frac{224.5 - 250}{15.811} \le Z \le \frac{275.5 - 250}{15.811}\right) = P(-1.61 \le Z \le 1.61) = .8926.$
- **c.** Using software, part (a) =  $\sum_{y=35}^{70} \frac{e^{-50} 50^y}{y!}$  = .9862 and part (b) =  $\sum_{y=225}^{275} \frac{e^{-250} 250^y}{y!}$  = .8934. Both of the approximations in (a) and (b) are correct to 2 decimal places.
- 57. With the parameters provided,  $E(X) = \alpha\beta = 100$  and  $V(X) = \alpha\beta^2 = 200$ . Using a normal approximation,  $P(X \le 125) \approx P\left(Z \le \frac{125 - 100}{\sqrt{200}}\right) = P(Z \le 1.77) = .9616.$

87

# Section 5.5

59.

- **a.**  $E(X_1 + X_2 + X_3) = 180, V(X_1 + X_2 + X_3) = 45, SD(X_1 + X_2 + X_3) = \sqrt{45} = 6.708.$   $P(X_1 + X_2 + X_3 \le 200) = P\left(Z \le \frac{200 - 180}{6.708}\right) = P(Z \le 2.98) = .9986.$  $P(150 \le X_1 + X_2 + X_3 \le 200) = P(-4.47 \le Z \le 2.98) \approx .9986.$
- **b.**  $\mu_{\bar{X}} = \mu = 60$  and  $\sigma_{\bar{X}} = \frac{\sigma_{\bar{X}}}{\sqrt{n}} = \frac{\sqrt{15}}{\sqrt{3}} = 2.236$ , so  $P(\bar{X} \ge 55) = P\left(Z \ge \frac{55 - 60}{2.236}\right) = P(Z \ge -2.236) = .9875$  and  $P(58 \le \bar{X} \le 62) = P(-.89 \le Z \le .89) = .6266.$
- c.  $E(X_1 .5X_2 .5X_3) = \mu .5 \mu .5 \mu = 0$ , while  $V(X_1 - .5X_2 - .5X_3) = \sigma_1^2 + .25\sigma_2^2 + .25\sigma_3^2 = 22.5 \Rightarrow SD(X_1 - .5X_2 - .5X_3) = 4.7434$ . Thus,  $P(-10 \le X_1 - .5X_2 - .5X_3 \le 5) = P\left(\frac{-10 - 0}{4.7434} \le Z \le \frac{5 - 0}{4.7434}\right) = P\left(-2.11 \le Z \le 1.05\right) = .8531 - .0174 = .8357.$
- **d.**  $E(X_1 + X_2 + X_3) = 150, V(X_1 + X_2 + X_3) = 36 \Rightarrow SD(X_1 + X_2 + X_3) = 6$ , so  $P(X_1 + X_2 + X_3 \le 200) = P\left(Z \le \frac{160 150}{6}\right) = P(Z \le 1.67) = .9525.$

Next, we want  $P(X_1 + X_2 \ge 2X_3)$ , or, written another way,  $P(X_1 + X_2 - 2X_3 \ge 0)$ .  $E(X_1 + X_2 - 2X_3) = 40 + 50 - 2(60) = -30$  and  $V(X_1 + X_2 - 2X_3) = \sigma_1^2 + \sigma_2^2 + 4\sigma_3^2 = 78 \Rightarrow$   $SD(X_1 + X_2 - 2X_3) = 8.832$ , so  $P(X_1 + X_2 - 2X_3 \ge 0) = P\left(Z \ge \frac{0 - (-30)}{8.832}\right) = P(Z \ge 3.40) = .0003$ .

61.

- **a.** The marginal pmfs of X and Y are given in the solution to Exercise 7, from which E(X) = 2.8, E(Y) = .7, V(X) = 1.66, and V(Y) = .61. Thus, E(X + Y) = E(X) + E(Y) = 3.5, V(X + Y) = V(X) + V(Y) = 2.27, and the standard deviation of X + Y is 1.51.
- **b.** E(3X + 10Y) = 3E(X) + 10E(Y) = 15.4, V(3X + 10Y) = 9V(X) + 100V(Y) = 75.94, and the standard deviation of revenue is 8.71.

63.

**a.** 
$$E(X_1) = 1.70, E(X_2) = 1.55, E(X_1X_2) = \sum_{x_1} \sum_{x_2} x_1 x_2 p(x_1, x_2) = \dots = 3.33$$
, so  
 $Cov(X_1, X_2) = E(X_1X_2) - E(X_1) E(X_2) = 3.33 - 2.635 = .695.$ 

**b.**  $V(X_1 + X_2) = V(X_1) + V(X_2) + 2Cov(X_1, X_2) = 1.59 + 1.0875 + 2(.695) = 4.0675$ . This is much larger than  $V(X_1) + V(X_2)$ , since the two variables are positively correlated.

65.

**a.** 
$$E(\overline{X} - \overline{Y}) = 0; V(\overline{X} - \overline{Y}) = \frac{\sigma^2}{25} + \frac{\sigma^2}{25} = .0032 \implies \sigma_{\overline{X} - \overline{Y}} = \sqrt{.0032} = .0566$$
  
 $\implies P(-.1 \le \overline{X} - \overline{Y} \le .1) = P(-1.77 \le Z \le 1.77) = .9232.$ 

- **b.**  $V(\overline{X} \overline{Y}) = \frac{\sigma^2}{36} + \frac{\sigma^2}{36} = .0022222 \Rightarrow \sigma_{\overline{X} \overline{Y}} = .0471$  $\Rightarrow P(-.1 \le \overline{X} - \overline{Y} \le .1) \approx P(-2.12 \le Z \le 2.12) = .9660$ . The normal curve calculations are still justified
  - $\Rightarrow P(-.1 \le X Y \le .1) \approx P(-2.12 \le 2 \le 2.12) = .9660$ . The normal curve calculations are still justified here, even though the populations are not normal, by the Central Limit Theorem (36 is a sufficiently "large" sample size).
- 67. Letting  $X_1$ ,  $X_2$ , and  $X_3$  denote the lengths of the three pieces, the total length is  $X_1 + X_2 X_3$ . This has a normal distribution with mean value 20 + 15 1 = 34 and variance .25 + .16 + .01 = .42 from which the standard deviation is .6481. Standardizing gives  $P(34.5 \le X_1 + X_2 X_3 \le 35) = P(.77 \le Z \le 1.54) = .1588$ .
- 69.
- **a.**  $E(X_1 + X_2 + X_3) = 800 + 1000 + 600 = 2400.$
- **b.** Assuming independence of  $X_1, X_2, X_3, V(X_1 + X_2 + X_3) = (16)^2 + (25)^2 + (18)^2 = 1205.$
- c.  $E(X_1 + X_2 + X_3) = 2400$  as before, but now  $V(X_1 + X_2 + X_3)$ =  $V(X_1) + V(X_2) + V(X_3) + 2Cov(X_1, X_2) + 2Cov(X_1, X_3) + 2Cov(X_2, X_3) = 1745$ , from which the standard deviation is 41.77.

#### 71.

**a.**  $M = a_1 X_1 + a_2 X_2 + W \int_0^{12} x dx = a_1 X_1 + a_2 X_2 + 72W$ , so E(M) = (5)(2) + (10)(4) + (72)(1.5) = 158 and  $\sigma_M^2 = (5)^2 (.5)^2 + (10)^2 (1)^2 + (72)^2 (.25)^2 = 430.25 \implies \sigma_M = 20.74.$ 

**b.** 
$$P(M \le 200) = P\left(Z \le \frac{200 - 158}{20.74}\right) = P(Z \le 2.03) = .9788$$

#### 73.

- a. Both are approximately normal by the Central Limit Theorem.
- **b.** The difference of two rvs is just an example of a linear combination, and a linear combination of normal rvs has a normal distribution, so  $\overline{X} \overline{Y}$  has approximately a normal distribution with  $\mu_{\overline{X}-\overline{Y}} = 5$

and 
$$\sigma_{\bar{X}-\bar{Y}} = \sqrt{\frac{8^2}{40} + \frac{6^2}{35}} = 1.621$$
.

c. 
$$P(-1 \le \overline{X} - \overline{Y} \le 1) \approx P\left(\frac{-1-5}{1.6213} \le Z \le \frac{1-5}{1.6213}\right) = P(-3.70 \le Z \le -2.47) \approx .0068$$

89

**d.**  $P(\overline{X} - \overline{Y} \ge 10) \approx P(Z \ge \frac{10-5}{1.6213}) = P(Z \ge 3.08) = .0010$ . This probability is quite small, so such an

occurrence is unlikely if  $\mu_1 - \mu_2 = 5$ , and we would thus doubt this claim.

# Supplementary Exercises

#### 75.

- **a.**  $p_X(x)$  is obtained by adding joint probabilities across the row labeled x, resulting in  $p_X(x) = .2, .5, .3$  for x = 12, 15, 20 respectively. Similarly, from column sums  $p_y(y) = .1, .35, .55$  for y = 12, 15, 20 respectively.
- **b.**  $P(X \le 15 \text{ and } Y \le 15) = p(12,12) + p(12,15) + p(15,12) + p(15,15) = .25.$
- c.  $p_X(12) \cdot p_Y(12) = (.2)(.1) \neq .05 = p(12,12)$ , so X and Y are not independent. (Almost any other (x, y) pair yields the same conclusion).

d. 
$$E(X+Y) = \sum \sum (x+y)p(x,y) = 33.35$$
 (or  $= E(X) + E(Y) = 33.35$ )

e.  $E(|X-Y|) = \sum \sum |x-y| p(x,y) = \dots = 3.85$ .

77.

**a.**  $1 = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y) dx dy = \int_{0}^{20} \int_{20-x}^{30-x} kxy dy dx + \int_{20}^{30} \int_{0}^{30-x} kxy dy dx = \frac{81,250}{3} \cdot k \Rightarrow k = \frac{3}{81,250} \cdot k \Rightarrow k = \frac{3}{81,250}$ 

**b.** 
$$f_{x}(x) = \begin{cases} \int_{20-x}^{30-x} kxy dy = k(250x - 10x^{2}) & 0 \le x \le 20 \\ \int_{0}^{30-x} kxy dy = k(450x - 30x^{2} + \frac{1}{2}x^{3}) & 20 \le x \le 30 \end{cases}$$

By symmetry,  $f_{x}(y)$  is obtained by substituting y for x in  $f_{x}(x)$ .

Since  $f_X(25) > 0$  and  $f_Y(25) > 0$ , but f(25, 25) = 0,  $f_X(x) \cdot f_Y(y) \neq f(x,y)$  for all (x, y), so X and Y are not independent.

e. 
$$P(X+Y \le 25) = \int_0^{20} \int_{20-x}^{25-x} kxy dy dx + \int_{20}^{25} \int_0^{25-x} kxy dy dx = \frac{3}{81,250} \cdot \frac{230,625}{24} = .355.$$

**d.** 
$$E(X+Y) = E(X) + E(Y) = 2E(X) = 2\left\{\int_{0}^{20} x \cdot k\left(250x - 10x^{2}\right)dx + \int_{20}^{30} x \cdot k\left(450x - 30x^{2} + \frac{1}{2}x^{3}\right)dx\right\} = 2k(351,666.67) = 25.969.$$

e.  $E(XY) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xy \cdot f(x, y) dx dy = \int_{0}^{20} \int_{20-x}^{30-x} kx^2 y^2 dy dx$ + $\int_{20}^{30} \int_{0}^{30-x} kx^2 y^2 dy dx = \frac{k}{3} \cdot \frac{33,250,000}{3} = 136.4103$ , so  $Cov(X, Y) = 136.4103 - (12.9845)^2 = -32.19$ .  $E(X^2) = E(Y^2) = 204.6154$ , so  $\sigma_X^2 = \sigma_Y^2 = 204.6154 - (12.9845)^2 = 36.0182$  and  $\rho = \frac{-32.19}{36.0182} = -.894$ .

**f.** 
$$V(X + Y) = V(X) + V(Y) + 2Cov(X, Y) = 7.66$$

79. 
$$E(\overline{X} + \overline{Y} + \overline{Z}) = 500 + 900 + 2000 =$$

$$V(\overline{X} + \overline{Y} + \overline{Z}) = \frac{50^2}{365} + \frac{100^2}{365} + \frac{180^2}{365} = 123.014 \implies SD(\overline{X} + \overline{Y} + \overline{Z}) = 11.09.$$
  
$$P(\overline{X} + \overline{Y} + \overline{Z} \le 3500) = P(Z \le 9.0) \approx 1.$$

3400.

81.

- **a.**  $E(N) \cdot \mu = (10)(40) = 400$  minutes.
- **b.** We expect 20 components to come in for repair during a 4 hour period, so  $E(N) \cdot \mu = (20)(3.5) = 70$ .

83. 
$$0.95 = P(\mu - .02 \le \overline{X} \le \mu + .02) = P\left(\frac{-.02}{.1/\sqrt{n}} \le Z \le \frac{.02}{.1/\sqrt{n}}\right) = P\left(-.2\sqrt{n} \le Z \le .2\sqrt{n}\right) \text{ ; since}$$

 $P(-1.96 \le Z \le 1.96) = .95$ ,  $.2\sqrt{n} = 1.96 \Rightarrow n = 97$ . The Central Limit Theorem justifies our use of the normal distribution here.

85. The expected value and standard deviation of volume are 87,850 and 4370.37, respectively, so  $P(\text{volume} \le 100,000) = P\left(Z \le \frac{100,000 - 87,850}{4370.37}\right) = P(Z \le 2.78) = .9973$ .

87.

**a.** 
$$P(12 < X < 15) = P\left(\frac{12 - 13}{4} < Z < \frac{15 - 13}{4}\right) = P(-0.25 < Z < 0.5) = .6915 - .4013 = .2092.$$

**b.** Since individual times are normally distributed,  $\overline{X}$  is also normal, with the same mean  $\mu = 13$  but with standard deviation  $\sigma_{\overline{X}} = \sigma / \sqrt{n} = 4 / \sqrt{16} = 1$ . Thus,

$$P(12 < \overline{X} < 15) = P\left(\frac{12 - 13}{1} < Z < \frac{15 - 13}{1}\right) = P(-1 < Z < 2) = .9772 - .1587 = .8185.$$

c. The mean is  $\mu = 13$ . A sample mean  $\overline{X}$  based on n = 16 observation is likely to be closer to the true mean than is a single observation X. That's because both are "centered" at  $\mu$ , but the decreased variability in  $\overline{X}$  gives it less ability to vary significantly from  $\mu$ .

**d.** 
$$P(\overline{X} > 20) = 1 - \Phi(7) \approx 1 - 1 = 0.$$

89.

**a.**  $V(aX+Y) = a^2 \sigma_X^2 + 2a \text{Cov}(X,Y) + \sigma_Y^2 = a^2 \sigma_X^2 + 2a \sigma_X \sigma_Y \rho + \sigma_Y^2.$ 

Substituting  $a = \frac{\sigma_{\gamma}}{\sigma_{\chi}}$  yields  $\sigma_{\gamma}^2 + 2\sigma_{\gamma}^2 \rho + \sigma_{\gamma}^2 = 2\sigma_{\gamma}^2 (1+\rho) \ge 0$ . This implies  $(1+\rho) \ge 0$ , or  $\rho \ge -1$ .

- **b.** The same argument as in a yields  $2\sigma_{\gamma}^2(1-\rho) \ge 0$ , from which  $\rho \le 1$ .
- c. Suppose  $\rho = 1$ . Then  $V(aX Y) = 2\sigma_Y^2(1 \rho) = 0$ , which implies that aX Y is a constant. Solve for Y and Y = aX (constant), which is of the form aX + b.
- 91.
- **a.** With  $Y = X_1 + X_2$ ,  $F_Y(y) = \int_0^y \left\{ \int_0^{y-x_1} \frac{1}{2^{v_1/2} \Gamma(v_1/2)} \cdot \frac{1}{2^{v_{21}/2} \Gamma(v_2/2)} \cdot x_1^{\frac{v_1}{2}} x_2^{\frac{v_2}{2}} e^{\frac{x_1+x_2}{2}} dx_2 \right\} dx_1$ . But the

inner integral can be shown to be equal to  $\frac{1}{2^{(\nu_1+\nu_2)/2}\Gamma((\nu_1+\nu_2)/2)}y^{\lfloor(\nu_1+\nu_2)/2\rfloor-1}e^{-y/2}$ , from which the result follows.

**b.** By a,  $Z_1^2 + Z_2^2$  is chi-squared with v = 2, so  $(Z_1^2 + Z_2^2) + Z_3^2$  is chi-squared with v = 3, etc., until  $Z_1^2 + ... + Z_n^2$  is chi-squared with v = n.

c.  $\frac{X_i - \mu}{\sigma}$  is standard normal, so  $\left[\frac{X_i - \mu}{\sigma}\right]^2$  is chi-squared with v = 1, so the sum is chi-squared with parameter v = n.

I Court V V)-

93.

**a.** 
$$V(X_1) = V(W + E_1) = \sigma_W^2 + \sigma_E^2 = V(W + E_2) = V(X_2)$$
 and  $Cov(X_1, X_2)$   
 $Cov(W + E_1, W + E_2) = Cov(W, W) + Cov(W, E_2) + Cov(E_1, W) + Cov(E_1, E_2) =$   
 $Cov(W, W) + 0 + 0 + 0 = V(W) = \sigma_W^2$ .  
Thus,  $\rho = \frac{\sigma_W^2}{\sqrt{\sigma_W^2 + \sigma_E^2} \cdot \sqrt{\sigma_W^2 + \sigma_E^2}} = \frac{\sigma_W^2}{\sigma_W^2 + \sigma_E^2}$ .

**b.** 
$$\rho = \frac{1}{1+.0001} = .9999.$$

95.

 $E(Y) \doteq h(\mu_1, \mu_2, \mu_3, \mu_4) = 120 \left[\frac{1}{10} + \frac{1}{15} + \frac{1}{20}\right] = 26.$ 

The partial derivatives of  $h(\mu_1, \mu_2, \mu_3, \mu_4)$  with respect to  $x_1, x_2, x_3$ , and  $x_4$  are  $-\frac{x_4}{x_1^2}, -\frac{x_4}{x_2^2}, -\frac{x_4}{x_3^2}$ , and

 $\frac{1}{x_1} + \frac{1}{x_2} + \frac{1}{x_3}$ , respectively. Substituting  $x_1 = 10$ ,  $x_2 = 15$ ,  $x_3 = 20$ , and  $x_4 = 120$  gives -1.2, -.5333, -.3000, and .2167, respectively, so  $V(Y) = (1)(-1.2)^2 + (1)(-.5333)^2 + (1.5)(-.3000)^2 + (4.0)(.2167)^2 = 2.6783$ , and the approximate sd of Y is 1.64.

97. Since X and Y are standard normal, each has mean 0 and variance 1.

**a.** Cov(X, U) = Cov(X, .6X + .8Y) = .6Cov(X, X) + .8Cov(X, Y) = .6V(X) + .8(0) = .6(1) = .6.The covariance of X and Y is zero because X and Y are independent. Also,  $V(U) = V(.6X + .8Y) = (.6)^2 V(X) + (.8)^2 V(Y) = (.36)(1) + (.64)(1) = 1$ . Therefore,

$$\operatorname{Corr}(X,U) = \frac{\operatorname{Cov}(X,U)}{\sigma_X \sigma_U} = \frac{.6}{\sqrt{1}\sqrt{1}} = .6, \text{ the coefficient on } X.$$

**b.** Based on part **a**, for any specified  $\rho$  we want  $U = \rho X + bY$ , where the coefficient *b* on *Y* has the feature that  $\rho^2 + b^2 = 1$  (so that the variance of *U* equals 1). One possible option for *b* is  $b = \sqrt{1 - \rho^2}$ , from which  $U = \rho X + \sqrt{1 - \rho^2} Y$ .

93

der midden bedri Enske of gun men Gregilich den Thriede an Initialia

# **CHAPTER 6**

# Section 6.1

1.

**a.** We use the sample mean,  $\overline{x}$ , to estimate the population mean  $\mu$ .  $\hat{\mu} = \overline{x} = \frac{\Sigma x_i}{n} = \frac{219.80}{27} = 8.1407$ .

**b.** We use the sample median,  $\tilde{x} = 7.7$  (the middle observation when arranged in ascending order).

c. We use the sample standard deviation, 
$$s = \sqrt{s^2} = \sqrt{\frac{1860.94 - \frac{(219.8)^2}{27}}{26}} = 1.660.$$

**d.** With "success" = observation greater than 10, x = # of successes = 4, and  $\hat{p} = \frac{x}{n} = \frac{4}{27} = .1481$ .

e. We use the sample (std dev)/(mean), or  $\frac{s}{\overline{x}} = \frac{1.660}{8.1407} = .2039$ .

3.

5.

- a. We use the sample mean,  $\overline{x} = 1.3481$ .
- **b.** Because we assume normality, the mean = median, so we also use the sample mean  $\overline{x} = 1.3481$ . We could also easily use the sample median.
- c. We use the 90<sup>th</sup> percentile of the sample:  $\hat{\mu} + (1.28)\hat{\sigma} = \overline{x} + 1.28s = 1.3481 + (1.28)(.3385) = 1.7814$ .
- d. Since we can assume normality,

$$P(X < 1.5) \approx P\left(Z < \frac{1.5 - \overline{x}}{s}\right) = P\left(Z < \frac{1.5 - 1.3481}{.3385}\right) = P(Z < .45) = .6736.$$

e. The estimated standard error of  $\overline{x} = \frac{\hat{\sigma}}{\sqrt{n}} = \frac{s}{\sqrt{n}} = \frac{.3385}{\sqrt{16}} = .0846.$ 

Let  $\theta$  = the total audited value. Three potential estimators of  $\theta$  are  $\hat{\theta}_1 = N\overline{X}$ ,  $\hat{\theta}_2 = T - N\overline{D}$ , and  $\hat{\theta}_3 = T \cdot \frac{\overline{X}}{\overline{Y}}$ From the data,  $\overline{y} = 374.6$ ,  $\overline{x} = 340.6$ , and  $\overline{d} = 34.0$ . Knowing N = 5,000 and T = 1,761,300, the three corresponding estimates are  $\hat{\theta}_1 = (5,000)(340.6) = 1,703,000$ ,  $\hat{\theta}_2 = 1,761,300 - (5,000)(34.0) = 1,591,300$ , and  $\hat{\theta}_3 = 1,761,300 \left(\frac{340.6}{374.6}\right) = 1,601,438.281$ .

**a.** 
$$\hat{\mu} = \overline{x} = \frac{\sum x_i}{n} = \frac{1206}{10} = 120.6.$$

**b.**  $\hat{\tau} = 10,000 \ \hat{\mu} = 1,206,000.$ 

- c. 8 of 10 houses in the sample used at least 100 therms (the "successes"), so  $\hat{p} = \frac{8}{10} = .80$ .
- **d.** The ordered sample values are 89, 99, 103, 109, 118, 122, 125, 138, 147, 156, from which the two middle values are 118 and 122, so  $\hat{\mu} = \tilde{x} = \frac{118 + 122}{2} = 120.0$ .

9.

7.

**a.**  $E(\overline{X}) = \mu = E(X)$ , so  $\overline{X}$  is an unbiased estimator for the Poisson parameter  $\mu$ . Since n = 150,  $\hat{\mu} = \overline{x} = \frac{\sum x_i}{n} = \frac{(0)(18) + (1)(37) + \dots + (7)(1)}{150} = \frac{317}{150} = 2.11$ .

**b.**  $\sigma_{\bar{x}} = \frac{\sigma}{\sqrt{n}} = \frac{\sqrt{\mu}}{\sqrt{n}}$ , so the estimated standard error is  $\sqrt{\frac{\hat{\mu}}{n}} = \frac{\sqrt{2.11}}{\sqrt{150}} = .119$ .

11.

**a.** 
$$E\left(\frac{X_1}{n_1}-\frac{X_2}{n_2}\right) = \frac{1}{n_1}E(X_1)-\frac{1}{n_2}E(X_2) = \frac{1}{n_1}(n_1p_1)-\frac{1}{n_2}(n_2p_2) = p_1-p_2.$$

**b.** 
$$V\left(\frac{X_1}{n_1} - \frac{X_2}{n_2}\right) = V\left(\frac{X_1}{n_1}\right) + V\left(\frac{X_2}{n_2}\right) = \left(\frac{1}{n_1}\right)^2 V(X_1) + \left(\frac{1}{n_2}\right)^2 V(X_2) = \frac{1}{n_1^2} (n_1 p_1 q_1) + \frac{1}{n_2^2} (n_2 p_2 q_2) = \frac{p_1 q_1}{n_1} + \frac{p_2 q_2}{n_2}, \text{ and the standard error is the square root of this quantity.}$$

c. With 
$$\hat{p}_1 = \frac{x_1}{n_1}$$
,  $\hat{q}_1 = 1 - \hat{p}_1$ ,  $\hat{p}_2 = \frac{x_2}{n_2}$ ,  $\hat{q}_2 = 1 - \hat{p}_2$ , the estimated standard error is  $\sqrt{\frac{\hat{p}_1\hat{q}_1}{n_1} + \frac{\hat{p}_2\hat{q}_2}{n_2}}$ .

**d.** 
$$(\hat{p}_1 - \hat{p}_2) = \frac{127}{200} - \frac{176}{200} = .635 - .880 = -.245$$

e. 
$$\sqrt{\frac{(.635)(.365)}{200} + \frac{(.880)(.120)}{200}} = .041$$

$$\mu = E(X) = \int_{-1}^{1} x \cdot \frac{1}{2} (1 + \theta x) dx = \frac{x^2}{4} + \frac{\theta x^3}{6} \Big|_{-1}^{1} = \frac{1}{3} \theta \Longrightarrow \theta = 3\mu \Longrightarrow$$
$$\hat{\theta} = 3\overline{X} \Longrightarrow E(\hat{\theta}) = E(3\overline{X}) = 3E(\overline{X}) = 3\mu = 3\left(\frac{1}{3}\right)\theta = \theta.$$

13.

15.

a.

$$E(X^{2}) = 2\theta \text{ implies that } E\left(\frac{X^{2}}{2}\right) = \theta \text{. Consider } \hat{\theta} = \frac{\sum X_{i}^{2}}{2n} \text{. Then}$$
$$E\left(\hat{\theta}\right) = E\left(\frac{\sum X_{i}^{2}}{2n}\right) = \frac{\sum E(X_{i}^{2})}{2n} = \frac{\sum 2\theta}{2n} = \frac{2n\theta}{2n} = \theta \text{, implying that } \hat{\theta} \text{ is an unbiased estimator for } \theta$$

**b.** 
$$\sum x_i^2 = 1490.1058$$
, so  $\hat{\theta} = \frac{1490.1058}{20} = 74.505$ .

17.

**a.** 
$$E(\hat{p}) = \sum_{x=0}^{\infty} \frac{r-1}{x+r-1} \cdot \binom{x+r-1}{x} \cdot p^r \cdot (1-p)^x$$
$$= p \sum_{x=0}^{\infty} \frac{(x+r-2)!}{x!(r-2)!} \cdot p^{r-1} \cdot (1-p)^x = p \sum_{x=0}^{\infty} \binom{x+r-2}{x} p^{r-1} (1-p)^x = p \sum_{x=0}^{\infty} nb(x;r-1,p) = p.$$

**b.** For the given sequence, x = 5, so  $\hat{p} = \frac{5-1}{5+5-1} = \frac{4}{9} = .444$ .

19.

**a.**  $\lambda = .5p + .15 \Longrightarrow 2\lambda = p + .3$ , so  $p = 2\lambda - .3$  and  $\hat{p} = 2\hat{\lambda} - .3 = 2\left(\frac{Y}{n}\right) - .3$ ; the estimate is  $2\left(\frac{20}{80}\right) - .3 = .2$ .

**b.**  $E(\hat{p}) = E(2\hat{\lambda} - .3) = 2E(\hat{\lambda}) - .3 = 2\lambda - .3 = p$ , as desired.

c. Here 
$$\lambda = .7 p + (.3)(.3)$$
, so  $p = \frac{10}{7}\lambda - \frac{9}{70}$  and  $\hat{p} = \frac{10}{7}\left(\frac{Y}{n}\right) - \frac{9}{70}$ .

# Section 6.2

a.

21.

$$E(X) = \beta \cdot \Gamma\left(1 + \frac{1}{\alpha}\right) \text{ and } E(X^2) = V(X) + [E(X)]^2 = \beta^2 \Gamma\left(1 + \frac{2}{\alpha}\right), \text{ so the moment estimators } \hat{\alpha} \text{ and } \hat{\beta} \text{ are the solution to } \overline{X} = \hat{\beta} \cdot \Gamma\left(1 + \frac{1}{\hat{\alpha}}\right), \frac{1}{n} \sum X_i^2 = \hat{\beta}^2 \Gamma\left(1 + \frac{2}{\hat{\alpha}}\right). \text{ Thus } \hat{\beta} = \frac{\overline{X}}{\Gamma\left(1 + \frac{1}{\hat{\alpha}}\right)}, \text{ so once } \hat{\alpha}$$
  
has been determined  $\Gamma\left(1 + \frac{1}{\hat{\alpha}}\right)$  is evaluated and  $\hat{\beta}$  then computed. Since  $\overline{X}^2 = \hat{\beta}^2 \cdot \Gamma^2\left(1 + \frac{1}{\hat{\alpha}}\right), \frac{1}{n} \sum \frac{X_i^2}{\overline{X}^2} = \frac{\Gamma\left(1 + \frac{2}{\hat{\alpha}}\right)}{\Gamma^2\left(1 + \frac{1}{\hat{\alpha}}\right)}, \text{ so this equation must be solved to obtain } \hat{\alpha}.$ 

**b.** From **a**, 
$$\frac{1}{20} \left( \frac{16,500}{28.0^2} \right) = 1.05 = \frac{\Gamma\left(1 + \frac{2}{\hat{\alpha}}\right)}{\Gamma^2\left(1 + \frac{1}{\hat{\alpha}}\right)}$$
, so  $\frac{1}{1.05} = .95 = \frac{\Gamma^2\left(1 + \frac{1}{\hat{\alpha}}\right)}{\Gamma\left(1 + \frac{2}{\hat{\alpha}}\right)}$ , and from the hint,  
 $\frac{1}{\hat{\alpha}} = .2 \Rightarrow \hat{\alpha} = 5$ . Then  $\hat{\beta} = \frac{\overline{x}}{\Gamma(1.2)} = \frac{28.0}{\Gamma(1.2)}$ .

23.

Determine the joint pdf (aka the likelihood function), take a logarithm, and then use calculus:  

$$f(x_1, ..., x_n \mid \theta) = \prod_{i=1}^n \frac{1}{\sqrt{2\pi\theta}} e^{-x_i^2/2\theta} = (2\pi\theta)^{-n/2} e^{-\sum x_i^2/2\theta}$$

$$\ell(\theta) = \ln[f(x_1, ..., x_n \mid \theta)] = -\frac{n}{2} \ln(2\pi) - \frac{n}{2} \ln(\theta) - \sum x_i^2/2\theta$$

$$\ell'(\theta) = 0 - \frac{n}{2\theta} + \sum x_i^2/2\theta^2 = 0 \Rightarrow -n\theta + \sum x_i^2 = 0$$

Solving for  $\theta$ , the maximum likelihood estimator is  $\hat{\theta} = \frac{1}{n} \sum X_i^2$ .

25.

**a.** 
$$\hat{\mu} = \overline{x} = 384.4; s^2 = 395.16$$
, so  $\frac{1}{n} \sum (x_i - \overline{x})^2 = \hat{\sigma}^2 = \frac{9}{10} (395.16) = 355.64$  and  $\hat{\sigma} = \sqrt{355.64} = 18.86$  (this is not s).

**b.** The 95<sup>th</sup> percentile is  $\mu + 1.645\sigma$ , so the mle of this is (by the invariance principle)  $\hat{\mu} + 1.645\hat{\sigma} = 415.42$ .

c. The mle of  $P(X \le 400)$  is, by the invariance principle,  $\Phi\left(\frac{400 - \hat{\mu}}{\hat{\sigma}}\right) = \Phi\left(\frac{400 - 384.4}{18.86}\right) = \Phi(0.83) = .7967.$ 

27.

29.

a. The joint pdf (likelihood function) is  $f(x_1,...,x_n;\lambda,\theta) = \begin{cases} \lambda^n e^{-\lambda \Sigma(x_i-\theta)} & x_1 \ge \theta,...,x_n \ge \theta \\ 0 & \text{otherwise} \end{cases}$ Notice that  $x_1 \ge \theta,...,x_n \ge \theta$  iff  $\min(x_i) \ge \theta$ , and that  $-\lambda \Sigma(x_i-\theta) = -\lambda \Sigma x_i + n\lambda \theta$ . Thus likelihood  $= \begin{cases} \lambda^n \exp(-\lambda \Sigma x_i) \exp(n\lambda \theta) & \min(x_i) \ge \theta \\ 0 & \min(x_i) < \theta \end{cases}$ 

Consider maximization with respect to  $\theta$ . Because the exponent  $n\lambda\theta$  is positive, increasing  $\theta$  will increase the likelihood provided that  $\min(x_i) \ge \theta$ ; if we make  $\theta$  larger than  $\min(x_i)$ , the likelihood drops to 0. This implies that the mle of  $\theta$  is  $\hat{\theta} = \min(x_i)$ . The log likelihood is now  $n\ln(\lambda) - \lambda\Sigma(x_i - \hat{\theta})$ . Equating the derivative w.r.t.  $\lambda$  to 0 and solving yields  $\hat{\lambda} = \frac{n}{\Sigma(x_i - \hat{\theta})} = \frac{n}{\Sigma x_i - n\hat{\theta}}$ 

**b.**  $\hat{\theta} = \min(x_i) = .64$ , and  $\Sigma x_i = 55.80$ , so  $\hat{\lambda} = \frac{10}{55.80 - 6.4} = .202$ 

# Supplementary Exercises

- 31. Substitute  $k = \varepsilon/\sigma_Y$  into Chebyshev's inequality to write  $P(|Y \mu_Y| \ge \varepsilon) \le 1/(\varepsilon/\sigma_Y)^2 = V(Y)/\varepsilon^2$ . Since  $E(\overline{X}) = \mu$  and  $V(\overline{X}) = \sigma^2 / n$ , we may then write  $P(|\overline{X} \mu| \ge \varepsilon) \le \frac{\sigma^2 / n}{\varepsilon^2}$ . As  $n \to \infty$ , this fraction converges to 0, hence  $P(|\overline{X} \mu| \ge \varepsilon) \to 0$ , as desired.
- 33. Let  $x_1$  = the time until the first birth,  $x_2$  = the elapsed time between the first and second births, and so on. Then  $f(x_1,...,x_n;\lambda) = \lambda e^{-\lambda x_1} \cdot (2\lambda) e^{-2\lambda x_2} ... (n\lambda) e^{-n\lambda x_n} = n!\lambda^n e^{-\lambda \Sigma k x_k}$ . Thus the log likelihood is  $\ln(n!) + n\ln(\lambda) - \lambda \Sigma k x_k$ . Taking  $\frac{d}{d\lambda}$  and equating to 0 yields  $\hat{\lambda} = \frac{n}{\Sigma k x_k}$ . For the given sample, n = 6,  $x_1 = 25.2$ ,  $x_2 = 41.7 - 25.2 = 16.5$ ,  $x_3 = 9.5$ ,  $x_4 = 4.3$ ,  $x_5 = 4.0$ ,  $x_6 = 2.3$ ; so  $\sum_{i=1}^{6} k x_k = (1)(25.2) + (2)(16.5) + ... + (6)(2.3) = 137.7$  and  $\hat{\lambda} = \frac{6}{137.7} = .0436$ .

-	6		
	P	5	
-	۰.	э	

$x_i + x_i$	23.5	26.3	28.0	28.2	29.4	29.5	30.6	31.6	33.9	49.3
23.5	23.5	24.9	25.75	25.85	26.45	26.5	27.05	27.55	28.7	36.4
26.3		26.3	27.15	27.25	27.85	27.9	28.45	28.95	30.1	37.8
28.0			28.0	28.1	28.7	28.75	29.3	29.8	30.95	38.65
28.2	5.00			28.2	28.8	28.85	29.4	29.9	31.05	38.75
29.4					29.4	29.45	30.0	30.5	30.65	39.35
29.5	and Down					29.5	30.05	30.55	31.7	39.4
30.6							30.6	31.1	32.25	39.95
31.6	05 0000							31.6	32.75	40.45
33.9									33.9	41.6
49.3	meth.									49.3

There are 55 averages, so the median is the 28<sup>th</sup> in order of increasing magnitude. Therefore,  $\hat{\mu} = 29.5$ .

37. Let  $c = \frac{\Gamma(\frac{n-1}{2})}{\Gamma(\frac{n}{2}) \cdot \sqrt{\frac{2}{n-1}}}$ . Then E(cS) = cE(S), and c cancels with the two  $\Gamma$  factors and the square root in E(S),

leaving just  $\sigma$ . When n = 20,  $c = \frac{\Gamma(9.5)}{\Gamma(10) \cdot \sqrt{\frac{2}{19}}} = \frac{(8.5)(7.5)\cdots(.5)\Gamma(.5)}{(10-1)!\sqrt{\frac{2}{19}}} = \frac{(8.5)(7.5)\cdots(.5)\sqrt{\pi}}{9!\sqrt{\frac{2}{19}}} = 1.0132.$ 

99

# CHAPTER 7

## Section 7.1

- **a.**  $z_{\alpha/2} = 2.81$  implies that  $\alpha/2 = 1 \Phi(2.81) = .0025$ , so  $\alpha = .005$  and the confidence level is  $100(1-\alpha)\% = 99.5\%$ .
- **b.**  $z_{\alpha/2} = 1.44$  implies that  $\alpha = 2[1 \Phi(1.44)] = .15$ , and the confidence level is  $100(1-\alpha)\% = 85\%$ .
- c. 99.7% confidence implies that  $\alpha = .003$ ,  $\alpha/2 = .0015$ , and  $z_{.0015} = 2.96$ . (Look for cumulative area equal to 1 .0015 = .9985 in the main body of table A.3.) Or, just use  $z \approx 3$  by the empirical rule.
- **d.** 75% confidence implies  $\alpha = .25$ ,  $\alpha/2 = .125$ , and  $z_{.125} = 1.15$ .
- 3.

1.

- **a.** A 90% confidence interval will be narrower. The z critical value for a 90% confidence level is 1.645, smaller than the z of 1.96 for the 95% confidence level, thus producing a narrower interval.
- b. Not a correct statement. Once and interval has been created from a sample, the mean μ is either enclosed by it, or not. We have 95% confidence in the general procedure, under repeated and independent sampling.
- c. Not a correct statement. The interval is an estimate for the population mean, not a boundary for population values.
- d. Not a correct statement. In theory, if the process were repeated an infinite number of times, 95% of the intervals would contain the population mean μ. We *expect* 95 out of 100 intervals will contain μ, but we don't know this to be true.
- 5.

a. 
$$4.85 \pm \frac{(1.96)(.75)}{\sqrt{20}} = 4.85 \pm .33 = (4.52, 5.18).$$

**b.** 
$$z_{\alpha/2} = z.01 = 2.33$$
, so the interval is  $4.56 \pm \frac{(2.33)(.75)}{\sqrt{16}} = (4.12, 5.00).$ 

**c.** 
$$n = \left[\frac{2(1.96)(.75)}{.40}\right]^2 = 54.02 \nearrow 55.$$

**d.** Width 
$$w = 2(.2) = .4$$
, so  $n = \left[\frac{2(2.58)(.75)}{.4}\right]^2 = 93.61 \nearrow 94$ .

100

## Chapter 7: Statistical Intervals Based on a Single Sample

7. If  $L = 2z_{4/2} \frac{\sigma}{\sqrt{n}}$  and we increase the sample size by a factor of 4, the new length is

 $L' = 2z_{\frac{q}{2}} \frac{\sigma}{\sqrt{4n}} = \left[2z_{\frac{q}{2}} \frac{\sigma}{\sqrt{n}}\right] \left(\frac{1}{2}\right) = \frac{L}{2}.$  Thus halving the length requires n to be increased fourfold. If n' = 25n, then  $L' = \frac{L}{5}$ , so the length is decreased by a factor of 5.

9.

D. P.

**a.** 
$$\left(\overline{x} - 1.645 \frac{\sigma}{\sqrt{n}}, \infty\right)$$
. From 5**a**,  $\overline{x} = 4.85$ ,  $\sigma = .75$ , and  $n = 20$ ;  $4.85 - 1.645 \frac{.75}{\sqrt{20}} = 4.5741$ , so the interval is  $(4.5741 \ \infty)$ 

**b.** 
$$\left(\overline{x}-z_{\alpha}\frac{\sigma}{\sqrt{n}},\infty\right)$$

c. 
$$\left(-\infty, \overline{x} + z_{\alpha} \frac{\sigma}{\sqrt{n}}\right)$$
; From 4a,  $\overline{x} = 58.3$ ,  $\sigma = 3.0$ , and  $n = 25$ ;  $58.3 + 2.33 \frac{3}{\sqrt{25}} = 59.70$ , so the interval is  $(-\infty, 59.70)$ .

11. *Y* is a binomial rv with n = 1000 and p = .95, so E(Y) = np = 950, the expected number of intervals that capture  $\mu$ , and  $\sigma_r = \sqrt{npq} = 6.892$ . Using the normal approximation to the binomial distribution,  $P(940 \le Y \le 960) = P(939.5 \le Y \le 960.5) \approx P(-1.52 \le Z \le 1.52) = .9357 - .0643 = .8714.$ 

# Section 7.2

13.

**a.**  $\overline{x} \pm z_{.025} \frac{s}{\sqrt{n}} = 654.16 \pm 1.96 \frac{164.43}{\sqrt{50}} = (608.58, 699.74)$ . We are 95% confident that the true average CO<sub>2</sub> level in this population of homes with gas cooking appliances is between 608.58ppm and 699.74ppm

**b.** 
$$w = 50 = \frac{2(1.96)(175)}{\sqrt{n}} \Rightarrow \sqrt{n} = \frac{2(1.96)(175)}{50} = 13.72 \Rightarrow n = (13.72)^2 = 188.24$$
, which rounds up to 189.

15.

- **a.**  $z_a = .84$ , and  $\Phi(.84) = .7995 \approx .80$ , so the confidence level is 80%.
- **b.**  $z_{\alpha} = 2.05$ , and  $\Phi(2.05) = .9798 \approx .98$ , so the confidence level is 98%.
- c.  $z_{\alpha} = .67$ , and  $\Phi(.67) = .7486 \approx .75$ , so the confidence level is 75%.

17.  $\overline{x} - z_{.01} \frac{s}{\sqrt{n}} = 135.39 - 2.33 \frac{4.59}{\sqrt{153}} = 135.39 - .865 = 134.53$ . We are 99% confident that the true average ultimate tensile strength is greater than 134.53.

$$\hat{p} = \frac{201}{356} = .5646 \text{ ; We calculate a 95\% confidence interval for the proportion of all dies that pass the probes}$$

$$\frac{.5646 + \frac{(1.96)^2}{2(356)} \pm 1.96\sqrt{\frac{(.5646)(.4354)}{356} + \frac{(1.96)^2}{4(356)^2}}}{1 + \frac{(1.96)^2}{356}} = \frac{.5700 \pm .0518}{1.01079} = (.513, .615) \text{ . The simpler CI formula}$$

$$(7.11) \text{ gives } .5646 \pm 1.96\sqrt{\frac{.5646(.4354)}{356}} = (.513, .616), \text{ which is almost identical.}$$

19.

For a one-sided bound, we need  $z_a = z_{.05} = 1.645$ ;  $\hat{p} = \frac{250}{1000} = .25$ ; and  $\tilde{p} = \frac{.25 + 1.645^2 / 2000}{1 + 1.645^2 / 1000} = .2507$ . The resulting 95% upper confidence bound for p, the true proportion of such consumers who never apply for a

rebate, is 
$$.2507 + \frac{1.645\sqrt{(.25)(.75)/1000 + (1.645)^2/(41000^2)}}{1+(1.645)^2/1000} = .2507 + .0225 = .2732.$$

Yes, there is compelling evidence the true proportion is less than 1/3 (.3333), since we are 95% confident this true proportion is less than .2732.

23.

25.

**a.** With such a large sample size, we can use the "simplified" CI formula (7.11). With  $\hat{p} = .25$ , n = 2003, and  $z_{n/2} = z_{.005} = 2.576$ , the 99% confidence interval for p is

$$\hat{p} \pm z_{a/2} \sqrt{\frac{\hat{p}\hat{q}}{n}} = .25 \pm 2.576 \sqrt{\frac{(.25)(.75)}{2003}} = .25 \pm .025 = (.225, .275).$$

**b.** Using the "simplified" formula for sample size and  $\hat{p} = \hat{q} = .5$ ,

$$n = \frac{4z^2 \hat{p}\hat{q}}{w^2} = \frac{4(2.576)^2 (.5)(.5)}{(.05)^2} = 2654.31$$

So, a sample of size at least 2655 is required. (We use  $\hat{p} = \hat{q} = .5$  here, rather than the values from the sample data, so that our CI has the desired width irrespective of what the true value of p might be. See the textbook discussion toward the end of Section 7.2.)

**a.** 
$$n = \frac{2(1.96)^2 (.25) - (1.96)^2 (.01) \pm \sqrt{4(1.96)^4 (.25)(.25 - .01) + .01(1.96)^4}}{.01} \approx 381$$
  
**b.** 
$$n = \frac{2(1.96)^2 (\frac{1}{3} \cdot \frac{2}{3}) - (1.96)^2 (.01) \pm \sqrt{4(1.96)^4 (\frac{1}{3} \cdot \frac{2}{3})(\frac{1}{3} \cdot \frac{2}{3} - .01) + .01(1.96)^4}}{.01} \approx 339$$

102

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.
Note that the midpoint of the new interval is  $\frac{x+z^2/2}{n+z^2}$ , which is roughly  $\frac{x+2}{n+4}$  with a confidence level of 27. 95% and approximating 1.96  $\approx$  2. The variance of this quantity is  $\frac{np(1-p)}{(n+z^2)^2}$ , or roughly  $\frac{p(1-p)}{n+4}$ . Now replacing p with  $\frac{x+2}{n+4}$ , we have  $\left(\frac{x+2}{n+4}\right) \pm z_{\frac{9}{2}} \sqrt{\frac{\left(\frac{x+2}{n+4}\right)\left(1-\frac{x+2}{n+4}\right)}{n+4}}$ . For clarity, let  $x^* = x+2$  and  $n^* = n + 4$ , then  $\hat{p}^* = \frac{x^*}{n^*}$  and the formula reduces to  $\hat{p}^* \pm z_{\frac{n}{2}} \sqrt{\frac{\hat{p}^* \hat{q}^*}{n^*}}$ , the desired conclusion. For further discussion, see the Agresti article.

# Section 7.3

29.				
	a.	$t_{.025,10} = 2.228$	d.	$t_{.005,50} = 2.678$
	b.	$t_{.025,20} = 2.086$	e.	$t_{.01,25} = 2.485$
	c.	$t_{.005,20} = 2.845$	f.	$-t_{.025,5} = -2.571$

#### 31.

a.	$t_{05,10} = 1.812$
b.	$t_{.05,15} = 1.753$
c.	$t_{.01,15} = 2.602$

 $t_{01.4} = 3.747$  $t_{02,24} \approx t_{.025,24} = 2.064$  $t_{0137} \approx 2.429$ 

#### 33.

The boxplot indicates a very slight positive skew, with no outliers. The data appears to center near 8. 438.



Based on a normal probability plot, it is reasonable to assume the sample observations came from a b. normal distribution.

- c. With df = n 1 = 16, the critical value for a 95% CI is  $t_{.025,16} = 2.120$ , and the interval is
  - $438.29 \pm (2.120) \left(\frac{15.14}{\sqrt{17}}\right) = 438.29 \pm 7.785 = (430.51, 446.08)$ . Since 440 is within the interval, 440 is

a plausible value for the true mean. 450, however, is not, since it lies outside the interval.

**35.** 
$$n = 15, \ \overline{x} = 25.0, \ s = 3.5; \ t_{.025,14} = 2.145$$

- **a.** A 95% CI for the mean:  $25.0 \pm 2.145 \frac{3.5}{\sqrt{15}} = (23.06, 26.94).$
- **b.** A 95% prediction interval:  $25.0 \pm 2.145(3.5)\sqrt{1 + \frac{1}{15}} = (17.25, 32.75)$ . The prediction interval is about 4 times wider than the confidence interval.

#### 37.

- a. A 95% CI:  $.9255 \pm 2.093(.0181) = .9255 \pm .0379 \Rightarrow (.8876, .9634)$
- **b.** A 95% P.I. : .9255 ± 2.093(.0809) $\sqrt{1 + \frac{1}{20}} = .9255 \pm .1735 \Rightarrow (.7520, 1.0990)$
- c. A tolerance interval is requested, with k = 99, confidence level 95%, and n = 20. The tolerance critical value, from Table A.6, is 3.615. The interval is  $.9255 \pm 3.615 (.0809) \Rightarrow (.6330, 1.2180)$ .
- 39.

a. Based on the plot, generated by Minitab, it is plausible that the population distribution is normal. Normal Probability Plot



**b.** We require a tolerance interval. From table A.6, with 95% confidence, k = 95, and n=13, the tolerance critical value is 3.081.  $\overline{x} \pm 3.081s = 52.231 \pm 3.081(14.856) = 52.231 \pm 45.771 \Rightarrow (6.460, 98.002)$ .

c. A prediction interval, with  $t_{.025,12} = 2.179$ :  $52.231 \pm 2.179(14.856)\sqrt{1 + \frac{1}{13}} = 52.231 \pm 33.593 \Rightarrow (18.638,85.824)$ 

# 104

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

41. The 20 df row of Table A.5 shows that 1.725 captures upper tail area .05 and 1.325 captures upper tail area .10 The confidence level for each interval is 100(central area)%. For the first interval, central area = 1 - sum of tail areas = 1 - (.25 + .05) = .70, and for the second and third intervals the central areas are 1 - (.20 + .10) = .70 and 1 - (.15 + .15) = 70. Thus each interval has

confidence level 70%. The width of the first interval is  $\frac{s(.687+1.725)}{\sqrt{n}} = 2.412 \frac{s}{\sqrt{n}}$ , whereas the widths of

the second and third intervals are 2.185 and 2.128 standard errors respectively. The third interval, with symmetrically placed critical values, is the shortest, so it should be used. This will always be true for a t interval.

# Section 7.4

43.

- **a.**  $\chi^2_{.05,10} = 18.307$
- **b.**  $\chi^2_{.95,10} = 3.940$
- c. Since  $10.987 = \chi^2_{.975,22}$  and  $36.78 = \chi^2_{.025,22}$ ,  $P(\chi^2_{.975,22} \le \chi^2 \le \chi^2_{.025,22}) = .95$ .
- **d.** Since  $14.611 = \chi^2_{.95,25}$  and  $37.652 = \chi^2_{.05,25}$ ,  $P(\chi^2 < 14.611 \text{ or } \chi^2 > 37.652) = 1 P(\chi^2 > 14.611) + P(\chi^2 > 37.652) = (1 .95) + .05 = .10$ .
- 45. For the n = 8 observations provided, the sample standard deviation is s = 8.2115. A 99% CI for the population variance,  $\sigma^2$ , is given by  $((n-1)s^2 / \chi^2_{005,n-1}, (n-1)s^2 / \chi^2_{.995,n-1}) = (7 \cdot 8.2115^2 / 20.276, 7 \cdot 8.2115^2 / 0.989) = (23.28, 477.25)$

Taking square roots, a 99% CI for  $\sigma$  is (4.82, 21.85). Validity of this interval requires that coating layer thickness be (at least approximately) normally distributed.

# Supplementary Exercises

47.

**a.** n = 48,  $\overline{x} = 8.079a$ ,  $s^2 = 23.7017$ , and s = 4.868. A 95% CI for  $\mu$  = the true average strength is

$$\overline{x} \pm 1.96 \frac{s}{\sqrt{n}} = 8.079 \pm 1.96 \frac{4.868}{\sqrt{48}} = 8.079 \pm 1.377 = (6.702, 9.456).$$

**b.** 
$$\hat{p} = \frac{13}{48} = .2708$$
. A 95% CI for *p* is

$$\frac{.2708 + \frac{1.96^2}{2(48)} \pm 1.96\sqrt{\frac{(.2708)(.7292)}{48} + \frac{1.96^2}{4(48)^2}}}{1 + \frac{1.96^2}{48}} = \frac{.3108 \pm .1319}{1.0800} = (.166, .410)$$

49. The sample mean is the midpoint of the interval:  $\overline{x} = \frac{60.2 + 70.6}{2} = 65.4$  N. The 95% confidence margin of error for the mean must have been  $\pm 5.2$ , so  $t \cdot s / \sqrt{n} = 5.2$ . The 95% confidence margin of error for a prediction interval (i.e., an individual) is  $t \cdot s \sqrt{1 + \frac{1}{n}} = \sqrt{n + 1} \cdot t \cdot s / \sqrt{n} = \sqrt{11 + 1}(5.2) = 18.0$ . Thus, the 95% PI is  $65.4 \pm 18.0 = (47.4 \text{ N}, 83.4 \text{ N})$ . (You could also determine *t* from *n* and *a*, then *s* separately.)

51.

**a.** With 
$$\hat{p} = 31/88 = .352$$
,  $\tilde{p} = \frac{.352 + 1.96^2 / 2(88)}{1 + 1.96^2 / 88} = .358$ , and the CI is

 $.358 \pm 1.96 \frac{\sqrt{(.352)(.648) + 1.96^2 / 4(88)^2}}{1 + 1.96^2 / 88} = (.260, .456)$ . We are 95% confident that between 26.0% and 45.6% of <u>all</u> athletes under these conditions have an exercise-induced laryngeal obstruction.

- **b.** Using the "simplified" formula,  $n = \frac{4z^2 \hat{p}\hat{q}}{w^2} = \frac{4(1.96)^2(.5)(.5)}{(.04)^2} = 2401$ . So, roughly 2400 people should be surveyed to assure a width no more than .04 with 95% confidence. Using Equation (7.12) gives the almost identical n = 2398.
- c. No. The upper bound in (a) uses a z-value of  $1.96 = z_{.025}$ . So, if this is used as an upper bound (and hence .025 equals  $\alpha$  rather than  $\alpha/2$ ), it gives a (1 .025) = 97.5% upper bound. If we want a 95% confidence upper bound for p, 1.96 should be replaced by the critical value  $z_{.05} = 1.645$ .

53. With 
$$\hat{\theta} = \frac{1}{3} \left( \overline{X}_1 + \overline{X}_2 + \overline{X}_3 \right) - \overline{X}_4, \ \sigma_{\hat{\theta}}^2 = \frac{1}{9} V \left( \overline{X}_1 + \overline{X}_2 + \overline{X}_3 \right) + V \left( \overline{X}_4 \right) = \frac{1}{9} \left( \frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2} + \frac{\sigma_3^2}{n_3} \right) + \frac{\sigma_4^2}{n_4}; \ \hat{\sigma}_{\hat{\theta}} \text{ is }$$

obtained by replacing each  $\sigma_i^2$  by  $s_i^2$  and taking the square root. The large-sample interval for  $\theta$  is then

$$\frac{1}{3}\left(\overline{x}_{1}+\overline{x}_{2}+\overline{x}_{3}\right)-\overline{x}_{4}\pm z_{\alpha/2}\sqrt{\frac{1}{9}\left(\frac{s_{1}^{2}}{n_{1}}+\frac{s_{2}^{2}}{n_{2}}+\frac{s_{3}^{2}}{n_{3}}\right)+\frac{s_{4}^{2}}{n_{4}}}$$

For the given data,  $\hat{\theta} = -.50$  and  $\hat{\sigma}_{\hat{\theta}} = .1718$ , so the interval is  $-.50 \pm 1.96(.1718) = (-.84, -.16)$ .

55. The specified condition is that the interval be length .2, so 
$$n = \left[\frac{2(1.96)(.8)}{.2}\right]^2 = 245.86 \nearrow 246$$
.

57. Proceeding as in Example 7.5 with  $T_r$  replacing  $\Sigma X_i$ , the CI for  $\frac{1}{\lambda}$  is  $\left(\frac{2t_r}{\chi^2_{1-\frac{1}{2},2r}},\frac{2t_r}{\chi^2_{\frac{1}{2},2r}}\right)$  where

 $t_r = y_1 + ... + y_r + (n-r)y_r$ . In Example 6.7, n = 20, r = 10, and  $t_r = 1115$ . With df = 20, the necessary critical values are 9.591 and 34.170, giving the interval (65.3, 232.5). This is obviously an extremely wide interval. The censored experiment provides less information about  $\frac{1}{\lambda}$  than would an uncensored experiment with n = 20.

59.

**a.**  $\int_{(\alpha/2)^{1/n}}^{(1-\alpha/2)^{1/n}} nu^{n-1} du = u^n \Big]_{(\alpha/2)^{1/n}}^{(1-\alpha/2)^{1/n}} = 1 - \frac{\alpha}{2} - \frac{\alpha}{2} = 1 - \alpha$ . From the probability statement,

 $\frac{\left(\frac{\alpha_{2}}{2}\right)^{\gamma_{n}}}{\max(X_{i})} \leq \frac{1}{\theta} \leq \frac{\left(1 - \frac{\alpha_{2}}{2}\right)^{\gamma_{n}}}{\max(X_{i})}$  with probability  $1 - \alpha$ , so taking the reciprocal of each endpoint and

interchanging gives the CI  $\left(\frac{\max(X_i)}{(1-\alpha_2')^{\frac{1}{2}}}, \frac{\max(X_i)}{(\alpha_2')^{\frac{1}{2}}}\right)$  for  $\theta$ .

**b.**  $\alpha^{\gamma_n} \leq \frac{\max(X_i)}{\theta} \leq 1$  with probability  $1 - \alpha$ , so  $1 \leq \frac{\theta}{\max(X_i)} \leq \frac{1}{\alpha^{\gamma_n}}$  with probability  $1 - \alpha$ , which yields

the interval  $\left(\max(X_i), \frac{\max(X_i)}{\alpha^{\frac{1}{\lambda}}}\right)$ .

- c. It is easily verified that the interval of **b** is shorter draw a graph of  $f_U(u)$  and verify that the shortest interval which captures area  $1 \alpha$  under the curve is the rightmost such interval, which leads to the CI of **b**. With  $\alpha = .05$ , n = 5, and  $\max(x_i) = 4.2$ , this yields (4.2, 7.65).
- 61.  $\tilde{x} = 76.2$ , the lower and upper fourths are 73.5 and 79.7, respectively, and  $f_s = 6.2$ . The robust interval is  $76.2 \pm (1.93) \left(\frac{6.2}{\sqrt{22}}\right) = 76.2 \pm 2.6 = (73.6, 78.8)$ .

 $\overline{x} = 77.33$ , s = 5.037, and  $t_{.025,21} = 2.080$ , so the *t* interval is

 $77.33 \pm (2.080) \left(\frac{5.037}{\sqrt{22}}\right) = 77.33 \pm 2.23 = (75.1, 79.6)$ . The *t* interval is centered at  $\overline{x}$ , which is pulled out

to the right of  $\tilde{x}$  by the single mild outlier 93.7; the interval widths are comparable.

A type I error constructed udging use of the two comprofets featered over the other viten in these there is a 30-50 when the solit in the portification. A type II error involves todaying the solit in the portification.

We expect 25(.5) = 11.5 "moreover" when N<sub>0</sub> is true 50, any X-values less than 6 are a least at contradictory to N<sub>0</sub> as x = 0. But areas the elementics hypothesis alsong to a.2. X-values that are just to fit away on the high do are equally contradictory. There are 19 and above, So, values at heat as contradictory to S<sub>0</sub> away to S<sub>0</sub> and above.

# **CHAPTER 8**

# Section 8.1

a.

1.

Yes. It is an assertion about the value of a parameter.

**b.** No. The sample median  $\tilde{x}$  is not a parameter.

c. No. The sample standard deviation s is not a parameter.

d. Yes. The assertion is that the standard deviation of population #2 exceeds that of population #1.

e. No.  $\overline{X}$  and  $\overline{Y}$  are statistics rather than parameters, so they cannot appear in a hypothesis.

f. Yes. *H* is an assertion about the value of a parameter.

3. We reject  $H_0$  iff P-value  $\leq \alpha = .05$ .

**a.** Reject  $H_0$  **b.** Reject  $H_0$  **c.** Do not reject  $H_0$  **d.** Reject  $H_0$  **e.** Do not reject  $H_0$ 

- 5. In this formulation,  $H_0$  states the welds do not conform to specification. This assertion will not be rejected unless there is strong evidence to the contrary. Thus the burden of proof is on those who wish to assert that the specification is satisfied. Using  $H_a$ :  $\mu < 100$  results in the welds being believed in conformance unless proved otherwise, so the burden of proof is on the non-conformance claim.
- 7. Let  $\sigma$  denote the population standard deviation. The appropriate hypotheses are  $H_0$ :  $\sigma = .05 \text{ v}$ .  $H_a$ :  $\sigma < .05$ . With this formulation, the burden of proof is on the data to show that the requirement has been met (the sheaths will not be used unless  $H_0$  can be rejected in favor of  $H_a$ . Type I error: Conclude that the standard deviation is < .05 mm when it is really equal to .05 mm. Type II error: Conclude that the standard deviation is .05 mm when it is really < .05.
- 9. A type I error here involves saying that the plant is not in compliance when in fact it is. A type II error occurs when we conclude that the plant is in compliance when in fact it isn't. Reasonable people may disagree as to which of the two errors is more serious. If in your judgment it is the type II error, then the reformulation  $H_0$ :  $\mu = 150$  v.  $H_a$ :  $\mu < 150$  makes the type I error more serious.
- 11.
- a. A type I error consists of judging one of the two companies favored over the other when in fact there is a 50-50 split in the population. A type II error involves judging the split to be 50-50 when it is not.
- **b.** We expect 25(.5) = 12.5 "successes" when  $H_0$  is true. So, any X-values less than 6 are at least as contradictory to  $H_0$  as x = 6. But since the alternative hypothesis states  $p \neq .5$ , X-values that are just as far away on the <u>high</u> side are equally contradictory. Those are 19 and above. So, values at least as contradictory to  $H_0$  as x = 6 are  $\{0,1,2,3,4,5,6,19,20,21,22,23,24,25\}$ .

- c. When  $H_0$  is true, X has a binomial distribution with n = 25 and p = .5. From part (b), P-value =  $P(X \le 6 \text{ or } X \ge 19) = B(6; 25, .5) + [1 - B(18; 25, .5)] = .014$ .
- **d.** Looking at Table A.1, a two-tailed *P*-value of .044 (2 × .022) occurs when x = 7. That is, saying we'll reject  $H_0$  iff *P*-value  $\leq$  .044 must be equivalent to saying we'll reject  $H_0$  iff  $X \leq 7$  or  $X \geq 18$  (the same distance from 12.5, but on the high side). Therefore, for any value of  $p \neq .5$ ,  $\beta(p) = P(\text{do not reject } H_0 \text{ when } X \sim \text{Bin}(25, p)) = P(7 < X < 18 \text{ when } X \sim \text{Bin}(25, p)) = B(17; 25, p) B(7; 25, p)$ .  $\beta(.4) = B(17; 25, .4) - B(7, 25, .4) = .845$ , while  $\beta(.3) = B(17; 25, .3) - B(7; 25, .3) = .488$ . By symmetry (or re-computation),  $\beta(.6) = .845$  and  $\beta(.7) = .488$ .
- e. From part (c), the *P*-value associated with x = 6 is .014. Since .014  $\leq$  .044, the procedure in (d) leads us to reject  $H_0$ .
- 13.

**a.**  $H_0: \mu = 10 \text{ v}. H_a: \mu \neq 10.$ 

**b.** Since the alternative is two-sided, values at least as contradictory to  $H_0$  as  $\overline{x} = 9.85$  are not only those less than 9.85 but also those equally far from  $\mu = 10$  on the high side: i.e.,  $\overline{x}$  values  $\ge 10.15$ .

When  $H_0$  is true,  $\overline{X}$  has a normal distribution with mean  $\mu = 10$  and sd  $\frac{\sigma}{\sqrt{n}} = \frac{.200}{\sqrt{25}} = .04$ . Hence,

 $P\text{-value} = P(\overline{X} \le 9.85 \text{ or } \overline{X} \ge 10.15 \text{ when } H_0 \text{ is true}) = 2P(\overline{X} \le 9.85 \text{ when } H_0 \text{ is true}) \text{ by symmetry}$  $= 2P\left(Z < \frac{9.85 - 10}{.04}\right) = 2\Phi(-3.75) \approx 0. \text{ (Software gives the more precise } P\text{-value .00018.)}$ 

In particular, since *P*-value  $\approx 0 < \alpha = .01$ , we reject  $H_0$  at the .01 significance level and conclude that the true mean measured weight differs from 10 kg.

c. To determine  $\beta(\mu)$  for any  $\mu \neq 10$ , we must first find the threshold between *P*-value  $\leq \alpha$  and *P*-value  $> \alpha$  in terms of  $\overline{x}$ . Parallel to part (b), proceed as follows:

 $.01 = P(\text{reject } H_0 \text{ when } H_0 \text{ is true}) = 2P(\overline{X} \le \overline{x} \text{ when } H_0 \text{ is true}) = 2\Phi\left(\frac{\overline{x} - 10}{.04}\right) \Rightarrow$ 

 $\Phi\left(\frac{\overline{x}-10}{.04}\right) = .005 \Rightarrow \frac{\overline{x}-10}{.04} = -2.58 \Rightarrow \overline{x} = 9.8968$ . That is, we'd reject  $H_0$  at the  $\alpha = .01$  level iff the

observed value of  $\overline{X}$  is  $\leq 9.8968$  — or, by symmetry,  $\geq 10 + (10 - 9.8968) = 10.1032$ . Equivalently, we do not reject  $H_0$  at the  $\alpha = .01$  level if  $9.8968 < \overline{X} < 10.1032$ .

Now we can determine the chance of a type II error:

 $\beta(10.1) = P(9.8968 < \overline{X} < 10.1032 \text{ when } \mu = 10.1) = P(-5.08 < Z < .08) = .5319.$ Similarly,  $\beta(9.8) = P(9.8968 < \overline{X} < 10.1032 \text{ when } \mu = 9.8) = P(2.42 < Z < 7.58) = .0078.$ 

### Section 8.2

15. In each case, the direction of  $H_a$  indicates that the *P*-value is  $P(Z \ge z) = 1 - \Phi(z)$ . a. *P*-value =  $1 - \Phi(1.42) = .0778$ .

- **b.** *P*-value =  $1 \Phi(0.90) = .1841$ .
- c. *P*-value =  $1 \Phi(1.96) = .0250$ .
- **d.** *P*-value =  $1 \Phi(2.48) = .0066$ .
- e. *P*-value =  $1 \Phi(-.11) = .5438$ .
- 17.
- **a.**  $z = \frac{30,960 30,000}{1500 / \sqrt{16}} = 2.56$ , so *P*-value =  $P(Z \ge 2.56) = 1 \Phi(2.56) = .0052$ . Since .0052 <  $\alpha = .01$ , reject  $H_0$ .
- **b.**  $z_a = z_{.01} = 2.33$ , so  $\beta(30500) = \Phi\left(2.33 + \frac{30000 30500}{1500 / \sqrt{16}}\right) = \Phi(1.00) = .8413$ .

c. 
$$z_{\alpha} = z_{.01} = 2.33$$
 and  $z_{\beta} = z_{.05} = 1.645$ . Hence,  $n = \left[\frac{1500(2.33 + 1.645)}{30,000 - 30,500}\right]^2 = 142.2$ , so use  $n = 143$ .

d. From (a), the *P*-value is .0052. Hence, the smallest  $\alpha$  at which  $H_0$  can be rejected is .0052.

- 19.
- a. Since the alternative hypothesis is two-sided, *P*-value =  $2 \cdot \left[ 1 \Phi \left( \frac{|94.32 95|}{1.20 / \sqrt{16}} \right) \right] = 2 \cdot [1 \Phi(2.27)] =$ 
  - 2(.0116) = .0232. Since  $.0232 > \alpha = .01$ , we do not reject  $H_0$  at the .01 significance level.

**b.** 
$$z_{\alpha/2} = z_{.005} = 2.58$$
, so  $\beta(94) = \Phi\left(2.58 + \frac{95 - 94}{1.20 / \sqrt{16}}\right) - \Phi\left(-2.58 + \frac{95 - 94}{1.20 / \sqrt{16}}\right) = \Phi(5.91) - \Phi(0.75) = .2266$ .

c. 
$$z_{\beta} = z_{.1} = 1.28$$
. Hence,  $n = \left[\frac{1.20(2.58 + 1.28)}{95 - 94}\right]^2 = 21.46$ , so use  $n = 22$ .

21. The hypotheses are  $H_0: \mu = 5.5$  v.  $H_a: \mu \neq 5.5$ . a. The *P*-value is  $2 \cdot \left[ 1 - \Phi \left( \left| \frac{5.25 - 5.5}{.3/\sqrt{16}} \right| \right) \right] = 2 \cdot [1 - \Phi(3.33)] = .0008$ . Since the *P*-value is smaller than

any reasonable significance level (.1, .05, .01, .001), we reject  $H_0$ .

**b.** The chance of detecting that  $H_0$  is false is the complement of the chance of a type II error. With  $z_{a/2} =$ 

$$z_{.005} = 2.58, \ 1 - \beta \left( 5.6 \right) = 1 - \left[ \Phi \left( 2.58 + \frac{5.5 - 5.6}{.3/\sqrt{16}} \right) - \Phi \left( -2.58 + \frac{5.5 - 5.6}{.3/\sqrt{16}} \right) \right] = 1 - \Phi (1.25) + \Phi (3.91) = 0$$

.1056.

c. 
$$n = \left[\frac{.3(2.58 + 2.33)}{5.5 - 5.6}\right]^2 = 216.97$$
, so use  $n = 217$ .

23.

- **a.** Using software,  $\bar{x} = 0.75$ ,  $\tilde{x} = 0.64$ , s = .3025,  $f_s = 0.48$ . These summary statistics, as well as a box plot (not shown) indicate substantial positive skewness, but no outliers.
- **b.** No, it is not plausible from the results in part **a** that the variable ALD is normal. However, since n = 49, normality is not required for the use of z inference procedures.

c. We wish to test  $H_0$ :  $\mu = 1.0$  versus  $H_a$ :  $\mu < 1.0$ . The test statistic is  $z = \frac{0.75 - 1.0}{.3025 / \sqrt{49}} = -5.79$ , and so the *P*-value is  $P(Z \le -5.79) \approx 0$ . At any reasonable significance level, we reject the null hypothesis.

Therefore, yes, the data provides strong evidence that the true average ALD is less than 1.0.

**d.** 
$$\overline{x} + z_{.05} \frac{s}{\sqrt{n}} = 0.75 + 1.645 \frac{.3025}{\sqrt{49}} = 0.821$$

25. Let  $\mu$  denote the true average task time. The hypotheses of interest are  $H_0$ :  $\mu = 2$  v.  $H_a$ :  $\mu < 2$ . Using z-based inference with the data provided, the *P*-value of the test is  $P\left(Z \le \frac{1.95 - 2}{.20/\sqrt{52}}\right) = \Phi(-1.80) = .0359$ . Since

.0359 > .01, at the  $\alpha = .01$  significance level we do not reject  $H_0$ . At the .01 level, we do not have sufficient evidence to conclude that the <u>true</u> average task time is less than 2 seconds.

27. 
$$\beta(\mu_0 - \Delta) = \Phi(z_{\alpha/2} + \Delta\sqrt{n} / \sigma) - \Phi(-z_{\alpha/2} + \Delta\sqrt{n} / \sigma) = 1 - \Phi(-z_{\alpha/2} - \Delta\sqrt{n} / \sigma) - [1 - \Phi(z_{\alpha/2} - \Delta\sqrt{n} / \sigma)] = \Phi(z_{\alpha/2} - \Delta\sqrt{n} / \sigma) - \Phi(-z_{\alpha/2} - \Delta\sqrt{n} / \sigma) = \beta(\mu_0 + \Delta).$$

# Section 8.3

- 29. The hypotheses are  $H_0$ :  $\mu = .5$  versus  $H_a$ :  $\mu \neq .5$ . Since this is a two-sided test, we must double the one-tail area in each case to determine the *P*-value.
  - **a.**  $n = 13 \Rightarrow df = 13 1 = 12$ . Looking at column 12 of Table A.8, the area to the right of t = 1.6 is .068. Doubling this area gives the two-tailed *P*-value of 2(.068) = .134. Since  $.134 > \alpha = .05$ , we do not reject  $H_0$ .
  - **b.** For a two-sided test, observing t = -1.6 is equivalent to observing t = 1.6. So, again the *P*-value is 2(.068) = .134, and again we do not reject  $H_0$  at  $\alpha = .05$ .

- c. df = n 1 = 24; the area to the left of -2.6 = the area to the right of 2.6 = .008 according to Table A.8. Hence, the two-tailed *P*-value is 2(.008) = .016. Since .016 > .01, we do not reject  $H_0$  in this case.
- **d.** Similar to part (c), Table A.8 gives a one-tail area of .000 for  $t = \pm 3.9$  at df = 24. Hence, the two-tailed *P*-value is 2(.000) = .000, and we reject  $H_0$  at any reasonable  $\alpha$  level.
- 31. This is an upper-tailed test, so the *P*-value in each case is  $P(T \ge \text{observed } t)$ . **a.** *P*-value =  $P(T \ge 3.2 \text{ with } \text{df} = 14) = .003 \text{ according to Table A.8. Since } .003 \le .05$ , we reject  $H_0$ .
  - **b.** *P*-value =  $P(T \ge 1.8 \text{ with } df = 8) = .055$ . Since .055 > .01, do not reject  $H_0$ .
  - c. *P*-value =  $P(T \ge -.2 \text{ with } df = 23) = 1 P(T \ge .2 \text{ with } df = 23)$  by symmetry = 1 .422 = .578. Since .578 is quite large, we would not reject  $H_0$  at any reasonable  $\alpha$  level. (Note that the sign of the observed *t* statistic contradicts  $H_{as}$  so we know immediately not to reject  $H_0$ .)

#### 33.

- a. It appears that the true average weight could be significantly off from the production specification of 200 lb per pipe. Most of the boxplot is to the right of 200.
- **b.** Let  $\mu$  denote the true average weight of a 200 lb pipe. The appropriate null and alternative hypotheses are  $H_0$ :  $\mu = 200$  and  $H_a$ :  $\mu \neq 200$ . Since the data are reasonably normal, we will use a one-sample t procedure. Our test statistic is  $t = \frac{206.73 200}{6.35/\sqrt{30}} = \frac{6.73}{1.16} = 5.80$ , for a *P*-value of  $\approx 0$ . So, we reject  $H_0$ .

At the 5% significance level, the test appears to substantiate the statement in part a.

#### 35.

**a.** The hypotheses are  $H_0$ :  $\mu = 200$  versus  $H_a$ :  $\mu > 200$ . With the data provided,

 $t = \frac{\overline{x} - \mu_0}{s / \sqrt{n}} = \frac{249.7 - 200}{145.1 / \sqrt{12}} = 1.2$ ; at df = 12 - 1 = 11, *P*-value = .128. Since .128 > .05, *H*<sub>0</sub> is not rejected

at the a = .05 level. We have insufficient evidence to conclude that the true average repair time exceeds 200 minutes.

b. With 
$$d = \frac{|\mu_0 - \mu|}{\sigma} = \frac{|200 - 300|}{150} = 0.67$$
, df = 11, and  $\alpha = .05$ , software calculates power  $\approx .70$ , so  $\beta(300) \approx .30$ .

and the big has the second of the light in the second in mathematical of the first way

112

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

37.

a.

The accompanying normal probability plot is acceptably linear, which suggests that a normal population distribution is quite plausible.



**b.** The parameter of interest is  $\mu$  = the true average compression strength (MPa) for this type of concrete. The hypotheses are  $H_0$ :  $\mu = 100$  versus  $H_a$ :  $\mu < 100$ .

Since the data come from a plausibly normal population, we will use the t procedure. The test statistic

is  $t = \frac{\overline{x} - \mu_0}{s / \sqrt{n}} = \frac{96.42 - 100}{8.26 / \sqrt{10}} = -1.37$ . The corresponding one-tailed *P*-value, at df = 10 - 1 = 9, is

 $P(T \leq -1.37) \approx .102.$ 

The *P*-value slightly exceeds .10, the largest  $\alpha$  level we'd consider using in practice, so the null hypothesis  $H_0$ :  $\mu = 100$  should not be rejected. This concrete should be used.

**39.** Software provides  $\overline{x} = 1.243$  and s = 0.448 for this sample.

The parameter of interest is  $\mu$  = the population mean expense ratio (%) for large-cap growth mutual funds. The hypotheses are  $H_0$ :  $\mu = 1$  versus  $H_a$ :  $\mu > 1$ .

We have a random sample, and a normal probability plot is reasonably linear, so the assumptions for a *t* procedure are met.

The test statistic is  $t = \frac{1.243 - 1}{0.448 / \sqrt{20}} = 2.43$ , for a *P*-value of  $P(T \ge 2.43 \text{ at } df = 19) \approx .013$ . Hence, we

(barely) fail to reject  $H_0$  at the .01 significance level. There is insufficient evidence, at the  $\alpha = .01$  level, to conclude that the population mean expense ratio for large-cap growth mutual funds exceeds 1%.

b. A Type I error would be to incorrectly conclude that the population mean expense ratio for large-cap growth mutual funds exceeds 1% when, in fact the mean is 1%. A Type II error would be to fail to recognize that the population mean expense ratio for large-cap growth mutual funds exceeds 1% when that's actually true.

Since we failed to reject  $H_0$  in (a), we potentially committed a Type II error there. If we later find out that, in fact,  $\mu = 1.33$ , so  $H_a$  was actually true all along, then yes we have committed a Type II error.

c. With n = 20 so df = 19,  $d = \frac{1.33 - 1}{.5} = .66$ , and  $\alpha = .01$ , software provides power  $\approx .66$ . (Note: it's

purely a coincidence that power and d are the same decimal!) This means that if the true values of  $\mu$  and  $\sigma$  are  $\mu = 1.33$  and  $\sigma = .5$ , then there is a 66% probability of correctly rejecting  $H_0$ :  $\mu = 1$  in favor of  $H_n$ :  $\mu > 1$  at the .01 significance level based upon a sample of size n = 20.

41.

 $\mu = \text{true average reading}, H_0: \mu = 70 \text{ v. } H_a: \mu \neq 70 \text{ , and } t = \frac{\overline{x} - 70}{s/\sqrt{n}} = \frac{75.5 - 70}{7/\sqrt{6}} = \frac{5.5}{2.86} = 1.92 \text{ .}$ 

From table A.8, df = 5, *P*-value =  $2[P(T > 1.92)] \approx 2(.058) = .116$ . At significance level .05, there is **not** enough evidence to conclude that the spectrophotometer needs recalibrating.

## Section 8.4

43.

a. The parameter of interest is p = the proportion of the population of female workers that have BMIs of at least 30 (and, hence, are obese). The hypotheses are H<sub>0</sub>: p = .20 versus H<sub>a</sub>: p > .20. With n = 541, np<sub>0</sub> = 541(.2) = 108.2 ≥ 10 and n(1 - p<sub>0</sub>) = 541(.8) = 432.8 ≥ 10, so the "large-sample" z procedure is applicable.

From the data provided,  $\hat{p} = \frac{120}{541} = .2218$ , so  $z = \frac{\hat{p} - p_0}{\sqrt{p_0(1 - p_0)/n}} = \frac{.2218 - .20}{\sqrt{.20(.80)/541}} = 1.27$  and *P*-value

 $= P(Z \ge 1.27) = 1 - \Phi(1.27) = .1020$ . Since .1020 > .05, we fail to reject  $H_0$  at the  $\alpha = .05$  level. We do not have sufficient evidence to conclude that more than 20% of the population of female workers is obese.

- **b.** A Type I error would be to incorrectly conclude that more than 20% of the population of female workers is obese, when the true percentage is 20%. A Type II error would be to fail to recognize that more than 20% of the population of female workers is obese when that's actually true.
- c. The question is asking for the chance of committing a Type II error when the true value of p is .25, i.e. β(.25). Using the textbook formula,

$$\beta(.25) = \Phi \left| \frac{.20 - .25 + 1.645 \sqrt{.20(.80) / 541}}{\sqrt{.25(.75) / 541}} \right| = \Phi(-1.166) \approx .121.$$

45.

Let p = true proportion of all donors with type A blood. The hypotheses are  $H_0$ : p = .40 versus  $H_a$ :  $p \neq .40$ . Using the one-proportion z procedure, the test statistic is  $z = \frac{82/150 - .40}{\sqrt{.40(.60)/150}} = \frac{.147}{.04} = 3.667$ , and the

corresponding *P*-value is  $2P(Z \ge 3.667) \approx 0$ . Hence, we reject  $H_0$ . The data does suggest that the percentage of all donors with type A blood differs from 40%. (at the .01 significance level). Since the *P*-value is also less than .05, the conclusion would not change.

- 47.
  - a. The parameter of interest is p = the proportion of all wine customers who would find screw tops acceptable. The hypotheses are  $H_0$ : p = .25 versus  $H_a$ : p < .25.
    - With n = 106,  $np_0 = 106(.25) = 26.5 \ge 10$  and  $n(1 p_0) = 106(.75) = 79.5 \ge 10$ , so the "large-sample" x procedure is applicable.

From the data provided,  $\hat{p} = \frac{22}{106} = .208$ , so  $z = \frac{.208 - .25}{\sqrt{.25(.75)/106}} = -1.01$  and *P*-value =  $P(Z \le -1.01) =$ 

 $\Phi(-1.01) = .1562.$ 

Since .1562 > .10, we fail to reject  $H_0$  at the  $\alpha = .10$  level. We do not have sufficient evidence to suggest that less than 25% of all customers find screw tops acceptable. Therefore, we recommend that the winery should switch to screw tops.

**b.** A Type I error would be to incorrectly conclude that less than 25% of all customers find screw tops acceptable, when the true percentage is 25%. Hence, we'd recommend not switching to screw tops when there use is actually justified. A Type II error would be to fail to recognize that less than 25% of all customers find screw tops acceptable when that's actually true. Hence, we'd recommend (as we did in (a)) that the winery switch to screw tops when the switch is not justified. Since we failed to reject  $H_0$  in (a), we may have committed a Type II error.

49.

a. Let p = true proportion of current customers who qualify. The hypotheses are  $H_0$ : p = .05 v.  $H_a$ :  $p \neq .05$ . The test statistic is  $z = \frac{.08 - .05}{\sqrt{.05(.95)/n}} = 3.07$ , and the *P*-value is  $2 \cdot P(Z \ge 3.07) = 2(.0011) = .0022$ .

Since  $.0022 \le \alpha = .01$ ,  $H_0$  is rejected. The company's premise is not correct.

b. 
$$\beta(.10) = \Phi\left[\frac{.05 - .10 + 2.58\sqrt{.05(.95)/500}}{\sqrt{.10(.90)/500}}\right] - \Phi\left[\frac{.05 - .10 - 2.58\sqrt{.05(.95)/500}}{\sqrt{.10(.90)/500}}\right]$$
  
 $\approx \Phi(-1.85) - 0 = .0332$ 

51. The hypotheses are  $H_0: p = .10 \text{ v}$ .  $H_a: p > .10$ , and we reject  $H_0$  iff  $X \ge c$  for some unknown c. The corresponding chance of a type I error is  $\alpha = P(X \ge c \text{ when } p = .10) = 1 - B(c - 1; 10, .1)$ , since the rv X has a Binomial(10, .1) distribution when  $H_0$  is true.

The values n = 10, c = 3 yield  $\alpha = 1 - B(2; 10, .1) = .07$ , while  $\alpha > .10$  for c = 0, 1, 2. Thus c = 3 is the best choice to achieve  $\alpha \le .10$  and simultaneously minimize  $\beta$ . However,  $\beta(.3) = P(X < c \text{ when } p = .3) = B(2; 10, .3) = .383$ , which has been deemed too high. So, the desired  $\alpha$  and  $\beta$  levels cannot be achieved with a sample size of just n = 10.

The values n = 20, c = 5 yield  $\alpha = 1 - B(4; 20, .1) = .043$ , but again  $\beta(.3) = B(4; 20, .3) = .238$  is too high. The values n = 25, c = 5 yield  $\alpha = 1 - B(4; 25, .1) = .098$  while  $\beta(.3) = B(4; 25, .3) = .090 \le .10$ , so n = 25 should be used. In that case and with the rule that we reject  $H_0$  iff  $X \ge 5$ ,  $\alpha = .098$  and  $\beta(.3) = .090$ .

#### Section 8.5

53.

a. The formula for  $\beta$  is  $1 - \Phi\left(-2.33 + \frac{\sqrt{n}}{9}\right)$ , which gives .8888 for n = 100, .1587 for n = 900, and .0006

for n = 2500.

- **b.** Z = -5.3, which is "off the z table," so P-value < .0002; this value of z is quite statistically significant.
- c. No. Even when the departure from  $H_0$  is insignificant from a practical point of view, a statistically significant result is highly likely to appear; the test is too likely to detect small departures from  $H_0$ .

- a. The chance of committing a type I error on a single test is .01. Hence, the chance of committing at least one type I error among *m* tests is  $P(\text{at least on error}) = 1 P(\text{no type I errors}) = 1 [P(\text{no type I} error)]^m$  by independence =  $1 .99^m$ . For m = 5, the probability is .049; for m = 10, it's .096.
- b. Set the answer from (a) to .5 and solve for m: 1 − .99<sup>m</sup> ≥ .5 ⇒ .99<sup>m</sup> ≤ .5 ⇒ m ≥ log(.5)/log(.99) = 68.97. So, at least 69 tests must be run at the α = .01 level to have a 50-50 chance of committing at least one type I error.

# Supplementary Exercises

57. Because n = 50 is large, we use a z test here. The hypotheses are  $H_0$ :  $\mu = 3.2$  versus  $H_a$ :  $\mu \neq 3.2$ . The computed z value is  $z = \frac{3.05 - 3.20}{.34/\sqrt{50}} = -3.12$ , and the *P*-value is  $2P(Z \ge |-3.12|) = 2(.0009) = 0018$ . Since .0018 < .05,  $H_0$  should be rejected in favor of  $H_a$ .

#### 59.

55.

- **a.**  $H_0: \mu = .85$  v.  $H_a: \mu \neq .85$
- **b.** With a *P*-value of .30, we would reject the null hypothesis at any reasonable significance level, which includes both .05 and .10.

#### 61.

- a. The parameter of interest is μ = the true average contamination level (Total Cu, in mg/kg) in this region. The hypotheses are H<sub>0</sub>: μ = 20 versus H<sub>a</sub>: μ > 20. Using a one-sample t procedure, with x̄ = 45.31 and SE(x̄) = 5.26, the test statistic is t = 45.31-20/5.26 = 3.86. That's a very large t-statistic; however, at df = 3 1 = 2, the P-value is P(T≥ 3.86) ≈ .03. (Using the tables with t = 3.9 gives a P-value of ≈ .02.) Since the P-value exceeds .01, we would fail to reject H<sub>0</sub> at the α = .01 level. This is quite surprising, given the large t-value (45.31 greatly exceeds 20), but it's a result of the very small n.
- **b.** We want the probability that we fail to reject  $H_0$  in part (a) when n = 3 and the true values of  $\mu$  and  $\sigma$  are  $\mu = 50$  and  $\sigma = 10$ , i.e.  $\beta(50)$ . Using software, we get  $\beta(50) \approx .57$ .
- 63. n=47,  $\bar{x}=215$  mg, s=235 mg, scope of values = 5 mg to 1,176 mg
  - a. No, the distribution does not appear to be normal. It appears to be skewed to the right, since 0 is less than one standard deviation below the mean. It is not necessary to assume normality if the sample size is large enough due to the central limit theorem. This sample size is large enough so we can conduct a hypothesis test about the mean.

**b.** The parameter of interest is  $\mu$  = true daily caffeine consumption of adult women, and the hypotheses are  $H_0$ :  $\mu = 200$  versus  $H_a$ :  $\mu > 200$ . The test statistic (using a z test) is  $z = \frac{215 - 200}{235 / \sqrt{47}} = .44$  with a

corresponding *P*-value of  $P(Z \ge .44) = 1 - \Phi(.44) = .33$ . We fail to reject  $H_0$ , because .33 > .10. The data do not provide convincing evidence that daily consumption of all adult women exceeds 200 mg.

65.

- **a.** From Table A.17, when  $\mu = 9.5$ , d = .625, and df = 9,  $\beta \approx .60$ . When  $\mu = 9.0$ , d = 1.25, and df = 9,  $\beta \approx .20$ .
- **b.** From Table A.17, when  $\beta = .25$  and d = .625,  $n \approx 28$ .

67.

**a.** With  $H_0: p = 1/75$  v.  $H_a: p \neq 1/75$ ,  $\hat{p} = \frac{16}{800} = .02$ ,  $z \frac{.02 - .01333}{\sqrt{\frac{.01333(.98667)}{800}}} = 1.645$ , and *P*-value = .10, we

fail to reject the null hypothesis at the  $\alpha = .05$  level. There is no significant evidence that the incidence rate among prisoners differs from that of the adult population.

The possible error we could have made is a type II.

- **b.** P-value =  $2[1 \Phi(1.645)] = 2[.05] = .10$ . Yes, since .10 < .20, we could reject  $H_0$ .
- 69. Even though the underlying distribution may not be normal, a z test can be used because n is large. The null hypothesis  $H_0$ :  $\mu = 3200$  should be rejected in favor of  $H_a$ :  $\mu < 3200$  if the P-value is less than .001.

The computed test statistic is  $z = \frac{3107 - 3200}{188 / \sqrt{45}} = -3.32$  and the *P*-value is  $\Phi(-3.32) = .0005 < .001$ , so  $H_0$ 

should be rejected at level .001.

71. We wish to test  $H_0$ :  $\mu = 4$  versus  $H_a$ :  $\mu > 4$  using the test statistic  $z = \frac{\overline{x} - 4}{\sqrt{4/n}}$ . For the given sample, n = 36

and 
$$\overline{x} = \frac{160}{36} = 4.444$$
, so  $z = \frac{4.444 - 4}{\sqrt{4/36}} = 1.33$ .

The *P*-value is  $P(Z \ge 1.33) = 1 - \Phi(1.33) = .0918$ . Since .0918 > .02,  $H_0$  should not be rejected at this level. We do not have significant evidence at the .02 level to conclude that the true mean of this Poisson process is greater than 4.

73. The parameter of interest is p = the proportion of all college students who have maintained lifetime abstinence from alcohol. The hypotheses are  $H_0$ : p = .1,  $H_a$ : p > .1. With n = 462,  $np_0 = 462(.1) = 46.2 \ge 10$   $n(1 - p_0) = 462(.9) = 415.8 \ge 10$ , so the "large-sample" z procedure is applicable.

From the data provided,  $\hat{p} = \frac{51}{462} = .1104$ , so  $z = \frac{.1104 - .1}{\sqrt{.1(.9)/462}} = 0.74$ .

The corresponding one-tailed *P*-value is  $P(Z \ge 0.74) = 1 - \Phi(0.74) = .2296$ . Since .2296 > .05, we fail to reject  $H_0$  at the  $\alpha = .05$  level (and, in fact, at any reasonable significance level). The data does not give evidence to suggest that more than 10% of all college students have completely abstained from alcohol use.

117

75. Since *n* is large, we'll use the one-sample *z* procedure. With  $\mu$  = population mean Vitamin D level for infants, the hypotheses are  $H_0$ :  $\mu = 20 \text{ v}$ .  $H_a$ :  $\mu > 20$ . The test statistic is  $z = \frac{21-20}{11/\sqrt{102}} = 0.92$ , and the upper-tailed *P*-value is  $P(Z \ge 0.92) = .1788$ . Since .1788 > .10, we fail to reject  $H_0$ . It cannot be concluded that  $\mu > 20$ .

- 77. The 20 df row of Table A.7 shows that  $\chi^2_{.99,20} = 8.26 < 8.58$  ( $H_0$  not rejected at level .01) and  $8.58 < 9.591 = \chi^2_{.975,20}$  ( $H_0$  rejected at level .025). Thus .01 < P-value < .025, and  $H_0$  cannot be rejected at level .01 (the P-value is the smallest  $\alpha$  at which rejection can take place, and this exceeds .01).
- 79.
- **a.** When  $H_0$  is true,  $2\lambda_0 \Sigma X_i = \frac{2}{\mu_0} \sum X_i$  has a chi-squared distribution with df = 2n. If the alternative is

 $H_a: \mu < \mu_0$ , then we should reject  $H_0$  in favor of  $H_a$  when the sample mean  $\overline{x}$  is small. Since  $\overline{x}$  is small exactly when  $\sum x_i$  is small, we'll reject  $H_0$  when the test statistic is small. In particular, the *P*-value

should be the area to the <u>left</u> of the observed value  $\frac{2}{\mu_0}\sum x_i$ .

**b.** The hypotheses are  $H_0$ :  $\mu = 75$  versus  $H_a$ :  $\mu < 75$ . The test statistic value is  $\frac{2}{\mu_0} \sum x_i = \frac{2}{75}(737) =$ 

19.65. At df = 2(10) = 20, the *P*-value is the area to the left of 19.65 under the  $\chi^2_{20}$  curve. From software, this is about .52, so  $H_0$  clearly should not be rejected (the *P*-value is very large). The sample data do not suggest that true average lifetime is less than the previously claimed value.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

的复数形式 化合同分子 化合同合物 网络小学家 网络小学家 网络小学家

**CHAPTER 9** 

# Section 9.1

1.

3.

5.

- **a.**  $E(\overline{X} \overline{Y}) = E(\overline{X}) E(\overline{Y}) = 4.1 4.5 = -.4$ , irrespective of sample sizes.
- **b.**  $V(\overline{X} \overline{Y}) = V(\overline{X}) + V(\overline{Y}) = \frac{\sigma_1^2}{m} + \frac{\sigma_2^2}{n} = \frac{(1.8)^2}{100} + \frac{(2.0)^2}{100} = .0724$ , and the SD of  $\overline{X} \overline{Y}$  is  $\overline{X} \overline{Y} = \sqrt{.0724} = .2691$ .
  - c. A normal curve with mean and sd as given in a and b (because m = n = 100, the CLT implies that both  $\overline{X}$  and  $\overline{Y}$  have approximately normal distributions, so  $\overline{X} \overline{Y}$  does also). The shape is not necessarily that of a normal curve when m = n = 10, because the CLT cannot be invoked. So if the two lifetime population distributions are not normal, the distribution of  $\overline{X} \overline{Y}$  will typically be quite complicated.
- Let  $\mu_1$  = the population mean pain level under the control condition and  $\mu_2$  = the population mean pain level under the treatment condition.
  - **a.** The hypotheses of interest are  $H_0: \mu_1 \mu_2 = 0$  versus  $H_a: \mu_1 \mu_2 > 0$ . With the data provided, the test statistic value is  $z = \frac{(5.2 3.1) 0}{\sqrt{\frac{2.3^2}{43} + \frac{2.3^2}{43}}} = 4.23$ . The corresponding *P*-value is  $P(Z \ge 4.23) = 1 \Phi(4.23) \approx 0$ .

Hence, we reject  $H_0$  at the  $\alpha = .01$  level (in fact, at any reasonable level) and conclude that the average pain experienced under treatment is less than the average pain experienced under control.

**b.** Now the hypotheses are  $H_0: \mu_1 - \mu_2 = 1$  versus  $H_a: \mu_1 - \mu_2 > 1$ . The test statistic value is

 $z = \frac{(5.2-3.1)-1}{\sqrt{2.3^2 + \frac{2.3^2}{2.3^2}}} = 2.22$ , and the *P*-value is  $P(Z \ge 2.22) = 1 - \Phi(2.22) = .0132$ . Thus we would reject

 $H_0$  at the  $\alpha = .05$  level and conclude that mean pain under control condition exceeds that of treatment condition by more than 1 point. However, we would not reach the same decision at the  $\alpha = .01$  level (because .0132  $\leq$  .05 but .0132 > .01).

a.  $H_a$  says that the average calorie output for sufferers is more than 1 cal/cm<sup>2</sup>/min <u>below</u> that for non-

sufferers.  $\sqrt{\frac{\sigma_1^2}{m} + \frac{\sigma_2^2}{n}} = \sqrt{\frac{(.2)^2}{10} + \frac{(.4)^2}{10}} = .1414$ , so  $z = \frac{(.64 - 2.05) - (-1)}{.1414} = -2.90$ . The *P*-value for this one-sided test is  $P(Z \le -2.90) = .0019 < .01$ . So, at level .01,  $H_0$  is rejected.

**b.** 
$$z_a = z_{.01} = 2.33$$
, and so  $\beta(-1.2) = 1 - \Phi\left(-2.33 - \frac{-1.2 + 1}{.1414}\right) = 1 - \Phi\left(-.92\right) = .8212.$ 

c. 
$$m = n = \frac{.2(2.33 + 1.28)^2}{(-.2)^2} = 65.15$$
, so use 66.

Let  $\mu_1$  denote the true mean course GPA for all courses taught by full-time faculty, and let  $\mu_2$  denote the true mean course GPA for all courses taught by part-time faculty. The hypotheses of interest are  $H_0$ :  $\mu_1 = \mu_2$  versus  $H_a$ :  $\mu_1 \neq \mu_2$ ; or, equivalently,  $H_0$ :  $\mu_1 - \mu_2 = 0$  v.  $H_a$ :  $\mu_1 - \mu_2 \neq 0$ .

The large-sample test statistic is 
$$z = \frac{(\overline{x} - \overline{y}) - \Delta_0}{\sqrt{\frac{s_1^2}{m} + \frac{s_2^2}{n}}} = \frac{(2.7186 - 2.8639) - 0}{\sqrt{\frac{(.63342)^2}{125} + \frac{(.49241)^2}{88}}} = -1.88$$
. The corresponding

two-tailed *P*-value is  $P(|Z| \ge |-1.88|) = 2[1 - \Phi(1.88)] = .0602$ . Since the *P*-value exceeds  $\alpha = .01$ , we fail to reject  $H_0$ . At the .01 significance level, there is insufficient evidence to conclude that the true mean course GPAs differ for these two populations of faculty.

9.

7.

**a.** Point estimate  $\overline{x} - \overline{y} = 19.9 - 13.7 = 6.2$ . It appears that there could be a difference.

**b.** 
$$H_0: \mu_1 - \mu_2 = 0, H_a: \mu_1 - \mu_2 \neq 0, z = \frac{(19.9 - 13.7)}{\sqrt{\frac{39.1^2}{60} + \frac{15.8^2}{60}}} = \frac{6.2}{5.44} = 1.14$$
, and the *P*-value = 2[*P*(*Z* > 1.14)] =

2(.1271) = .2542. The *P*-value is larger than any reasonable  $\alpha$ , so we do not reject  $H_0$ . There is no statistically significant difference.

- c. No. With a normal distribution, we would expect most of the data to be within 2 standard deviations of the mean, and the distribution should be symmetric. Two sd's above the mean is 98.1, but the distribution stops at zero on the left. The distribution is positively skewed.
- **d.** We will calculate a 95% confidence interval for  $\mu$ , the true average length of stays for patients given the treatment.  $19.9 \pm 1.96 \frac{39.1}{\sqrt{60}} = 19.9 \pm 9.9 = (10.0, 21.8).$
- 11.
- $(\overline{x}-\overline{y})\pm z_{\alpha/2}\sqrt{\frac{s_1^2}{m}+\frac{s_2^2}{n}} = (\overline{x}-\overline{y})\pm z_{\alpha/2}\sqrt{(SE_1)^2+(SE_2)^2}$ . Using  $\alpha = .05$  and  $z_{\alpha/2} = 1.96$  yields  $(5.5-3.8)\pm 1.96\sqrt{(0.3)^2+(0.2)^2} = (0.99, 2.41)$ . We are 95% confident that the true average blood lead level for male workers is between 0.99 and 2.41 higher than the corresponding average for female workers.
- 13.  $\sigma_1 = \sigma_2 = .05, d = .04, \alpha = .01, \beta = .05, \text{ and the test is one-tailed} \Rightarrow$  $n = \frac{(.0025 + .0025)(2.33 + 1.645)^2}{.0016} = 49.38, \text{ so use } n = 50.$

15.

**a.** As either *m* or *n* increases, *SD* decreases, so  $\frac{\mu_1 - \mu_2 - \Delta_0}{SD}$  increases (the numerator is positive), so

$$\left(z_{\alpha} - \frac{\mu_{1} - \mu_{2} - \Delta_{0}}{SD}\right)$$
 decreases, so  $\beta = \Phi\left(z_{\alpha} - \frac{\mu_{1} - \mu_{2} - \Delta_{0}}{SD}\right)$  decreases.

**b.** As  $\beta$  decreases,  $z_{\beta}$  increases, and since  $z_{\beta}$  is the numerator of *n*, *n* increases also.

### Chapter 9: Inferences Based on Two Samples

# Section 9.2

17.

prior is a particular to a second normal probability picks we are the both of the line of the solitoness.

a. 
$$v = \frac{\left(5^2/10 + 6^2/10\right)^2}{\left(5^2/10\right)^2/9 + \left(6^2/10\right)^2/9} = \frac{37.21}{.694 + 1.44} = 17.43 \approx 17.$$
  
b.  $v = \frac{\left(5^2/10 + 6^2/15\right)^2}{\left(5^2/10\right)^2/9 + \left(6^2/15\right)^2/14} = \frac{24.01}{.694 + .411} = 21.7 \approx 21.$   
c.  $v = \frac{\left(2^2/10 + 6^2/15\right)^2}{\left(2^2/10\right)^2/9 + \left(6^2/15\right)^2/14} = \frac{7.84}{.018 + .411} = 18.27 \approx 18.$   
d.  $v = \frac{\left(5^2/12 + 6^2/24\right)^2}{\left(5^2/12\right)^2/11 + \left(6^2/24\right)^2/23} = \frac{12.84}{.395 + .098} = 26.05 \approx 26.$ 

19. For the given hypotheses, the test statistic is  $t = \frac{115.7 - 129.3 + 10}{\sqrt{\frac{5.03^2}{6} + \frac{5.38^2}{6}}} = \frac{-3.6}{3.007} = -1.20$ , and the df is

$$\nu = \frac{\left(4.2168 + 4.8241\right)^2}{\left(\frac{4.2168}{5}\right)^2} + \frac{\left(4.8241\right)^2}{5} = 9.96$$
, so use df = 9. The *P*-value is  $P(T \le -1.20$  when  $T \sim t_9) \approx .130$ .

Since .130 > .01, we don't reject  $H_0$ .

21. Let  $\mu_1$  = the true average gap detection threshold for normal subjects, and  $\mu_2$  = the corresponding value for CTS subjects. The relevant hypotheses are  $H_0$ :  $\mu_1 - \mu_2 = 0$  v.  $H_a$ :  $\mu_1 - \mu_2 < 0$ , and the test statistic is

CTS subjects. The relevant hypotheses are  $n_0, p_1 = p_2$  $t = \frac{1.71 - 2.53}{\sqrt{.0351125 + .07569}} = \frac{-.82}{.3329} = -2.46$ . Using df  $v = \frac{(.0351125 + .07569)^2}{(.0351125)^2} = 15.1$ , or 15, the *P*-value

is  $P(T \le -2.46$  when  $T \sim t_{15}) \approx .013$ . Since .013 > .01, we fail to reject  $H_0$  at the  $\alpha = .01$  level. We have insufficient evidence to claim that the true average gap detection threshold for CTS subjects exceeds that for normal subjects.

23.

a.

Using Minitab to generate normal probability plots, we see that both plots illustrate sufficient linearity. Therefore, it is plausible that both samples have been selected from normal population distributions.



b. The comparative boxplot does not suggest a difference between average extensibility for the two types of fabrics.



Comparative Box Plot for High Quality and Poor Quality Fabric

c. We test  $H_0: \mu_1 - \mu_2 = 0$  v.  $H_a: \mu_1 - \mu_2 \neq 0$ . With degrees of freedom  $\nu = \frac{(.0433265)^2}{.00017906} = 10.5$ 

(which we round down to 10) and test statistic is  $t = \frac{-.08}{\sqrt{(.0433265)}} = -.38 \approx -0.4$ , the *P*-value is

2(.349) = .698. Since the *P*-value is very large, we do not reject  $H_0$ . There is insufficient evidence to claim that the true average extensibility differs for the two types of fabrics.

## Chapter 9: Inferences Based on Two Samples

#### 25.

- **a.** Normal probability plots of both samples (not shown) exhibit substantial linear patterns, suggesting that the normality assumption is reasonable for both populations of prices.
  - **b.** The comparative boxplots below suggest that the average price for a wine earning  $a \ge 93$  rating is much higher than the average price earning  $a \le 89$  rating.



c. From the data provided,  $\bar{x} = 110.8$ ,  $\bar{y} = 61.7$ ,  $s_1 = 48.7$ ,  $s_2 = 23.8$ , and  $v \approx 15$ . The resulting 95% CI for the difference of population means is  $(110.8 - 61.7) \pm t_{.025,15} \sqrt{\frac{48.7^2}{12} + \frac{23.8^2}{14}} = (16.1, 82.0)$ . That is, we

are 95% confident that wines rated  $\geq$  93 cost, on average, between \$16.10 and \$82.00 more than wines rated  $\leq$  89. Since the CI does not include 0, this certainly contradicts the claim that price and quality are unrelated.

#### 27.

a. Let's construct a 99% CI for μ<sub>AN</sub>, the true mean intermuscular adipose tissue (IAT) under the described AN protocol. Assuming the data comes from a normal population, the CI is given by

$$\overline{x} \pm t_{\alpha/2,n-1} \frac{s}{\sqrt{n}} = .52 \pm t_{.005,15} \frac{.26}{\sqrt{16}} = .52 \pm 2.947 \frac{.26}{\sqrt{16}} = (.33, .71)$$
. We are 99% confident that the true mean

IAT under the AN protocol is between .33 kg and .71 kg.

**b.** Let's construct a 99% CI for  $\mu_{AN} - \mu_C$ , the difference between true mean AN IAT and true mean control IAT. Assuming the data come from normal populations, the CI is given by

$$(\overline{x} - \overline{y}) \pm t_{\alpha/2,\nu} \sqrt{\frac{s_1^2}{m} + \frac{s_2^2}{n}} = (.52 - .35) \pm t_{.005,21} \sqrt{\frac{(.26)^2}{16} + \frac{(.15)^2}{8}} = .17 \pm 2.831 \sqrt{\frac{(.26)^2}{16} + \frac{(.15)^2}{8}} = (-.07, .41).$$

Since this CI includes zero, it's plausible that the difference between the two true means is zero (i.e.,  $\mu_{AN} - \mu_C = 0$ ). [Note: the df calculation  $\nu = 21$  comes from applying the formula in the textbook.]

29. Let  $\mu_1$  = the true average compression strength for strawberry drink and let  $\mu_2$  = the true average compression strength for cola. A lower tailed test is appropriate. We test  $H_0$ :  $\mu_1 - \mu_2 = 0$  v.  $H_a$ :  $\mu_1 - \mu_2 < 0$ .

The test statistic is 
$$t = \frac{-14}{\sqrt{29.4 + 15}} = -2.10$$
;  $v = \frac{(44.4)^2}{(\frac{29.4}{14})^2} + \frac{(15)^2}{14} = \frac{1971.36}{77.8114} = 25.3$ , so use df=25.

The *P*-value  $\approx P(t < -2.10) = .023$ . This *P*-value indicates strong support for the alternative hypothesis. The data does suggest that the extra carbonation of cola results in a higher average compression strength.

#### 31.

a. The most notable feature of these boxplots is the larger amount of variation present in the mid-range data compared to the high-range data. Otherwise, both look reasonably symmetric with no outliers present.



**b.** Using df = 23, a 95% confidence interval for  $\mu_{\text{mid-range}} - \mu_{\text{high-range}}$  is

$$(38.3-437.45) \pm 2.069\sqrt{\frac{15.1^2}{17}} + \frac{6.83^2}{11} = .85 \pm 8.69 = (-7.84, 9.54)$$
. Since plausible values for

 $\mu_{\text{mid-range}} - \mu_{\text{high-range}}$  are both positive and negative (i.e., the interval spans zero) we would conclude that there is not sufficient evidence to suggest that the average value for mid-range and the average value for high-range differ.

33.

Let  $\mu_1$  and  $\mu_2$  represent the true mean body mass decrease for the vegan diet and the control diet, respectively. We wish to test the hypotheses  $H_0$ :  $\mu_1 - \mu_2 \le 1$  v.  $H_a$ :  $\mu_1 - \mu_2 \ge 1$ . The relevant test statistic is

 $t = \frac{(5.8-3.8)-1}{\sqrt{3.2^2+2.8^2}} = 1.33$ , with estimated df = 60 using the formula. Rounding to t = 1.3, Table A.8 gives a

(4

one-sided *P*-value of .098 (a computer will give the more accurate *P*-value of .094). Since our *P*-value >  $\alpha = .05$ , we fail to reject  $H_0$  at the 5% level. We do not have statistically significant evidence that the true average weight loss for the vegan diet exceeds the true average weight loss for the control diet by more than 1 kg.

35.

There are two changes that must be made to the procedure we currently use. First, the equation used to

compute the value of the *t* test statistic is:  $t = \frac{(\overline{x} - \overline{y}) - \Delta}{s_p \sqrt{\frac{1}{m} + \frac{1}{n}}}$  where  $s_p$  is defined as in Exercise 34. Second,

the degrees of freedom = m + n - 2. Assuming equal variances in the situation from Exercise 33, we

- calculate  $s_p$  as follows:  $s_p = \sqrt{\left(\frac{7}{16}\right)\left(2.6\right)^2 + \left(\frac{9}{16}\right)\left(2.5\right)^2} = 2.544$ . The value of the test statistic is,
- then,  $t = \frac{(32.8 40.5) (-5)}{2.544\sqrt{\frac{1}{8} + \frac{1}{10}}} = -2.24 \approx -2.2$  with df = 16, and the *P*-value is P(T < -2.2) = .021. Since
- .021 > .01, we fail to reject  $H_0$ .

### Chapter 9: Inferences Based on Two Samples

# Section 9.3

- 37.
- a. This exercise calls for paired analysis. First, compute the difference between indoor and outdoor concentrations of hexavalent chromium for each of the 33 houses. These 33 differences are summarized as follows: n = 33,  $\overline{d} = -.4239$ ,  $s_D = .3868$ , where d = (indoor value outdoor value). Then  $t_{.025,32} = 2.037$ , and a 95% confidence interval for the population mean difference between indoor

and outdoor concentration is  $-.4239 \pm (2.037) \left( \frac{.3868}{\sqrt{33}} \right) = -.4239 \pm .13715 = (-.5611, -.2868)$ . We can

be highly confident, at the 95% confidence level, that the true average concentration of hexavalent chromium outdoors exceeds the true average concentration indoors by between .2868 and .5611 nanograms/m<sup>3</sup>.

**b.** A 95% prediction interval for the difference in concentration for the 34<sup>th</sup> house is  $\overline{d} \pm t_{.025,32} \left( s_D \sqrt{1 + \frac{1}{n}} \right) = -.4239 \pm (2.037) \left( .3868 \sqrt{1 + \frac{1}{33}} \right) = (-1.224, .3758)$ . This prediction interval means that the indoor concentration may exceed the outdoor concentration by as much as .3758 nanograms/m<sup>3</sup> and that the outdoor concentration may exceed the indoor concentration by a much as 1.224 nanograms/m<sup>3</sup>, for the 34<sup>th</sup> house. Clearly, this is a wide prediction interval, largely because of the amount of variation in the differences.

39.

The accompanying normal probability plot shows that the <u>differences</u> are consistent with a normal population distribution.



**b.** We want to test  $H_0: \mu_D = 0$  versus  $H_a: \mu_D \neq 0$ . The test statistic is  $t = \frac{\overline{d} - 0}{s_D / \sqrt{n}} = \frac{167.2 - 0}{228 / \sqrt{14}} = 2.74$ , and

the two-tailed *P*-value is given by  $2[P(T > 2.74)] \approx 2[P(T > 2.7)] = 2[.009] = .018$ . Since .018 < .05, we reject  $H_0$ . There is evidence to support the claim that the true average difference between intake values measured by the two methods is <u>not</u> 0.

41.

a. Let  $\mu_D$  denote the true mean change in total cholesterol under the aripiprazole regimen. A 95% CI for

 $\mu_D$ , using the "large-sample" method, is  $\overline{d} \pm z_{\alpha/2} \frac{s_D}{\sqrt{n}} = 3.75 \pm 1.96(3.878) = (-3.85, 11.35).$ 

**b.** Now let  $\mu_D$  denote the true mean change in total cholesterol under the quetiapine regimen. The hypotheses are  $H_0$ :  $\mu_D = 0$  versus  $H_a$ :  $\mu_D > 0$ . Assuming the distribution of cholesterol changes under this regimen is normal, we may apply a paired *t* test:

$$t = \frac{d - \Delta_0}{s_p / \sqrt{n}} = \frac{9.05 - 0}{4.256} = 2.126 \Rightarrow P \text{-value} = P(T_{35} \ge 2.126) \approx P(T_{35} \ge 2.1) = .02.$$

Our conclusion depends on our significance level. At the  $\alpha = .05$  level, there is evidence that the true mean change in total cholesterol under the quetiapine regimen is positive (i.e., there's been an increase); however, we do not have sufficient evidence to draw that conclusion at the  $\alpha = .01$  level.

- c. Using the "large-sample" procedure again, the 95% CI is  $\overline{d} \pm 1.96 \frac{s_D}{\sqrt{n}} = \overline{d} \pm 1.96 SE(\overline{d})$ . If this equals
  - (7.38, 9.69), then midpoint =  $\overline{d}$  = 8.535 and width = 2(1.96 SE( $\overline{d}$ )) = 9.69 7.38 = 2.31  $\Rightarrow$
  - $SE(\overline{d}) = \frac{2.31}{2(1.96)} = .59$ . Now, use these values to construct a 99% CI (again, using a "large-sample" z method):  $\overline{d} \pm 2.576SE(\overline{d}) = 8.535 \pm 2.576(.59) = 8.535 \pm 1.52 = (7.02, 10.06)$ .
- 43.

**a.** Although there is a "jump" in the middle of the Normal Probability plot, the data follow a reasonably straight path, so there is no strong reason for doubting the normality of the population of differences.

b. A 95% lower confidence bound for the population mean difference is:

$$\overline{d} - t_{.05,14} \left(\frac{s_d}{\sqrt{n}}\right) = -38.60 - (1.761) \left(\frac{23.18}{\sqrt{15}}\right) = -38.60 - 10.54 = -49.14$$
. We are 95% confident that the

true mean difference between age at onset of Cushing's disease symptoms and age at diagnosis is greater than -49.14.

- c. A 95% upper confidence bound for the population mean difference is 38.60 + 10.54 = 49.14.
- 45.
- **a.** Yes, it's quite plausible that the population distribution of differences is normal, since the accompanying normal probability plot of the differences is quite linear.



www.konkur.in

#### Chapter 9: Inferences Based on Two Samples

**b.** No. Since the data is paired, the sample means and standard deviations are not useful summaries for inference. Those statistics would only be useful if we were analyzing two <u>independent</u> samples of data. (We could deduce  $\overline{d}$  by subtracting the sample means, but there's no way we could deduce  $s_D$  from the separate sample standard deviations.)

c. The hypotheses corresponding to an upper-tailed test are  $H_0$ :  $\mu_D = 0$  versus  $H_a$ :  $\mu_D > 0$ . From the data

provided, the paired t test statistic is  $t = \frac{\overline{d} - \Delta_0}{s_D / \sqrt{n}} = \frac{82.5 - 0}{87.4 / \sqrt{15}} = 3.66$ . The corresponding P-value is

 $P(T_{14} \ge 3.66) \approx P(T_{14} \ge 3.7) = .001$ . While the *P*-value stated in the article is inaccurate, the conclusion remains the same: we have strong evidence to suggest that the mean difference in ER velocity and IR velocity is positive. Since the measurements were <u>negative</u> (e.g. -130.6 deg/sec and -98.9 deg/sec), this actually means that the magnitude of IR velocity is significantly higher, on average, than the magnitude of ER velocity, as the authors of the article concluded.

#### 47. From the data, n = 12, $\overline{d} = -0.73$ , $s_D = 2.81$ .

**a.** Let  $\mu_D$  = the true mean difference in strength between curing under moist conditions and laboratory drying conditions. A 95% CI for  $\mu_D$  is  $\overline{d} \pm t_{.025,11}s_D/\sqrt{n} = -0.73 \pm 2.201(2.81)/\sqrt{10} = (-2.52 \text{ MPa}, 1.05 \text{ MPa})$ . In particular, this interval estimate includes the value zero, suggesting that true mean strength is not significantly different under these two conditions.

**b.** Since n = 12, we must check that the <u>differences</u> are plausibly from a normal population. The normal probability plot below strongly substantiates that condition.



### Section 9.4

Let  $p_1$  denote the true proportion of correct responses to the first question; define  $p_2$  similarly. The hypotheses of interest are  $H_0$ :  $p_1 - p_2 = 0$  versus  $H_a$ :  $p_1 - p_2 > 0$ . Summary statistics are  $n_1 = n_2 = 200$ ,

 $\hat{p}_1 = \frac{164}{200} = .82$ ,  $\hat{p}_2 = \frac{140}{200} = .70$ , and the pooled proportion is  $\hat{p} = .76$ . Since the sample sizes are large, we

may apply the two-proportion z test procedure.

The calculated test statistic is 
$$z = \frac{(.82 - .70) - 0}{\sqrt{(.76)(.24)\left[\frac{1}{200} + \frac{1}{200}\right]}} = 2.81$$
, and the *P*-value is  $P(Z \ge 2.81) = .0025$ .

Since  $.0025 \le .05$ , we reject  $H_0$  at the  $\alpha = .05$  level and conclude that, indeed, the true proportion of correct answers to the context-free question is higher than the proportion of right answers to the contextual one.

51. Let  $p_1$  = the true proportion of patients that will experience erectile dysfunction when given no counseling, and define  $p_2$  similarly for patients receiving counseling about this possible side effect. The hypotheses of interest are  $H_0$ :  $p_1 - p_2 = 0$  versus  $H_a$ :  $p_1 - p_2 < 0$ .

The actual data are 8 out of 52 for the first group and 24 out of 55 for the second group, for a pooled

proportion of  $\hat{p} = \frac{8+24}{52+55} = .299$ . The two-proportion z test statistic is  $\frac{(.153-.436)-0}{\sqrt{(.299)(.701)\left[\frac{1}{52}+\frac{1}{55}\right]}} = -3.20$ , and

the *P*-value is  $P(Z \le -3.20) = .0007$ . Since .0007 < .05, we reject  $H_0$  and conclude that a higher proportion of men will experience erectile dysfunction if told that it's a possible side effect of the BPH treatment, than if they weren't told of this potential side effect.

53.

p

b

**a.** Let  $p_1$  and  $p_2$  denote the true incidence rates of GI problems for the olestra and control groups, respectively. We wish to test  $H_0: p_1 - \mu_2 = 0$  v.  $H_a: p_1 - p_2 \neq 0$ . The pooled proportion is

$$=\frac{529(.176)+563(.158)}{529+563} = .1667$$
, from which the relevant test statistic is  $z =$ 

$$176 - .158 = 0.78$$
. The two-sided *P*-value is  $2P(Z \ge 0.78) = .433 > \alpha = .05$ ,

 $\sqrt{(.1667)(.8333)[529^{-1}+563^{-1}]}$ 

hence we fail to reject the null hypothesis. The data do not suggest a statistically significant difference between the incidence rates of GI problems between the two groups.

$$n = \frac{\left(1.96\sqrt{(.35)(1.65)/2} + 1.28\sqrt{(.15)(.85) + (.2)(.8)}\right)^2}{(.05)^2} = 1210.39$$
, so a common sample size of  $m = n = \frac{(.05)^2}{(.05)^2}$ 

1211 would be required.

55.

**a.** A 95% large sample confidence interval formula for  $\ln(\theta)$  is  $\ln(\hat{\theta}) \pm z_{\alpha/2} \sqrt{\frac{m-x}{mx} + \frac{n-y}{ny}}$ . Taking the antilogs of the upper and lower bounds gives the confidence interval for  $\theta$  itself.

<sup>49.</sup> 

#### Chapter 9: Inferences Based on Two Samples

$$\hat{\theta} = \frac{\frac{189}{11,034}}{\frac{104}{11,037}} = 1.818, \ \ln(\hat{\theta}) = .598, \text{ and the standard deviation is}$$

 $\sqrt{\frac{10,845}{(11,034)(189)}} + \frac{10,933}{(11,037)(104)} = .1213$ , so the CI for  $\ln(\theta)$  is  $.598 \pm 1.96(.1213) = (.360,.836)$ .

Then taking the antilogs of the two bounds gives the CI for  $\theta$  to be (1.43, 2.31). We are 95% confident that people who do not take the aspirin treatment are between 1.43 and 2.31 times more likely to suffer a heart attack than those who do. This suggests aspirin therapy may be effective in reducing the risk of a heart attack.

57. 
$$\hat{p}_1 = \frac{15+7}{40} = .550$$
,  $\hat{p}_2 = \frac{29}{42} = .690$ , and the 95% CI is  $(.550 - .690) \pm 1.96(.106) = -.14 \pm .21 = (-.35, .07)$ .

## Section 9.5

ł

#### 59.

- **a.** From Table A.9, column 5, row 8,  $F_{.01,5,8} = 3.69$ .
- **b.** From column 8, row 5,  $F_{.01,8,5} = 4.82$ .

c. 
$$F_{.95,5,8} = \frac{1}{F_{.05,8,5}} = .207$$
.

**d.** 
$$F_{.95,8,5} = \frac{1}{F_{.05,5,8}} = .271$$

**e.**  $F_{.01,10,12} = 4.30$ 

**f.** 
$$F_{.99,10,12} = \frac{1}{F_{.01,12,10}} = \frac{1}{4.71} = .212$$
.

g. 
$$F_{.05,6,4} = 6.16$$
, so  $P(F \le 6.16) = .95$ .

**h.** Since  $F_{.99,10,5} = \frac{1}{5.64} = .177$ ,  $P(.177 \le F \le 4.74) = P(F \le 4.74) - P(F \le .177) = .95 - .01 = .94$ .

61. We test 
$$H_0: \sigma_1^2 = \sigma_2^2$$
 v.  $H_a: \sigma_1^2 \neq \sigma_2^2$ . The calculated test statistic is  $f = \frac{(2.75)^2}{(4.44)^2} = .384$ . To use Table

A.9, take the reciprocal: 1/f = 2.61. With numerator df = m - 1 = 5 - 1 = 4 and denominator df = n - 1 = 10 - 1 = 9 after taking the reciprocal, Table A.9 indicates the one-tailed probability is slightly more than .10, and so the two-sided *P*-value is slightly more than 2(.10) = .20.

Since .20 > .10, we do not reject  $H_0$  at the  $\alpha = .1$  level and conclude that there is no significant difference between the two standard deviations.

63. Let  $\sigma_1^2$  = variance in weight gain for low-dose treatment, and  $\sigma_2^2$  = variance in weight gain for control condition. We wish to test  $H_0: \sigma_1^2 = \sigma_2^2$  v.  $H_a: \sigma_1^2 > \sigma_2^2$ . The test statistic is  $f = \frac{s_1^2}{s_2^2} = \frac{54^2}{32^2} = 2.85$ . From

Table A.9 with df =  $(19, 22) \approx (20, 22)$ , the *P*-value is approximately .01, and we reject  $H_0$  at level .05. The data do suggest that there is more variability in the low-dose weight gains.

65.

67.

 $P\left(F_{1-\alpha/2,m-1,n-1} \le \frac{S_1^2 / \sigma_1^2}{S_2^2 / \sigma_2^2} \le F_{\alpha/2,m-1,n-1}\right) = 1 - \alpha \text{. The set of inequalities inside the parentheses is clearly}$ 

equivalent to  $\frac{S_2^2 F_{1-\alpha/2,m-1,n-1}}{S_1^2} \le \frac{\sigma_2^2}{\sigma_1^2} \le \frac{S_2^2 F_{\alpha/2,m-1,n-1}}{S_1^2}$ . Substituting the sample values  $s_1^2$  and  $s_2^2$  yields the

confidence interval for  $\frac{\sigma_2^2}{\sigma_1^2}$ , and taking the square root of each endpoint yields the confidence interval for

 $\frac{\sigma_2}{\sigma_1}$ . With m = n = 4, we need  $F_{.05,3,3} = 9.28$  and  $F_{.95,3,3} = \frac{1}{9.28} = .108$ . Then with  $s_1 = .160$  and  $s_2 = .074$ , the CI for  $\frac{\sigma_2^2}{\sigma_1^2}$  is (.023, 1.99), and for  $\frac{\sigma_2}{\sigma_1}$  is (.15, 1.41).

# Supplementary Exercises

We test 
$$H_0: \mu_1 - \mu_2 = 0$$
 v.  $H_a: \mu_1 - \mu_2 \neq 0$ . The test statistic is  
 $t = \frac{(\overline{x} - \overline{y}) - \Delta}{\sqrt{\frac{s_1^2}{m} + \frac{s_2^2}{n}}} = \frac{807 - 757}{\sqrt{\frac{27^2}{10} + \frac{41^2}{10}}} = \frac{50}{\sqrt{241}} = \frac{50}{15.524} = 3.22$ . The approximate df is  
 $v = \frac{(241)^2}{\frac{(72.9)^2}{9} + \frac{(168.1)^2}{9}} = 15.6$ , which we round down to 15. The *P*-value for a two-tailed test is

approximately 2P(T > 3.22) = 2(.003) = .006. This small of a *P*-value gives strong support for the alternative hypothesis. The data indicates a significant difference. Due to the small sample sizes (10 each), we are assuming here that compression strengths for both fixed and floating test platens are normally distributed. And, as always, we are assuming the data were randomly sampled from their respective populations.

69. Let  $p_1$  = true proportion of returned questionnaires that included no incentive;  $p_2$  = true proportion of returned questionnaires that included an incentive. The hypotheses are  $H_0: p_1 - p_2 = 0$  v.  $H_a: p_1 - p_2 < 0$ .

The test statistic is 
$$z = \frac{\hat{p}_1 - \hat{p}_2}{\sqrt{\hat{p}\hat{q}\left(\frac{1}{m} + \frac{1}{n}\right)}}$$

 $\hat{p}_1 = \frac{75}{110} = .682$  and  $\hat{p}_2 = \frac{66}{98} = .673$ ; at this point, you might notice that since  $\hat{p}_1 > \hat{p}_2$ , the numerator of the

z statistic will be > 0, and since we have a lower tailed test, the *P*-value will be > .5. We fail to reject  $H_0$ . This data does not suggest that including an incentive increases the likelihood of a response.

#### Chapter 9: Inferences Based on Two Samples

- 71. The center of any confidence interval for  $\mu_1 \mu_2$  is always  $\overline{x}_1 \overline{x}_2$ , so  $\overline{x}_1 \overline{x}_2 = \frac{-473.3 + 1691.9}{2} = 609.3$ . Furthermore, half of the width of this interval is  $\frac{1691.9 - (-473.3)}{2} = 1082.6$ . Equating this value to the expression on the right of the 95% confidence interval formula, we find  $\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}} = \frac{1082.6}{1.96} = 552.35$ . For a 90% interval, the associated z value is 1.645, so the 90% confidence interval is then  $609.3 \pm (1.645)(552.35) = 609.3 \pm 908.6 = (-299.3, 1517.9)$ .
- 73. Let  $\mu_1$  and  $\mu_2$  denote the true mean zinc mass for Duracell and Energizer batteries, respectively. We want to test the hypotheses  $H_0$ :  $\mu_1 \mu_2 = 0$  versus  $H_a$ :  $\mu_1 \mu_2 \neq 0$ . Assuming that both zinc mass distributions are normal, we'll use a two-sample *t* test; the test statistic is  $t = \frac{(\overline{x} \overline{y}) \Delta_0}{\sqrt{\frac{s_1^2}{m} + \frac{s_2^2}{n}}} = \frac{(138.52 149.07) 0}{\sqrt{\frac{(7.76)^2}{15} + \frac{(1.52)^2}{20}}} = -5.19.$

The textbook's formula for df gives v = 14. The *P*-value is  $P(T_{14} \le -5.19) \approx 0$ . Hence, we strongly reject  $H_0$  and we conclude the mean zinc mass content for Duracell and Energizer batteries are <u>not</u> the same (they do differ).

75. Since we can assume that the distributions from which the samples were taken are normal, we use the twosample t test. Let  $\mu_1$  denote the true mean headability rating for aluminum killed steel specimens and  $\mu_2$ denote the true mean headability rating for silicon killed steel. Then the hypotheses are  $H_0: \mu_1 - \mu_2 = 0$  v.

$$H_{\rm a}: \mu_1 - \mu_2 \neq 0$$
. The test statistic is  $t = \frac{-.66}{\sqrt{.03888 + .047203}} = \frac{-.66}{\sqrt{.086083}} = -2.25$ . The approximate

degrees of freedom are  $v = \frac{(.086083)^2}{(.03888)^2 + (.047203)^2} = 57.5 > 57$ . The two-tailed *P*-value  $\approx 2(.014) = .028$ ,

which is less than the specified significance level, so we would reject  $H_0$ . The data supports the article's authors' claim.

#### 77.

**a.** The relevant hypotheses are  $H_0: \mu_1 - \mu_2 = 0$  v.  $H_a: \mu_1 - \mu_2 \neq 0$ . Assuming both populations have normal distributions, the two-sample *t* test is appropriate. m = 11,  $\overline{x} = 98.1$ ,  $s_1 = 14.2$ , n = 15,

$$\overline{y} = 129.2$$
,  $s_2 = 39.1$ . The test statistic is  $t = \frac{-31.1}{\sqrt{18.3309 + 101.9207}} = \frac{-31.1}{\sqrt{120.252}} = -2.84$ . The

approximate degrees of freedom 
$$v = \frac{(120.252)^2}{\frac{(18.3309)^2}{10} + \frac{(101.9207)^2}{14}} = 18.64 \searrow 18$$
. From Table A.8, the

two-tailed P-value  $\approx 2(.006) = .012$ . No, obviously the results are different.

**b.** For the hypotheses  $H_0: \mu_1 - \mu_2 = -25$  v.  $H_a: \mu_1 - \mu_2 < -25$ , the test statistic changes to

$$t = \frac{-31.1 - (-25)}{\sqrt{120.252}} = -.556$$
. With df = 18, the *P*-value  $\approx P(T < -.6) = .278$ . Since the *P*-value is greater

than any sensible choice of  $\alpha$ , we fail to reject  $H_0$ . There is insufficient evidence that the true average strength for males exceeds that for females by more than 25 N.

79. To begin, we must find the % difference for each of the 10 meals! For the first meal, the % difference is  $\frac{\text{measured} - \text{stated}}{\text{stated}} = \frac{212 - 180}{180} = .1778$ , or 17.78%. The other nine percentage differences are 45%,

21.58%, 33.04%, 5.5%, 16.49%, 15.2%, 10.42%, 81.25%, and 26.67%.

We wish to test the hypotheses  $H_0$ :  $\mu = 0$  versus  $H_a$ :  $\mu \neq 0$ , where  $\mu$  denotes the true average percent difference for all supermarket convenience meals. A normal probability plot of these 10 values shows some noticeable deviation from linearity, so a t-test is actually of questionable validity here, but we'll proceed just to illustrate the method.

For this sample, 
$$n = 10$$
,  $\bar{x} = 27.29\%$ , and  $s = 22.12\%$ , for a *t* statistic of  $t = \frac{27.29 - 0}{22.12/\sqrt{10}} = 3.90$ .

At df = n - 1 = 9, the *P*-value is  $2P(T_9 \ge 3.90) \approx 2(.002) = .004$ . Since this is smaller than any reasonable significance level, we reject  $H_0$  and conclude that the true average percent difference between meals' stated energy values and their measured values is non-zero.

The normal probability plot below indicates the data for good visibility does not come from a normal 81. distribution. Thus, a t-test is not appropriate for this small a sample size. (The plot for poor visibility isn't as bad.) That is, a pooled t test should not be used here, nor should an "unpooled" two-sample t test be used (since it relies on the same normality assumption).



83. We wish to test  $H_0$ :  $\mu_1 = \mu_2$  versus  $H_a$ :  $\mu_1 \neq \mu_2$ Unpooled:

With  $H_0$ :  $\mu_1 - \mu_2 = 0$  v.  $H_a$ :  $\mu_1 - \mu_2 \neq 0$ , we will reject  $H_0$  if  $p - value < \alpha$ .

$$v = \frac{\left(\frac{.79^2}{14} + \frac{1.52^2}{12}\right)^2}{\left(\frac{.79^2}{14}\right)^2} = 15.95 \downarrow 15, \text{ and the test statistic } t = \frac{8.48 - 9.36}{\sqrt{\frac{.79^2}{14} + \frac{1.52^2}{12}}} = \frac{-.88}{.4869} = -1.81 \text{ leads to a } P$$

value of about  $2P(T_{15} > 1.8) = 2(.046) = .092$ . Pooled:

The degrees of freedom are v = m + n - 2 = 14 + 12 - 2 = 24 and the pooled variance

is 
$$\left(\frac{13}{24}\right)(.79)^2 + \left(\frac{11}{24}\right)(1.52)^2 = 1.3970$$
, so  $s_p = 1.181$ . The test statistic is  
 $t = \frac{-.88}{1.181\sqrt{\frac{1}{14} + \frac{1}{12}}} = \frac{-.88}{.465} \approx -1.89$ . The *P*-value =  $2P(T_{24} > 1.9) = 2(.035) = .070$ .

132

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

#### Chapter 9: Inferences Based on Two Samples

With the pooled method, there are more degrees of freedom, and the *P*-value is smaller than with the unpooled method. That is, if we are willing to assume equal variances (which might or might not be valid here), the pooled test is more capable of detecting a significant difference between the sample means.

85.

**a.** With *n* denoting the second sample size, the first is m = 3n. We then wish  $20 = 2(2.58)\sqrt{\frac{900}{3n} + \frac{400}{n}}$ , which yields n = 47, m = 141.

**b.** We wish to find the n which minimizes  $2z_{a/2}\sqrt{\frac{900}{400-n} + \frac{400}{n}}$ , or equivalently, the *n* which minimizes

 $\frac{900}{400-n} + \frac{400}{n}$ . Taking the derivative with respect to *n* and equating to 0 yields  $900(400-n)^{-2} - 400n^{-2} = 0$ , whence  $9n^2 = 4(400-n)^2$ , or  $5n^2 + 3200n - 640,000 = 0$ . The solution is n = 160, and thus m = 400 - n = 240.

87. We want to test the hypothesis  $H_0: \mu_1 \le 1.5\mu_2 \text{ v}$ .  $H_a: \mu_1 > 1.5\mu_2$  — or, using the hint,  $H_0: \theta \le 0 \text{ v}$ .  $H_a: \theta > 0$ . Our point estimate of  $\theta$  is  $\hat{\theta} = \overline{X}_1 - 1.5\overline{X}_2$ , whose estimated standard error equals  $s(\hat{\theta}) = \sqrt{\frac{s_1^2}{n_1} + (1.5)^2 \frac{s_2^2}{n_2}}$ ,

using the fact that  $V(\hat{\theta}) = \frac{\sigma_1^2}{n_1} + (1.5)^2 \frac{\sigma_2^2}{n_2}$ . Plug in the values provided to get a test statistic t =

 $\frac{22.63 - 1.5(14.15) - 0}{\sqrt{2.8975}} \approx 0.83. \text{ A conservative df estimate here is } v = 50 - 1 = 49. \text{ Since } P(T \ge 0.83) \approx .20$ 

and .20 > .05, we fail to reject  $H_0$  at the 5% significance level. The data does not suggest that the average tip after an introduction is more than 50% greater than the average tip without introduction.

89.  $\Delta_0 = 0, \ \sigma_1 = \sigma_2 = 10, \ d = 1, \ \sigma = \sqrt{\frac{200}{n}} = \frac{14.142}{\sqrt{n}}, \text{ so } \beta = \Phi\left(1.645 - \frac{\sqrt{n}}{14.142}\right), \text{ giving } \beta = .9015, .8264,$ 

.0294, and .0000 for n = 25, 100, 2500, and 10,000 respectively. If the  $\mu_1$ s referred to true average IQs resulting from two different conditions,  $\mu_1 - \mu_2 = 1$  would have little practical significance, yet very large sample sizes would yield statistical significance in this situation.

91.  $H_0: p_1 = p_2$  will be rejected at level  $\alpha$  in favor of  $H_a: p_1 > p_2$  if  $z \ge z_{\alpha}$ . With  $\hat{p}_1 = \frac{250}{2500} = .10$  and  $\hat{p}_2 = \frac{167}{2500} = .0668$ ,  $\hat{p} = .0834$  and  $z = \frac{.0332}{.0079} = 4.2$ , so  $H_0$  is rejected at any reasonable  $\alpha$  level. It appears that a response is more likely for a white name than for a black name.

133

93.

**a.** Let  $\mu_1$  and  $\mu_2$  denote the true average weights for operations 1 and 2, respectively. The relevant hypotheses are  $H_0: \mu_1 - \mu_2 = 0$  v.  $H_a: \mu_1 - \mu_2 \neq 0$ . The value of the test statistic is

$$t = \frac{(1402.24 - 1419.63)}{\sqrt{\frac{(10.97)^2}{30} + \frac{(9.96)^2}{30}}} = \frac{-17.39}{\sqrt{4.011363 + 3.30672}} = \frac{-17.39}{\sqrt{7.318083}} = -6.43.$$
  
At df =  $v = \frac{(7.318083)^2}{\frac{(4.011363)^2}{29} + \frac{(3.30672)^2}{29}} = 57.5 \searrow 57$ ,  $2P(T \le -6.43) \approx 0$ , so we can reject  $H_0$  at level

.05. The data indicates that there is a significant difference between the true mean weights of the packages for the two operations.

- **b.**  $H_0: \mu_1 = 1400$  will be tested against  $H_a: \mu_1 > 1400$  using a one-sample t test with test statistic
  - $t = \frac{\overline{x} 1400}{s_1 / \sqrt{m}}$ . With degrees of freedom = 29, we reject  $H_0$  if  $t \ge t_{.05,29} = 1.699$ . The test statistic value
  - is  $t = \frac{1402.24 1400}{10.97 / \sqrt{30}} = \frac{2.24}{2.00} = 1.1$ . Because 1.1 < 1.699,  $H_0$  is not rejected. True average weight does not appear to exceed 1400.

95.

A large-sample confidence interval for  $\lambda_1 - \lambda_2$  is  $(\hat{\lambda}_1 - \hat{\lambda}_2) \pm z_{\alpha/2} \sqrt{\frac{\hat{\lambda}_1}{m} + \frac{\hat{\lambda}_2}{n}}$ , or  $(\overline{x} - \overline{y}) \pm z_{\alpha/2} \sqrt{\frac{\overline{x}}{m} + \frac{\overline{y}}{n}}$ . With  $\overline{x} = 1.616$  and  $\overline{y} = 2.557$ , the 95% confidence interval for  $\lambda_1 - \lambda_2$  is  $-.94 \pm 1.96(.177) = -.94 \pm .35 = (-1.29, -.59)$ .

134

Exercises of the adjustment of the restaure of the

# **CHAPTER 10**

# Section 10.1

1. The computed value of F is  $f = \frac{\text{MSTr}}{\text{MSE}} = \frac{2673.3}{1094.2} = 2.44$ . Degrees of freedom are I - 1 = 4 and I(J - 1) = (5)(3) = 15. From Table A.9,  $F_{.05,4,15} = 3.06$  and  $F_{.10,4,15} = 2.36$ ; since our computed value of 2.44 is between those values, it can be said that .05 < P-value < .10. Therefore,  $H_0$  is <u>not</u> rejected at the  $\alpha = .05$  level. The data do not provide statistically significant evidence of a difference in the mean tensile strengths of the different types of copper wires.

3. With  $\mu_i$  = true average lumen output for brand *i* bulbs, we wish to test  $H_0: \mu_1 = \mu_2 = \mu_3$  v.  $H_a$ : at least two

- $\mu_i$ 's are different. MSTr =  $\hat{\sigma}_B^2 = \frac{591.2}{2} = 295.60$ , MSE =  $\hat{\sigma}_W^2 = \frac{4773.3}{21} = 227.30$ , so
- $f = \frac{\text{MSTr}}{\text{MSE}} = \frac{295.60}{227.30} = 1.30$ .

For finding the *P*-value, we need degrees of freedom I - 1 = 2 and I(J - 1) = 21. In the 2<sup>nd</sup> row and 21<sup>st</sup> column of Table A.9, we see that  $1.30 < F_{10,2,21} = 2.57$ , so the *P*-value > .10. Since .10 is not < .05, we cannot reject  $H_0$ . There are no statistically significant differences in the average lumen outputs among the three brands of bulbs.

 $\mu_i$  = true mean modulus of elasticity for grade *i* (*i* = 1, 2, 3). We test  $H_0: \mu_1 = \mu_2 = \mu_3$  vs.  $H_a:$  at least two  $\mu_i$ 's are different. Grand mean = 1.5367,

$$MSTr = \frac{10}{2} \Big[ (1.63 - 1.5367)^2 + (1.56 - 1.5367)^2 + (1.42 - 1.5367)^2 \Big] = .1143,$$
  

$$MSE = \frac{1}{3} \Big[ (.27)^2 + (.24)^2 + (.26)^2 \Big] = .0660, \ f = \frac{MSTr}{MSE} = \frac{.1143}{.0660} = 1.73. \ \text{At df} = (2,27), \ 1.73 < 2.51 \Rightarrow \text{the}$$

*P*-value is more than .10. Hence, we fail to reject  $H_0$ . The three grades do not appear to differ significantly.

7. Let  $\mu_i$  denote the true mean electrical resistivity for the *i*th mixture (*i* = 1, ..., 6).

The hypotheses are  $H_0: \mu_1 = ... = \mu_6$  versus  $H_u:$  at least two of the  $\mu_i$ 's are different. There are I = 6 different mixtures and J = 26 measurements for each mixture. That information provides the df values in the table. Working backwards, SSE = I(J - 1)MSE = 2089.350; SSTr = SST - SSE = 3575.065; MSTr = SSTr/(I - 1) = 715.013; and, finally, f = MSTr/MSE = 51.3.

Source	df	SS	MS	f
Treatments	5	3575.065	715.013	51.3
Error	150	2089.350	13.929	
Total	155	5664.415		

The *P*-value is  $P(F_{5,150} \ge 51.3) \approx 0$ , and so  $H_0$  will be rejected at any reasonable significance level. There is strong evidence that true mean electrical resistivity is <u>not</u> the same for all 6 mixtures.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

5.

9.

The summary quantities are  $x_{1.} = 34.3$ ,  $x_{2.} = 39.6$ ,  $x_{3.} = 33.0$ ,  $x_{4.} = 41.9$ ,  $x_{..} = 148.8$ ,  $\Sigma\Sigma x_{ij}^2 = 946.68$ , so  $CF = \frac{(148.8)^2}{24} = 922.56$ , SST = 946.68 - 922.56 = 24.12,  $SSTr = \frac{(34.3)^2 + ... + (41.9)^2}{6} - 922.56 = 8.98$ , SSE = 24.12 - 8.98 = 15.14.

Source	df	SS	MS	F
Treatments	3	8.98	2.99	3.95
Error	20	15.14	.757	
Total	23	24.12		

Since  $3.10 = F_{053,20} < 3.95 < 4.94 = F_{013,20}$ , .01 < P-value < .05, and  $H_0$  is rejected at level .05.

# Section 10.2

11.  $Q_{.05,5,15} = 4.37$ ,  $w = 4.37\sqrt{\frac{272.8}{4}} = 36.09$ . The brands seem to divide into two groups: 1, 3, and 4; and 2 and 5; with no significant differences within each group but all between group differences are significant.  $3 \qquad 1 \qquad 4 \qquad 2 \qquad 5$   $437.5 \qquad 462.0 \qquad 469.3 \qquad 512.8 \qquad 532.1$ 

13. Brand 1 does not differ significantly from 3 or 4, 2 does not differ significantly from 4 or 5, 3 does not differ significantly from 1, 4 does not differ significantly from 1 or 2, 5 does not differ significantly from 2, but all other differences (e.g., 1 with 2 and 5, 2 with 3, etc.) do appear to be significant.

427.5	462.0	469.3	502.8	532.1
3	1	4	2	5

15. In Exercise 10.7, I = 6 and J = 26, so the critical value is  $Q_{.05,6,150} \approx Q_{.05,6,120} = 4.10$ , and MSE = 13.929. So,  $w \approx 4.10 \sqrt{\frac{13.929}{26}} = 3.00$ . So, sample means less than 3.00 apart will belong to the same underscored set.

Three distinct groups emerge: the first mixture (in the above order), then mixtures 2-4, and finally mixtures 5-6.

111 - (1.4.1. Skin + (No.1.) (0.1.4 (1.4.2.1-53.0) - = 1234

17.

 $\theta = \Sigma c_i \mu_i$  where  $c_1 = c_2 = .5$  and  $c_3 = -1$ , so  $\hat{\theta} = .5\overline{x}_{1.} + .5\overline{x}_{2.} - \overline{x}_{3.} = -.527$  and  $\Sigma c_i^2 = 1.50$ . With  $t_{.025,27} = 2.052$  and MSE = .0660, the desired CI is (from (10.5))

$$-.527 \pm (2.052) \sqrt{\frac{(.0660)(1.50)}{10}} = -.527 \pm .204 = (-.731, -.323).$$

## Chapter 10: The Analysis of Variance

MSTr = 140, error df = 12, so 
$$f = \frac{140}{\text{SSE}/12} = \frac{1680}{\text{SSE}}$$
 and  $F_{.05,2,12} = 3.89$ .

$$v = Q_{.05,3,12} \sqrt{\frac{\text{MSE}}{J}} = 3.77 \sqrt{\frac{\text{SSE}}{60}} = .4867 \sqrt{\text{SSE}}$$
. Thus we wish  $\frac{1680}{SSE} > 3.89$  (significant f) and

.4867 $\sqrt{\text{SSE}} > 10 \ (= 20 - 10)$ , the difference between the extreme  $\overline{x_i}$  's, so no significant differences are identified). These become 431.88 > SSE and SSE > 422.16, so SSE = 425 will work.

21.

**a.** The hypotheses are  $H_0$ :  $\mu_1 = \dots = \mu_6 v$ .  $H_a$ : at least two of the  $\mu_i$ 's are different. Grand mean = 222.167, MSTr = 38,015.1333, MSE = 1,681.8333, and f = 22.6. At df = (5, 78)  $\approx$  (5, 60), 22.6  $\geq$  4.76  $\Rightarrow$  *P*-value < .001. Hence, we reject  $H_0$ . The data indicate there is

a dependence on injection regimen.

**b.** Assume  $t_{.005,78} \approx 2.645$ .

) Confidence interval for 
$$\mu_1 - \frac{1}{5}(\mu_2 + \mu_3 + \mu_4 + \mu_5 + \mu_6)$$
:  $\Sigma c_i \overline{x}_i \pm t_{\alpha/2, I(J-1)} \sqrt{\frac{\text{MSE}(\Sigma c_i^2)}{J}}$ 

$$= -67.4 \pm (2.645) \sqrt{\frac{1,681.8333(1.2)}{14}} = (-99.16, -35.64).$$

ii) Confidence interval for  $\frac{1}{4}(\mu_2 + \mu_3 + \mu_4 + \mu_5) - \mu_6$ :

$$= 61.75 \pm (2.645) \sqrt{\frac{1,681.8333(1.25)}{14}} = (29.34,94.16)$$

# Section 10.3

23.  $J_{1} = 5, J_{2} = 4, J_{3} = 4, J_{4} = 5, \ \overline{x}_{1} = 58.28, \ \overline{x}_{2} = 55.40, \ \overline{x}_{3} = 50.85, \ \overline{x}_{4} = 45.50, \ \text{MSE} = 8.89.$ With  $W_{ij} = Q_{.05,4,14} \cdot \sqrt{\frac{\text{MSE}}{2} \left(\frac{1}{J_{i}} + \frac{1}{J_{j}}\right)} = 4.11 \sqrt{\frac{8.89}{2} \left(\frac{1}{J_{i}} + \frac{1}{J_{j}}\right)},$   $\overline{x}_{1} - \overline{x}_{2} \pm W_{12} = (2.88) \pm (5.81); \ \overline{x}_{1} - \overline{x}_{3} \pm W_{13} = (7.43) \pm (5.81) *; \ \overline{x}_{1} - \overline{x}_{4} \pm W_{14} = (12.78) \pm (5.48) *;$   $\overline{x}_{2} - \overline{x}_{3} \pm W_{23} = (4.55) \pm (6.13); \ \overline{x}_{2} - \overline{x}_{4} \pm W_{24} = (9.90) \pm (5.81) *; \ \overline{x}_{3} - \overline{x}_{4} \pm W_{34} = (5.35) \pm (5.81).$ 

A \* indicates an interval that doesn't include zero, corresponding to  $\mu$ 's that are judged significantly different. This underscoring pattern does not have a very straightforward interpretation.

25.

- a. The distributions of the polyunsaturated fat percentages for each of the four regimens must be normal with equal variances.
- **b.** We have all the  $\overline{x}_i$  s, and we need the grand mean:

$$\overline{x}_{11} = \frac{8(43.0) + 13(42.4) + 17(43.1) + 14(43.5)}{52} = \frac{2236.9}{52} = 43.017;$$

$$SSTr = \sum J_i (\overline{x}_i - \overline{x}_i)^2 = 8(43.0 - 43.017)^2 + 13(42.4 - 43.017)^2$$
  
+17(43.1 - 43.017)<sup>2</sup> + 13(43.5 - 43.017)<sup>2</sup> = 8.334 and MSTr =  $\frac{8.334}{3}$  = 2.778  
$$SSE = \sum (J_i - 1)s^2 = 7(1.5)^2 + 12(1.3)^2 + 16(1.2)^2 + 13(1.2)^2 = 77.79 \text{ and } MSE = \frac{77.79}{48} = 1.621. \text{ Then}$$
  
$$f = \frac{MSTr}{MSE} = \frac{2.778}{1.621} = 1.714$$

Since  $1.714 < F_{.10,3,50} = 2.20$ , we can say that the *P*-value is > .10. We do not reject the null hypothesis at significance level .10 (or any smaller), so we conclude that the data suggests no difference in the percentages for the different regimens.

27.

**a.** Let  $\mu_i$  = true average folacin content for specimens of brand *i*. The hypotheses to be tested are  $H_0: \mu_1 = \mu_2 = \mu_3 = \mu_4$  vs.  $H_a:$  at least two of the  $\mu_i$ 's are different.  $\Sigma \Sigma x_{ii}^2 = 1246.88$  and

$\frac{x_{}^2}{x_{}^2} = ($	$(168.4)^2 = 1181.61$ so SST = 65.27:	$\Sigma x_{i}^2$	$(57.9)^2$	$(37.5)^2$	$(38.1)^2$	$(34.9)^2$	-1205 10 50
n	24 24	J	7	5	6	6	- 1203.10, 30
SSTr	= 1205.10 - 1181.61 = 23.49.						

Source	df	SS	MS	F
Treatments	3	23.49	7.83	3.75
Error	20	41.78	2.09	
Total	23	65.27		

With numerator df = 3 and denominator df = 20,  $F_{.05,3,20} = 3.10 < 3.75 < F_{.01,3,20} = 4.94$ , so the *P*-value is between .01 and .05. We reject  $H_0$  at the .05 level: at least one of the pairs of brands of green tea has different average folacin content.

**b.** With  $\overline{x}_{i} = 8.27, 7.50, 6.35$ , and 5.82 for i = 1, 2, 3, 4, we calculate the residuals  $x_{ij} - \overline{x}_{i}$  for all

observations. A normal probability plot appears below and indicates that the distribution of residuals could be normal, so the normality assumption is plausible. The sample standard deviations are 1.463, 1.681, 1.060, and 1.551, so the equal variance assumption is plausible (since the largest sd is less than twice the smallest sd).



Normal Probability Plot for ANOVA Residuals
### Chapter 10: The Analysis of Variance

Pair	Interval	Pair	Interval	The hypotheses he H = a +
1,2	$.77 \pm 2.37$	2,3	1.15±2.45a	test statute in the state of the states.
1,3	$1.92 \pm 2.25$	2,4	$1.68\pm2.45$	
1,4	2.45 ± 2.25 *	3,4	.53±2.34	

Only Brands 1 and 4 are significantly different from each other.

$$29. \qquad E\left(\text{SSTr}\right) = E\left(\sum_{i} J_{i} \overline{X}_{i}^{2} - n \overline{X}_{-}^{2}\right) = \sum J_{i} E\left(\overline{X}_{i}^{2}\right) - n E\left(\overline{X}_{-}^{2}\right)$$
$$= \sum J_{i} \left[V\left(\overline{X}_{i}\right) + \left(E\left(\overline{X}_{i}\right)\right)^{2}\right] - n \left[V\left(\overline{X}_{-}\right) + \left(E\left(\overline{X}_{-}\right)\right)^{2}\right] = \sum J_{i} \left[\frac{\sigma^{2}}{J_{i}} + \mu_{i}^{2}\right] - n \left[\frac{\sigma^{2}}{n} + \left(\frac{\sum J_{i} \mu_{i}}{n}\right)^{2}\right]$$
$$= I\sigma^{2} + \sum J_{i} \left(\mu + \alpha_{i}\right)^{2} - \sigma^{2} - \frac{1}{n} \left[\sum J_{i} \left(\mu + \alpha_{i}\right)\right]^{2} = (I-1)\sigma^{2} + \sum J_{i}\mu^{2} + 2\mu \sum J_{i}\alpha_{i}^{2} - \frac{1}{n} \left[n\mu + 0\right]^{2}$$
$$= (I-1)\sigma^{2} + \mu^{2}n + 2\mu 0 + \sum J_{i}\alpha_{i}^{2} - n\mu^{2} = (I-1)\sigma^{2} + \sum J_{i}\alpha_{i}^{2}, \text{ from which } E(\text{MSTr}) \text{ is obtained through division by } (I-1).$$

31. With 
$$\sigma = 1$$
 (any other  $\sigma$  would yield the same  $\phi$ ),  $\alpha_1 = -1$ ,  $\alpha_2 = \alpha_3 = 0$ ,  $\alpha_4 = 1$ ,  
 $\phi^2 = \frac{1(5(-1)^2 + 4(0)^2 + 4(0)^2 + 5(1)^2)}{4} = 2.5$ ,  $\phi = 1.58$ ,  $v_1 = 3$ ,  $v_2 = 14$ , and power  $\approx .65$ .

33.  $g(x) = x\left(1-\frac{x}{n}\right) = nu(1-u)$  where  $u = \frac{x}{n}$ , so  $h(x) = \int \left[u(1-u)\right]^{-1/2} du$ . From a table of integrals, this gives  $h(x) = \arcsin\left(\sqrt{u}\right) = \arcsin\left(\sqrt{\frac{x}{n}}\right)$  as the appropriate transformation.

bis is a modern effects intuition.  $R_1 \ge e_1^2 = 0$  intris that variation in Liberatories down? contribute to mixton in percentage. SCT = 16,078,5897 – 36,077,2024 = 1,7673, SSTe = 1,0559, and SSE = ,7116, 14f – (2, 8), 2.92 < 3.96 < 4.07 mb, 05  $\times$  P value < 10, so  $R_0$  cannot be rejected at level .05. Variation bound that does not appear to be present.

139

### Supplementary Exercises

35.

- a. The hypotheses are H<sub>0</sub>: μ<sub>1</sub> = μ<sub>2</sub> = μ<sub>3</sub> = μ<sub>4</sub> v. H<sub>a</sub>: at least two of the μ<sub>i</sub>'s are different. The calculated test statistic is f = 3.68. Since F<sub>.05,3,20</sub> = 3.10 < 3.68 < F<sub>.01,3,20</sub> = 4.94, the P-value is between .01 and .05. Thus, we fail to reject H<sub>0</sub> at α = .01. At the 1% level, the means do not appear to differ significantly.
- **b.** We reject  $H_0$  when the *P*-value  $\leq \alpha$ . Since .029 is not < .01, we still fail to reject  $H_0$ .
- 37. Let  $\mu_i$  = true average amount of motor vibration for each of five bearing brands. Then the hypotheses are  $H_0: \mu_1 = ... = \mu_5$  vs.  $H_a$ : at least two of the  $\mu_i$ 's are different. The ANOVA table follows:

Source	df	SS	MS	F
Treatments	4	30.855	7.714	8.44
Error	25	22.838	0.914	
Total	29	53.694		

 $8.44 > F_{.001,4,25} = 6.49$ , so *P*-value < .001 < .05, so we reject  $H_0$ . At least two of the means differ from one another. The Tukey multiple comparisons are appropriate.  $Q_{.05,5,25} = 4.15$  from Minitab output; or, using

Table A.10, we can approximate with  $Q_{.05,5,24} = 4.17$ .  $W_{ij} = 4.15\sqrt{.914/6} = 1.620$ .

Pair	$\overline{x}_{i} - \overline{x}_{j}$	Pair	$\overline{X}_{i\cdot} - \overline{X}_{j\cdot}$
1,2	-2.267*	2,4	1.217
1,3	0.016	2,5	2.867*
1,4	-1.050	3,4	-1.066
1,5	0.600	3,5	0.584
2,3	2.283*	4,5	1.650*

\*Indicates significant pairs.

- 39.  $\hat{\theta} = 2.58 \frac{2.63 + 2.13 + 2.41 + 2.49}{4} = .165, t_{025,25} = 2.060$ , MSE = .108, and  $\Sigma c_i^2 = (1)^2 + (-.25)^2 + (-.25)^2 + (-.25)^2 = 1.25$ , so a 95% confidence interval for  $\theta$  is  $.165 \pm 2.060 \sqrt{\frac{(.108)(1.25)}{6}} = .165 \pm .309 = (-.144,.474)$ . This interval does include zero, so 0 is a plausible value for  $\theta$ .
- 41. This is a random effects situation.  $H_0: \sigma_A^2 = 0$  states that variation in laboratories doesn't contribute to variation in percentage. SST = 86,078.9897 86,077.2224 = 1.7673, SSTr = 1.0559, and SSE = .7114. At df = (3, 8), 2.92 < 3.96 < 4.07  $\Rightarrow$  .05 < *P*-value < .10, so  $H_0$  cannot be rejected at level .05. Variation in laboratories does not appear to be present.

www.konkur.in

45.

## Chapter 10: The Analysis of Variance

43. 
$$\sqrt{(I-1)(\text{MSE})(F_{.05,I-1,n-I})} = \sqrt{(2)(2.39)(3.63)} = 4.166. \text{ For } \mu_1 - \mu_2, c_1 = 1, c_2 = -1, \text{ and } c_3 = 0, \text{ so}$$
$$\sqrt{\sum \frac{c_i^2}{J_i}} = \sqrt{\frac{1}{8} + \frac{1}{5}} = .570. \text{ Similarly, for } \mu_1 - \mu_3, \sqrt{\sum \frac{c_i^2}{J_i}} = \sqrt{\frac{1}{8} + \frac{1}{6}} = .540; \text{ for } \mu_2 - \mu_3,$$
$$\sqrt{\sum \frac{c_i^2}{J_i}} = \sqrt{\frac{1}{5} + \frac{1}{6}} = .606, \text{ and for } .5\mu_2 + .5\mu_2 - \mu_3, \sqrt{\sum \frac{c_i^2}{J_i}} = \sqrt{\frac{.5^2}{8} + \frac{.5^2}{5} + \frac{(-1)^2}{6}} = .498.$$

Contrast Estimate		Interval
$\mu_1 - \mu_2$	25.59 - 26.92 = -1.33	$(-1.33)\pm(.570)(4.166)=(-3.70,1.04)$
$\mu_1 - \mu_3$	25.59 - 28.17 = -2.58	$(-2.58)\pm(.540)(4.166)=(-4.83,33)$
$\mu_2 - \mu_3$	26.92 - 28.17 = -1.25	$(-1.25) \pm (.606)(4.166) = (-3.77, 1.27)$
$.5\mu_2 + .5\mu_2 - \mu_3$	-1.92	$(-1.92)\pm(.498)(4.166)=(-3.99,0.15)$

The contrast between  $\mu_1$  and  $\mu_3$ , since the calculated interval is the only one that does not contain 0.

 $Y_{ij} - \overline{Y}_{a} = c \left( X_{ij} - \overline{X}_{a} \right)$  and  $\overline{Y}_{i} - \overline{Y}_{a} = c \left( \overline{X}_{i} - \overline{X}_{a} \right)$ , so each sum of squares involving Y will be the corresponding sum of squares involving X multiplied by  $c^{2}$ . Since F is a ratio of two sums of squares,  $c^{2}$  appears in both the numerator and denominator. So  $c^{2}$  cancels, and F computed from  $Y_{ij}$ 's = F computed from  $X_{ij}$ 's.

## **CHAPTER 11**

## Section 11.1

1.

**a.** The test statistic is  $f_A = \frac{\text{MSA}}{\text{MSE}} = \frac{\text{SSA}/(I-1)}{\text{SSE}/(I-1)(J-1)} = \frac{442.0/(4-1)}{123.4/(4-1)(3-1)} = 7.16$ . Compare this to the

*F* distribution with df = (4 - 1, (4 - 1)(3 - 1)) = (3, 6):  $4.76 < 7.16 < 9.78 \Rightarrow .01 < P$ -value < .05. In particular, we reject  $H_{0A}$  at the .05 level and conclude that at least one of the factor A means is different (equivalently, at least one of the  $a_i$ 's is not zero).

**b.** Similarly,  $f_B = \frac{\text{SSB}/(J-1)}{\text{SSE}/(I-1)(J-1)} = \frac{428.6/(3-1)}{123.4/(4-1)(3-1)} = 10.42$ . At df = (2, 6), 5.14 < 10.42 < 10.92

 $\Rightarrow .01 < P$ -value < .05. In particular, we reject  $H_{0B}$  at the .05 level and conclude that at least one of the factor B means is different (equivalently, at least one of the  $\beta_j$ 's is not zero).

3.

a.	The entries of	this AN	<b>IOVA</b> table	were p	produced	with software.	
----	----------------	---------	-------------------	--------	----------	----------------	--

Source	df	SS	MS	F
Medium	1	0.053220	0.0532195	18.77
Current	3	0.179441	0.0598135	21.10
Error	3	0.008505	0.0028350	
Total	7	0.241165		
				10 A 10 AN 100

To test  $H_{0,4}$ :  $\alpha_1 = \alpha_2 = 0$  (no liquid medium effect), the test statistic is  $f_A = 18.77$ ; at df = (1, 3), the *P*-value is .023 from software (or between .01 and .05 from Table A.9). Hence, we reject  $H_{0,4}$  and conclude that medium (oil or water) affects mean material removal rate.

To test  $H_{0B}$ :  $\beta_1 = \beta_2 = \beta_3 = \beta_4 = 0$  (no current effect), the test statistic is  $f_B = 21.10$ ; at df = (3, 3), the *P*-value is .016 from software (or between .01 and .05 from Table A.9). Hence, we reject  $H_{0B}$  and conclude that working current affects mean material removal rate as well.

**b.** Using a .05 significance level, with J = 4 and error df = 3 we require  $Q_{.05,4,3} = 6.825$ . Then, the metric for significant differences is  $w = 6.825\sqrt{0.0028530/2} = 0.257$ . The means happen to increase with current; sample means and the underscore scheme appear below.

Current:	10	15	20	25
$\overline{x}_{j}$ :	0.201	0.324	0.462	0.602

www.konkur.

5.

7.

9.

## Chapter 11: Multifactor Analysis of Variance

Source	df	SS	MS	F	
Angle	3	58.16	19.3867	2.5565	
Connector	4	246.97	61.7425	8.1419	
Error	12	91.00	7.5833		
Total	19	396.13			

We're interested in  $H_0: \alpha_1 = \alpha_2 = \alpha_3 = \alpha_4 = 0$  versus  $H_a$ : at least one  $\alpha_i \neq 0$ .  $f_A = 2.5565 < F_{.01,3,12} = 5.95 \Rightarrow$ *P*-value > .01, so we fail to reject  $H_0$ . The data fails to indicate any effect due to the angle of pull, at the .01 significance level.

a. The entries of this ANOVA table were produced with software.

Source	df	SS	MS	F
Brand	2	22.8889	11.4444	8.96
Operator	2	27.5556	13.7778	10.78
Error	4	5.1111	1.2778	
Total	8	55.5556		

The calculated test statistic for the *F*-test on brand is  $f_A = 8.96$ . At df = (2, 4), the *P*-value is .033 from software (or between .01 and .05 from Table A.9). Hence, we reject  $H_0$  at the .05 level and conclude that lathe brand has a statistically significant effect on the percent of acceptable product.

**b.** The block-effect test statistic is f = 10.78, which is quite large (a *P*-value of .024 at df = (2, 4)). So, yes, including this operator blocking variable was a good idea, because there is significant variation due to different operators. If we had not controlled for such variation, it might have affected the analysis and conclusions.

## The entries of this ANOVA table were produced with software.

Source	df	SS	MS	F
Treatment	3	81.1944	27.0648	22.36
Block	8	66.5000	8.3125	6.87
Error	24	29.0556	1.2106	
Total	35	176.7500		

At df = (3, 24),  $f = 22.36 > 7.55 \Rightarrow P$ -value < .001. Therefore, we strongly reject  $H_{0A}$  and conclude that there is an effect due to treatments. We follow up with Tukey's procedure:

 $Q_{.05,4,24} = 3.90; w = 3.90\sqrt{1.2106/9} = 1.43$ 

1	4	3	2
8.56	9.22	10.78	12.44

**11.** The residual, percentile pairs are (-0.1225, -1.73), (-0.0992, -1.15), (-0.0825, -0.81), (-0.0758, -0.55), (-0.0750, -0.32), (0.0117, -0.10), (0.0283, 0.10), (0.0350, 0.32), (0.0642, 0.55), (0.0708, 0.81), (0.0875, 1.15), (0.1575, 1.73).



The pattern is sufficiently linear, so normality is plausible.

13.

- **a.** With  $Y_{ij} = X_{ij} + d$ ,  $\overline{Y}_{i.} = \overline{X}_{i.} + d$  and  $\overline{Y}_{j.} = \overline{X}_{.j} + d$  and  $\overline{Y}_{..} = \overline{X}_{..} + d$ , so all quantities inside the parentheses in (11.5) remain unchanged when the Y quantities are substituted for the corresponding X's (e.g.,  $\overline{Y}_{i.} \overline{Y}_{..} = \overline{X}_{i.} \overline{X}_{..}$ , etc.).
- **b.** With  $Y_{ij} = cX_{ij}$ , each sum of squares for Y is the corresponding SS for X multiplied by  $c^2$ . However, when F ratios are formed the  $c^2$  factors cancel, so all F ratios computed from Y are identical to those computed from X. If  $Y_{ij} = cX_{ij} + d$ , the conclusions reached from using the Y's will be identical to those reached using the X's.
- 15.
- a.  $\Sigma \alpha_i^2 = 24$ , so  $\phi^2 = \left(\frac{3}{4}\right) \left(\frac{24}{16}\right) = 1.125$ ,  $\phi = 1.06$ ,  $v_1 = 3$ ,  $v_2 = 6$ , and from Figure 10.5, power  $\approx .2$ . For the second alternative,  $\phi = 1.59$ , and power  $\approx .43$ .
- **b.**  $\phi^2 = \left(\frac{I}{J}\right) \sum \frac{\beta_j^2}{\sigma^2} = \left(\frac{4}{5}\right) \left(\frac{20}{16}\right) = 1.00$ , so  $\phi = 1.00$ ,  $v_1 = 4$ ,  $v_2 = 12$ , and power  $\approx .3$ .

www.konkur.i

## Chapter 11: Multifactor Analysis of Variance

## Section 11.2

a.

17.

Source	df	SS	MS	F	P-value
Sand	2	705	352.5	3.76	.065
Fiber	2	1,278	639.0	6.82	.016
Sand × Fiber	4	279	69.75	0.74	.585
Error	9	843	93.67		
Total	17	3,105			

*P*-values were obtained from software; approximations can also be acquired using Table A.9. There appears to be an effect due to carbon fiber addition, but not due to any other effect (interaction effect or sand addition main effect).

Source	df	SS	MS	F	P-value
Sand	2	106.78	53.39	6.54	.018
Fiber	2	87.11	43.56	5.33	.030
Sand × Fiber	4	8.89	2.22	0.27	.889
Error	9	73.50	8.17		
Total	17	276.28			

There appears to be an effect due to both sand and carbon fiber addition to casting hardness, but no interaction effect.

-		
Ŀ		
•		

b.

Sand%	0	15	30	0	15	30	0	15	30
Fiber%	0	0	0	0.25	0.25	0.25	0.5	0.5	0.5
ī	62	68	69.5	69	71.5	73	68	71.5	74

The plot below indicates some effect due to sand and fiber addition with no significant interaction. This agrees with the statistical analysis in part **b**.



145

٠		•		
	£			
	- 2	,	-	
•	-		٠	

Source	df	SS	MS	F
Farm Type	2	35.75	17.875	0.94
Fractor Maint. Method	5	861.20	172.240	9.07
Type × Method	10	603.51	60.351	3.18
Error	18	341.82	18.990	
Total	35	1842.28		

For the interaction effect,  $f_{AB} = 3.18$  at df = (10, 18) gives *P*-value = .016 from software. Hence, we do not reject  $H_{0AB}$  at the .01 level (although just barely). This allows us to proceed to the main effects.

For the factor A main effect,  $f_A = 0.94$  at df = (2, 18) gives *P*-value = .41 from software. Hence, we clearly fail to reject  $H_{0A}$  at the .01 level — there is not statistically significant effect due to type of farm.

Finally,  $f_B = 9.07$  at df = (5, 18) gives *P*-value < .0002 from software. Hence, we strongly reject  $H_{0B}$  at the .01 level — there is a statistically significant effect due to tractor maintenance method.

21.

From the provided SS, SSAB = 64,954.70 - [22,941.80 + 22,765.53 + 15,253.50] = 3993.87. This allows us to complete the ANOVA table below.

df	SS	MS	F
2	22,941.80	11,470.90	22.98
4	22,765.53	5691.38	11.40
8	3993.87	499.23	.49
15	15,253.50	1016.90	
29	64,954.70		
	df 2 4 8 15 29	df         SS           2         22,941.80           4         22,765.53           8         3993.87           15         15,253.50           29         64,954.70	df         SS         MS           2         22,941.80         11,470.90           4         22,765.53         5691.38           8         3993.87         499.23           15         15,253.50         1016.90           29         64,954.70         64,954.70

 $f_{AB} = .49$  is clearly not significant. Since  $22.98 \ge F_{.05,2,8} = 4.46$ , the *P*-value for factor A is < .05 and  $H_{0A}$  is rejected. Since  $11.40 \ge F_{.05,4,8} = 3.84$ , the *P*-value for factor B is < .05 and  $H_{0B}$  is also rejected. We conclude that the different cement factors affect flexural strength differently and that batch variability contributes to variation in flexural strength.

23.

of concrete cylinders.

Summary quantities include  $x_{1_{...}} = 9410$ ,  $x_{2_{...}} = 8835$ ,  $x_{3_{...}} = 9234$ ,  $x_{1_{...}} = 5432$ ,  $x_{2_{...}} = 5684$ ,  $x_{3_{...}} = 5619$ ,  $x_{4_{...}} = 5567$ ,  $x_{5_{...}} = 5177$ ,  $x_{...} = 27,479$ , CF = 16,779,898.69,  $\Sigma x_{i_{...}}^2 = 251,872,081$ ,  $\Sigma x_{j_{...}}^2 = 151,180,459$ , resulting in the accompanying ANOVA table.

in the accompany	ng Anovr	i table.		
Source	df	SS	MS	F
А	2	11,573.38	5786.69	$\frac{MSA}{MSAB} = 26.70$
В	4	17,930.09	4482.52	$\frac{MSB}{MSAB} = 20.68$
AB	8	1734.17	216.77	$\frac{MSAB}{MSE} = 1.38$
Error	30	4716.67	157.22	
Total	44	35,954.31		

Since  $1.38 < F_{.01,8,30} = 3.17$ , the interaction *P*-value is > .01 and  $H_{0G}$  cannot be rejected. We continue:  $26.70 \ge F_{.01,2,8} = 8.65 \implies \text{factor A } P\text{-value} < .01 \text{ and } 20.68 \ge F_{.01,4,8} = 7.01 \implies \text{factor B } P\text{-value} < .01$ , so both  $H_{0A}$  and  $H_{0B}$  are rejected. Both capping material and the different batches affect compressive strength

www.konkur.i

# Chapter 11: Multifactor Analysis of Variance

25. With 
$$\theta = \alpha_i - \alpha'_i$$
,  $\hat{\theta} = \overline{X}_{i...} - \overline{X}_{i'...} = \frac{1}{JK} \sum_{j \ k} [X_{ijk} - X_{i'jk}]$ , and since  $i \neq i'$ ,  $X_{ijk}$  and  $X_{i'jk}$  are independent  
for every  $j$ ,  $k$ . Thus,  $V(\hat{\theta}) = V(\overline{X}_{i..}) + V(\overline{X}_{i'..}) = \frac{\sigma^2}{JK} + \frac{\sigma^2}{JK} = \frac{2\sigma^2}{JK}$  (because  $V(\overline{X}_{i...}) = V(\overline{\varepsilon}_{i...})$  and  
 $V(\varepsilon_{ijk}) = \sigma^2$ ) so  $\hat{\sigma}_{\hat{\theta}} = \sqrt{\frac{2MSE}{JK}}$ . The appropriate number of df is  $IJ(K-1)$ , so the CI is  
 $(\overline{x}_{i...} - \overline{x}_{i'...}) \pm t_{\alpha/2,IJ(K-1)} \sqrt{\frac{2MSE}{JK}}$ . For the data of exercise 19,  $\overline{x}_{2...} = 8.192, \overline{x}_{3...} = 8.395, MSE = .0170,$   
 $t_{.025,9} = 2.262, J = 3, K = 2$ , so the 95% C.I. for  $\alpha_2 - \alpha_3$  is  $(8.182 - 8.395) \pm 2.262 \sqrt{\frac{.0340}{6}} = -0.203 \pm 0.170$   
 $= (-0.373, -0.033).$ 

### Section 11.3

### 27.

Source	df	SS	MS	F	$F_{.05,\mathrm{num}}$ df, den df
Δ	2	14,144,44	7072.22	61.06	3.35
B	2	5.511.27	2755.64	23.79	3.35
C	2	244 696 39	122.348.20	1056.24	3.35
AB	4	1.069.62	267.41	2.31	2.73
AC	4	62.67	15.67	.14	2.73
BC	4	331.67	82.92	.72	2.73
ABC	8	1,080.77	135.10	1.17	2.31
Error	27	3,127.50	115.83		

270,024.33

- b. The computed F-statistics for all four interaction terms (2.31, .14, .72, 1.17) are less than the tabled values for statistical significance at the level .05 (2.73 for AB/AC/BC, 2.31 for ABC). Hence, all four P-values exceed .05. This indicates that none of the interactions are statistically significant.
- **c.** The computed *F*-statistics for all three main effects (61.06, 23.79, 1056.24) exceed the tabled value for significance at level .05 ( $3.35 = F_{.05,2,27}$ ). Hence, all three *P*-values are less than .05 (in fact, all three *P*-values are less than .001), which indicates that all three main effects are statistically significant.

**d.** Since  $Q_{.05,3,27}$  is not tabled, use  $Q_{.05,3,24} = 3.53$ ,  $w = 3.53\sqrt{\frac{115.83}{(3)(3)(2)}} = 8.95$ . All three levels differ

significantly from each other.

53

Total

147

Source	df	SS	MS	F	P-value
A	2	1043.27	521.64	110.69	<.001
В	1	112148.10	112148.10	23798.01	<.001
С	2	3020.97	1510.49	320.53	<.001
AB	2	373.52	186.76	39.63	<.001
AC	4	392.71	98.18	20.83	<.001
BC	2	145.95	72.98	15.49	<.001
ABC	4	54.13	13.53	2.87	.029
Error	72	339.30	4.71		
Total	89	117517.95			

*P*-values were obtained using software. At the .01 significance level, all main and two-way interaction effects are statistically significant (in fact, extremely so), but the three-way interaction is not statistically significant (.029 > .01).

**b.** The means provided allow us to construct an AB interaction plot and an AC interaction plot. Based on the first plot, it's actually surprising that the AB interaction effect is significant: the "bends" of the two paths (B = 1, B = 2) are different but not that different. The AC interaction effect is more clear: the effect of C = 1 on mean response decreases with A (= 1, 2, 3), while the pattern for C = 2 and C = 3 is very different (a sharp up-down-up trend).





a. The following ANOVA table was created with software.

Source	df	SS	MS	F	P-value
A	2	124.60	62.30	4.85	.042
В	2	20.61	10.30	0.80	.481
С	2	356.95	178.47	13.89	.002
AB	4	57.49	14.37	1.12	.412
AC	4	61.39	15.35	1.19	.383
BC	4	11.06	2.76	0.22	.923
Error	8	102.78	12.85		
Total	26	734.87			

**b.** The *P*-values for the *AB*, *AC*, and *BC* interaction effects are provided in the table. All of them are much greater than .1, so none of the interaction terms are statistically significant.

29.

a.

www.konkur.in

## Chapter 11: Multifactor Analysis of Variance

- c. According to the *P*-values, the factor *A* and *C* main effects are statistically significant at the .05 level. The factor *B* main effect is not statistically significant.
- **d.** The paste thickness (factor *C*) means are 38.356, 35.183, and 29.560 for thickness .2, .3, and .4, respectively. Applying Tukey's method,  $Q_{.05,3,8} = 4.04 \Rightarrow w = 4.04\sqrt{12.85/9} = 4.83$ .

Thickness:	.4	.3	.2
Mean:	29,560	35.183	38.356
Ivicali.	27.000		and the second se

33. The various sums of squares yield the accompanying ANOVA table.

Source	df	SS	MS	F
A	6	67.32	11.02	
В	6	51.06	8.51	
Ċ	6	5.43	.91	.61
Error	30	44.26	1.48	
Total	48	168.07		

We're interested in factor C. At df = (6, 30),  $.61 < F_{.05,6,30} = 2.42 \Rightarrow P$ -value > .05. Thus, we fail to reject  $H_{0C}$  and conclude that heat treatment had no effect on aging.

	1	2	3	4	5	
$x_i$	40.68	30.04	44.02	32.14	33.21	$\Sigma x_{i}^2 = 6630.91$
<i>x</i> .,	29.19	31.61	37.31	40.16	41.82	$\Sigma x_{.j.}^2 = 6605.02$
<i>xk</i>	36.59	36.67	36.03	34.50	36.30	$\Sigma x_{k}^2 = 6489.92$

x = 180.09, CF = 1297.30,  $\Sigma\Sigma x_{ii(k)}^2 = 1358.60$ 

N.S.	Source	df	SS	MS	F
-	A	4	28.89	7.22	10.71
	B D	4	23.71	5.93	8.79
	C	emod 4	0.63	0.16	0.23
	Error	12	8.09	0.67	
	Total	24	61.30		

 $F_{.05,4,12} = 3.26$ , so the *P*-values for factor A and B effects are < .05 (10.71 > 3.26, 8.79 > 3.26), but the *P*-value for the factor C effect is > .05 (0.23 < 3.26). Both factor A (plant) and B(leaf size) appear to affect moisture content, but factor C (time of weighing) does not.

Source	df	MS	F	F.01,num df, den df*
Α	2	2207.329	2259.29	5.39
В	1	47.255	48.37	7.56
С	2	491.783	503.36	5.39
D	1	.044	.05	7.56
AB	2	15.303	15.66	5.39
AC	4	275.446	281.93	4.02
AD	2	.470	.48	5.39
BC	2	2.141	2.19	5.39
BD	1	.273	.28	7.56
CD	2	.247	.25	5.39
ABC	4	3.714	3.80	4.02
ABD	2	4.072	4.17	5.39
ACD	4	.767	.79	4.02
BCD	2	.280	.29	5.39
ABCD	4	.347	.355	4.02
Error	36	.977	and marked	
Total	71		and the second second	

37.	SST = (71)(93.621) = 6,647.091. Computing all other sums of squares and adding them up = 6,645.702.
	Thus $SSABCD = 6,647.091 - 6,645.702 = 1.389$ and $MSABCD = 1.389/4 = .347$ .

\*Because denominator df for 36 is not tabled, use df = 30.

To be significant at the .01 level (*P*-value < .01), the calculated *F* statistic must be greater than the .01 critical value in the far right column. At level .01 the statistically significant main effects are A, B, C. The interaction AB and AC are also statistically significant. No other interactions are statistically significant.

## Section 11.4

39.

Start by applying Yates' method. Each sum of squares is given by  $SS = (effect contrast)^2/24$ .

	Total			Effect	
Condition	X <sub>iik</sub>	1	2	Contrast	SS
(1)	315	927	2478	5485	
а	612	1551	3007	1307	SSA = 71,177.04
b	584	1163	680	1305	SSB = 70,959.38
ab	967	1844	627	199	SSAB = 1650.04
с	453	297	624	529	SSC = 11,660.04
ac	710	383	681	-53	SSAC = 117.04
bc	737	257	86	57	SSBC = 135.38
abc	1107	370	113	27	SSABC = 30.38

a. Totals appear above. From these,

 $\hat{\beta}_{1} = \overline{x}_{2.} - \overline{x}_{...} = \frac{584 + 967 + 737 + 1107 - 315 - 612 - 453 - 710}{24} = 54.38;$  $\hat{\gamma}_{11}^{AC} = \frac{315 - 612 + 584 - 967 - 453 + 710 - 737 + 1107}{24} = 2.21; \quad \hat{\gamma}_{21}^{AC} = -\hat{\gamma}_{11}^{AC} = 2.21.$ 

# Chapter 11: Multifactor Analysis of Variance

**b.** Factor sums of squares appear in the preceding table. From the original data,  $\sum \sum x_{ijkl}^2 = 1,411,889$  and x... = 5485, so SST = 1,411,889 - 5485<sup>2</sup>/24 = 158,337.96, from which SSE = 2608.7 (the remainder).

Source	df	SS	MS	F	P-value
A	1	71.177.04	71,177.04	435.65	<.001
R	06.565.4	70 959 38	70,959.38	435.22	<.001
AD	09.21	1650.04	1650.04	10.12	.006
AD	44.23	11 660 04	11.660.04	71.52	<.001
AC	10,001	117.04	117.04	0.72	.409
AC PC	1 1	135 38	135.38	0.83	.376
APC	10,00	30.38	30.38	0.19	.672
Error	16	2608.7	163.04		
Total	23	158,337.96			

*P*-values were obtained from software. Alternatively, a *P*-value less than .05 requires an *F* statistic greater than  $F_{.05,1,16} = 4.49$ . We see that the AB interaction and all the main effects are significant.

c. Yates' algorithm generates the 15 effect SS's in the ANOVA table; each SS is (effect contrast)<sup>2</sup>/48. From the original data,  $\sum \sum \sum x_{ijklm}^2 = 3,308,143$  and  $x.... = 11,956 \Rightarrow SST = 3,308,143 - 11,956^2/48$ 328,607.98. SSE is the remainder: SSE = SST - [sum of effect SS's] = ... = 4,339.33.

Source	df	SS	MS	F
A	1	136,640.02	136,640.02	1007.6
B	i	139,644,19	139,644.19	1029.8
C	i	24,616.02	24,616.02	181.5
D	i	20.377.52	20,377.52	150.3
AB	i	2.173.52	2,173.52	16.0
AD	1	2.52	2.52	0.0
AD	6 N (TEC) 19	58.52	58.52	0.4
AD PC	î.	165.02	165.02	1.2
BC	1	9.19	9.19	0.1
CD CD	1	17.52	17.52	0.1
ADC	1	42 19	42.19	0.3
ABC	1	117.19	117.19	0.9
ABD	1	188.02	188.02	1.4
ACD	1	13.02	13.02	0.1
BCD	1	204 19	204.19	1.5
ABCD	22	1 220 33	135.60	
Error	32	120 607 08	155.00	
Total	4/	520,007.90		

In this case, a *P*-value less than .05 requires an *F* statistic greater than  $F_{.05,1,32} \approx 4.15$ . Thus, all four main effects and the AB interaction effect are statistically significant at the .05 level (and no other effects are).

The accompanying ANOVA table was created using software. All F statistics are quite large (some extremely so) and all P-values are very small. So, in fact, all seven effects are statistically significant for predicting quality.

Source	df	SS	MS	F	P-value
A	1	.003906	.003906	25.00	.001
B	1	.242556	.242556	1552.36	<.001
C		.003906	.003906	25.00	.001
AR	SI THE	178506	.178506	1142.44	<.001
AC	1	.002256	.002256	14.44	.005
BC	13.21	178506	.178506	1142.44	<.001
ABC	1	.002256	.002256	14.44	.005
Error	8	.000156	.000156		
Total	15	.613144			

43.

Condition/ Effect	$SS = \frac{(contrast)^2}{16}$	F	Condition/ Effect	$SS = \frac{(contrast)^2}{16}$	F
(1)	a go aparata sa <u>an</u> at	VOT STORY	D	414.123	850.77
A	.436	<1	AD	.017	< 1
B	.099	<1	BD	.456	< 1
AB	.003	<1	ABD	.990	-
C	.109	<1	CD	2.190	4.50
AC	.078	<1	ACD	1.020	
BC	1 404	3.62	BCD	.133	
ABC	.286	-	ABCD	.004	

SSE = .286 + .990 + 1.020 + .133 + .004 = 2.433, df = 5, so MSE = .487, which forms the denominators of the F values above. A P-value less than .05 requires an F statistic greater than  $F_{.05,1,5} = 6.61$ , so only the D main effect is significant.

45.

The allocation of treatments to blocks is as given in the answer section (see back of book), with block a. #1 containing all treatments having an even number of letters in common with both ab and cd, block #2 those having an odd number in common with ab and an even number with cd, etc.

**b.** 
$$\Sigma\Sigma\Sigma\Sigma x_{ijklm}^2 = 9,035,054 \text{ and } x_{....} = 16,898, \text{ so } SST = 9,035,054 - \frac{16,898^2}{32} = 111,853.875$$
. The eight

block-replication totals are 2091 (= 618 + 421 + 603 + 449), the sum of the four observations in block #1 on replication #1), 2092, 2133, 2145, 2113, 2080, 2122, and 2122, so

$$SSB1 = \frac{2091^2}{4} + ... + \frac{2122^2}{4} - \frac{16,898^2}{32} = 898.875$$
. The effect SS's can be computed via Yates'

algorithm; those we keep appear below. SSE is computed by SST - [sum of all other SS]. MSE = 5475.75/12 = 456.3125, which forms the denominator of the F ratios below. With  $F_{.01,1,12} = 9.33$ , only

the A and B main effects are significant.

<sup>41.</sup> 

1382.7

Chapter	11:	Multifactor	Ana	lysis of	Variance
---------	-----	-------------	-----	----------	----------

Source	df	SS	F
A	1.1	12403.125	27.18
В	1	92235.125	202.13
C	1	3.125	0.01
D	1	60.500	0.13
AC	1	10.125	0.02
BC	1	91.125	0.20
AD	1	50.000	0.11
BC	1	420.500	0.92
ABC	1	3.125	0.01
ABD	1	0.500	0.00
ACD	1	200.000	0.44
BCD	1	2.000	0.00
Block	7	898.875	0.28
Error	12	5475.750	
Total	31	111853.875	

47.

- a. The third nonestimable effect is (ABCDE)(CDEFG) = ABFG. The treatments in the group containing (1) are (1), ab, cd, ce, de, fg, acf, adf, adg, aef, acg, aeg, bcg, bcf, bdf, bdg, bef, beg, abcd, abce, abde, abfg, cdfg, cefg, defg, acdef, acdeg, bcdef, bcdeg, abcdfg, abcefg, abdefg. The alias groups of the seven main effects are {A, BCDE, ACDEFG, BFG}, {B, ACDE, BCDEFG, AFG}, {C, ABDE, DEFG, ABCFG}, {D, ABCE, CEFG, ABDFG}, {E, ABCD, CDFG, ABEFG}, {F, ABCDEF, CDEG, ABG}, and {G, ABCDEG, CDEF, ABF}.
- b. 1: (1), aef, beg, abcd, abfg, cdfg, acdeg, bcdef; 2: ab, cd, fg, aeg, bef, acdef, bcdeg, abcdfg; 3: de, acg, adf, bcf, bdg, abce, cefg, abdefg; 4: ce, acf, adg, bcg, bdf, abde, defg, abcefg.

49.

		A	в	С	D	Е	AB	AC	AD	AE	BC	BD	BE	CD	CE	DE
a	70.4	+	-	-	-	-	-	-	+	-	+	+	+	+	+	+
b	72.1	-	+	-	-	-	-	+	+	+	-	-	-	+	+	+
c	70.4	_	-	+	-	-	+	-	+	+	-	+	+	-	-	+
abc	73.8	+	+	+		-	+	+	-	-	+	-	-		-	+
d	67.4	-	-	-	+	11	+	+	12	+	+	-	+	-	+	-
abd	67.0	+	+	-	+	12	+	-	+	-	-	+	-	-	+	
acd	66.6	+	1	+	+	-	-	+	+	-	-	-	+	+	-	-
hed	66.8		+	+	+	-	-	-	-	+	+	+	-	+	-	-
e	68.0	-	_	-	-	+	+	+	+	-	+	+	-	+	-	-
aha	67.8	+	+	-		+	+	-	-	+	-	-	+	+	-	-
ace	67.5	+	_	+	_	+	_	+	-	+	_	+	-	-	+	-
han	70.3	5	+	+	1	+	-	12	+	-	+	-	+	-	+	-
oce	64.0	+			+	+	-	_	+	+	+	-	-	-	-	+
ade	67.0		+	1	+	+	_	+	-	-	-	+	+	-	-	+
bde	61.9	-		+	+	+	+	1.5	_	-	-	-	-	+	+	+
abcde	68.0	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+

Thus  $SSA = \frac{(70.4 - 72.1 - 70.4 + ... + 68.0)^2}{16} = 2.250$ , SSB = 7.840, SSC = .360, SSD = 52.563, SSE = .360

10.240, SSAB = 1.563, SSAC = 7.563, SSAD = .090, SSAE = 4.203, SSBC = 2.103, SSBD = .010, SSBE = .123, SSCD = .010, SSCE = .063, SSDE = 4.840, Error SS = sum of two factor SS's = 20.568, Error MS = 2.057,  $F_{0.11,10} = 10.04$ , so only the D main effect is significant.

## **Supplementary Exercises**

-	٠.		
-			
-	_	τ.	

Source	df	SS	MS	F
А	1	322.667	322.667	980.38
В	3	35.623	11.874	36.08
AB	3	8.557	2.852	8.67
Error	16	5.266	.329	
Total	23	372.113		

We first test the null hypothesis of no interactions ( $H_{0AB}$ :  $\gamma_{ij} = 0$  for all *i*, *j*). At df = (3, 16), 5.29 < 8.67 < 9.01  $\Rightarrow$  .01 < *P*-value < .001. Therefore,  $H_0$  is rejected. Because we have concluded that interaction is present, tests for main effects are not appropriate.

53. Let A = spray volume, B = belt speed, C = brand. The Yates table and ANOVA table are below. At degrees of freedom = (1, 8), a *P*-value less than .05 requires  $F > F_{.05,1,8} = 5.32$ . So all of the main effects are significant at level .05, but none of the interactions are significant.

Condition	Total	1	2	Contrast	$SS = \frac{(contrast)^2}{16}$
(1)	76	129	289	592	21,904.00
A	53	160	303	22	30.25
В	62	143	13	48	144.00
AB	98	160	9	134	1122.25
С	88	-23	31	14	12.25
AC	55	36	17	-4	1.00
BC	59	-33	59	-14	12.25
ABC	101	42	75	16	16.00
	Effect	df	M	IS	F
	А	1	30.2	25	6.72
	В	1	144.0	00	32.00
	AB	1	1122.2	25	249.39
С		1	12.2	25	2.72
	AC	1	1.0	00	.22
	BC	1	12.2	25	2.72
	ABC	1	16.0	00	3.56
	Error	8	4.:	50	
	Total	15			

www.konkur.ir

# Chapter 11: Multifactor Analysis of Variance

### 55.

Effect	0/ Inon		-			
	7011'01	194.00	2	3	Contrast	SS
16.5	7	18	37	174	684	
А	11	19	137	510	144	1296
В	7	62	169	50	36	81
AB	12	75	341	94	0	06 (
C	21	79	9	14	272	4624
AC	41	90	41	22	32	64
BC	27	165	47	2	12	hero i
ABC	48	176	47	-2	-4	
D	28	4	1	100	336	705
AD	51	5	13	172	44	12
BD	33	20	11	32	8	linght vite
ABD	57	21	11	0	0	1
CD	70	23	1	12	72	32
ACD	95	24	1	0	-32	6
BCD	77	25	1	0	-12	lingle on a
ABCD	99	22	-3	-4	-4	
	A B AB C AC BC ABC D AD BD ABD CD ACD BCD ABCD	A       11         B       7         AB       12         C       21         AC       41         BC       27         ABC       48         D       28         AD       51         BD       33         ABD       57         CD       70         ACD       95         BCD       77         ABCD       99	A       11       19         B       7       62         AB       12       75         C       21       79         AC       41       90         BC       27       165         ABC       48       176         D       28       4         AD       51       5         BD       33       20         ABD       57       21         CD       70       23         ACD       95       24         BCD       77       25         ABCD       99       22	A       11       19       137         B       7       62       169         AB       12       75       341         C       21       79       9         AC       41       90       41         BC       27       165       47         ABC       48       176       47         D       28       4       1         AD       51       5       13         BD       33       20       11         ABD       57       21       11         CD       70       23       1         ACD       95       24       1         BCD       77       25       1         ABCD       99       22       -3	A         11         19         137         510           B         7         62         169         50           AB         12         75         341         94           C         21         79         9         14           AC         41         90         41         22           BC         27         165         47         2           ABC         48         176         47         -2           D         28         4         1         100           AD         51         5         13         172           BD         33         20         11         32           ABD         57         21         11         0           CD         70         23         1         12           ACD         95         24         1         0           BCD         77         25         1         0           BCD         99         22         -3         -4	A         11         19         137         510         144           B         7         62         169         50         36           AB         12         75         341         94         0           C         21         79         9         14         272           AC         41         90         41         22         32           BC         27         165         47         2         12           ABC         48         176         47         -2         -4           D         28         4         1         100         336           AD         51         5         13         172         44           BD         33         20         11         32         8           ABD         57         21         11         0         0           CD         70         23         1         12         72           ACD         95         24         1         0         -32           BCD         77         25         1         0         -12           ABCD         99         22         -3         <

$$\hat{\delta}_{1} = \frac{272}{16} = 17.00, \ \hat{\gamma}_{1} = \frac{336}{16} = 21.00. \text{ Similarly, } \left(\hat{\alpha\beta}\right)_{11} = 0, \ \left(\hat{\alpha\delta}\right)_{11} = 2.00, \ \left(\hat{\alpha\gamma}\right)_{11} = 2.75 \\ \left(\hat{\beta\delta}\right)_{11} = .75, \ \left(\hat{\beta\gamma}\right)_{11} = .50, \text{ and } \left(\hat{\delta\gamma}\right)_{11} = 4.50. \end{aligned}$$

The plot suggests main effects A, C, and D are quite important, and perhaps the interaction CD as well. b. In fact, pooling the 4 three-factor interaction SS's and the four-factor interaction SS to obtain an SSE based on 5 df and then constructing an ANOVA table suggests that these are the most important effects.

.



Source	df	SS	MS	F	$F_{.01, \text{ num df, den df}}$
А	2	67553	33777	11.37	5.49
В	2	72361	36181	12.18	5.49
С	2	442111	221056	74.43	5.49
AB	4	9696	2424	0.82	4.11
AC	4	6213	1553	0.52	4.11
BC	4	34928	8732	2.94	4.11
ABC	8	33487	4186	1.41	3.26
Error	27	80192	2970	1.1	
Total	53	746542		27.5	

57. The ANOVA table is:

A *P*-value less than .01 requires an *F* statistic greater than the  $F_{.01}$  value at the appropriate df (see the far right column). All three main effects are statistically significant at the 1% level, but no interaction terms are statistically significant at that level.

59. Based on the *P*-values in the ANOVA table, statistically significant factors at the level .01 are adhesive type and cure time. The conductor material does not have a statistically significant effect on bond strength. There are no significant interactions.

51.	$SSA = \sum_{i} \sum_{j} ($	$(\overline{X}_{i\ldots} - \overline{X}_{i\ldots})$	$\left(\right)^{2} = \frac{1}{N} \Sigma \lambda$	$\zeta_{i\ldots}^2 - \frac{X_{\ldots}^2}{N}$	, with simi	ilar express	ions for SSB, SSC, a	and SSD, each having
	N-1 df. $SST = \sum \sum_{n=1}^{\infty} ($	$(X_{ij(kl)} - \bar{\lambda})$	$\left(\overline{c}_{m}\right)^2 = \sum_{i=1}^{n} \left(\overline{c}_{m}\right)^2$	$\sum X_{ij(kl)}^2 -$	$\frac{X^2}{N}$ with	$N^2 - 1$ df,	leaving $N^2 - 1 - 4(N)$	(-1) df for error.
	( )	1	2	3	4	5	$\Sigma x^2$	
	<i>x</i> <sub><i>i</i></sub> :	482	446	464	468	434	1,053,916	
	<i>x</i> <sub>.j.</sub> :	470	451	440	482	451	1,053,626	
	<i>x</i> :	372	429	484	528	481	1,066,826	
	x ,:	340	417	466	537	534	1,080,170	

Also,  $\Sigma\Sigma x_{ij(kl)}^2 = 220,378$ ,  $x_{iii} = 2294$ , and CF = 210,497.44.

	Source	df	SS	MS	F
-	A	4	285.76	71.44	.594
	В	4	227.76	56.94	.473
	С	4	2867.76	716.94	5.958
	D	4	5536.56	1384.14	11.502
	Error	8	962.72	120.34	
	Total	24			

At df = (4, 8), a *P*-value less than .05 requires an *F*-statistic greater than  $F_{.05,4,8} = 3.84$ .  $H_{0A}$  and  $H_{0B}$  cannot be rejected, while  $H_{0C}$  and  $H_{0D}$  are rejected.

ww.konku

a maintain and a bits man loss most iterate being million and an

# CHAPTER 12

### Section 12.1

1.

a. Stem and Leaf display of temp:

17 0	
17 23	stem = tens
17 445	leaf = ones
17 67	
17	
18 0000011	
18 2222	
18 445	
18 6	
18 8	
	17 0 17 23 17 445 17 67 17 18 0000011 18 2222 18 445 18 6 18 8

180 appears to be a typical value for this data. The distribution is reasonably symmetric in appearance and somewhat bell-shaped. The variation in the data is fairly small since the range of values (188 - 170 = 18) is fairly small compared to the typical value of 180.

0	889	
1	0000	stem = ones
1	3	leaf = tenths
1	4444	
1	66	
1	8889	
2	11	
2		
2	5	
2	6	
2	1.28-19	
3	00	

For the ratio data, a typical value is around 1.6 and the distribution appears to be positively skewed. The variation in the data is large since the range of the data (3.08 - .84 = 2.24) is very large compared to the typical value of 1.6. The two largest values could be outliers.

b. The efficiency ratio is not uniquely determined by temperature since there are several instances in the data of equal temperatures associated with different efficiency ratios. For example, the five observations with temperatures of 180 each have different efficiency ratios.

157

c. A scatter plot of the data appears below. The points exhibit quite a bit of variation and do not appear to fall close to any straight line or simple curve.



A scatter plot of the data appears below. The points fall very close to a straight line with an intercept of approximately 0 and a slope of about 1. This suggests that the two methods are producing substantially the same concentration measurements.



For derains outer a typical solar is information of the fields on the province provincing the series of the set The series of the dama prove size and the province of the story of the second second second second second second

The efficiency relief includence determines for the project and an extinct a market of infances in the data of equal temperature above and only infantite for each other of a market for each of the off end one with consentions of the act only off councils each one.

158

C 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

3.

a. The scatter plot with axes intersecting at (0,0) is shown below.



The scatter plot with axes intersecting at (55, 100) is shown below.



c. A parabola appears to provide a good fit to both graphs.

7.

b.

www.konkur.in

5.

**a.**  $\mu_{\gamma \cdot 2500} = 1800 + 1.3(2500) = 5050$ 

- **b.** expected change = slope =  $\beta_1 = 1.3$
- c. expected change =  $100\beta_1 = 130$
- **d.** expected change =  $-100\beta_1 = -130$

9.

- **a.**  $\beta_1$  = expected change in flow rate (y) associated with a one inch increase in pressure drop (x) = .095.
- **b.** We expect flow rate to decrease by  $5\beta_1 = .475$ .

159

c. 
$$\mu_{\gamma_{10}} = -.12 + .095(10) = .83$$
, and  $\mu_{\gamma_{15}} = -.12 + .095(15) = 1.305$ .

**d.** 
$$P(Y > .835) = P\left(Z > \frac{.835 - .830}{.025}\right) = P(Z > .20) = .4207$$
.  
 $P(Y > .840) = P\left(Z > \frac{.840 - .830}{.025}\right) = P(Z > .40) = .3446$ .

e. Let  $Y_1$  and  $Y_2$  denote pressure drops for flow rates of 10 and 11, respectively. Then  $\mu_{Y_11} = .925$ , so  $Y_1 - Y_2$  has expected value .830 - .925 = -.095, and sd  $\sqrt{(.025)^2 + (.025)^2} = .035355$ . Thus  $P(Y_1 > Y_2) = P(Y_1 - Y_2 > 0) = P\left(z > \frac{0 - (-.095)}{.035355}\right) = P(Z > 2.69) = .0036$ .

11.

- a.  $\beta_1$  = expected change for a one degree increase = -.01, and  $10\beta_1$  = -.1 is the expected change for a 10 degree increase.
- **b.**  $\mu_{\gamma_{200}} = 5.00 .01(200) = 3$ , and  $\mu_{\gamma_{250}} = 2.5$ .
- c. The probability that the first observation is between 2.4 and 2.6 is

 $P(2.4 \le Y \le 2.6) = P\left(\frac{2.4 - 2.5}{.075} \le Z \le \frac{2.6 - 2.5}{.075}\right) = P(-1.33 \le Z \le 1.33) = .8164$ . The probability that any particular one of the other four observations is between 2.4 and 2.6 is also .8164, so the probability that all five are between 2.4 and 2.6 is (.8164)<sup>5</sup> = .3627.

**d.** Let  $Y_1$  and  $Y_2$  denote the times at the higher and lower temperatures, respectively. Then  $Y_1 - Y_2$  has expected value 5.00 - .01(x+1) - (5.00 - .01x) = -.01. The standard deviation of  $Y_1 - Y_2$  is

$$\sqrt{(.075)^2 + (.075)^2} = .10607$$
. Thus  $P(Y_1 - Y_2 > 0) = P\left(Z > \frac{-(-.01)}{.10607}\right) = P(Z > .09) = .4641$ .

### Section 12.2

13. For this data, 
$$n = 4$$
,  $\Sigma x_i = 200$ ,  $\Sigma y_i = 5.37$ ,  $\Sigma x_i^2 = 12.000$ ,  $\Sigma y_i^2 = 9.3501$ ,  $\Sigma x_i y_i = 333 \Rightarrow$   
 $S_{xx} = 12,000 - \frac{(200)^2}{4} = 2000$ ,  $SST = S_{yy} = 9.3501 - \frac{(5.37)^2}{4} = 2.140875$ ,  $S_{xy} = 333 - \frac{(200)(5.37)}{4} = 64.5$   
 $\Rightarrow \hat{\beta}_1 = \frac{S_{yy}}{S_{xx}} = \frac{64.5}{2000} = .03225 \Rightarrow SSE = S_{yy} - \hat{\beta}_1 S_{xy} = 2.14085 - (.03225)(64.5) = .060750$ . From these calculations,  $r^2 = 1 - \frac{SSE}{SST} = 1 - \frac{.060750}{2.14085} = .972$ . This is a very high value of  $r^2$ , which confirms the

authors' claim that there is a strong linear relationship between the two variables. (A scatter plot also shows a strong, linear relationship.)

15.

# Chapter 12: Simple Linear Regression and Correlation

a. The following stem and leaf display shows that: a typical value for this data is a number in the low 40's. There is some positive skew in the data. There are some potential outliers (79.5 and 80.0), and there is a reasonably large amount of variation in the data (e.g., the spread 80.0-29.8 = 50.2 is large compared with the typical values in the low 40's).

2	9	
3	33	stem = tens
3	5566677889	leaf = ones
4	1223	
4	56689	
5	1	
5	a di na tina tatan	
6	2	
6	9	
7	a treviseloto a	
7	9	
8	0	
	and a start of the	

- b. No, the strength values are not uniquely determined by the MoE values. For example, note that the two pairs of observations having strength values of 42.8 have different MoE values.
- c. The least squares line is  $\hat{y} = 3.2925 + .10748x$ . For a beam whose modulus of elasticity is x = 40, the predicted strength would be  $\hat{y} = 3.2925 + .10748(40) = 7.59$ . The value x = 100 is far beyond the range of the x values in the data, so it would be dangerous (i.e., potentially misleading) to extrapolate the linear relationship that far.
- **d.** From the output, SSE = 18.736, SST = 71.605, and the coefficient of determination is  $r^2$  = .738 (or 73.8%). The  $r^2$  value is large, which suggests that the linear relationship is a useful approximation to the true relationship between these two variables.

#### 17.

**a.** From software, the equation of the least squares line is  $\hat{y} = 118.91 - .905x$ . The accompanying fitted line plot shows a very strong, linear association between unit weight and porosity. So, yes, we anticipate the linear model will explain a great deal of the variation in y.



161

- **b.** The slope of the line is  $b_1 = -.905$ . A one-pcf increase in the unit weight of a concrete specimen is associated with a .905 percentage point decrease in the specimen's predicted porosity. (Note: slope is not ordinarily a percent decrease, but the units on porosity, y, are percentage points.)
- c. When x = 135, the predicted porosity is  $\hat{y} = 118.91 .905(135) = -3.265$ . That is, we get a negative prediction for y, but in actuality y cannot be negative! This is an example of the perils of extrapolation; notice that x = 135 is outside the scope of the data.
- **d.** The first observation is (99.0, 28.8). So, the actual value of y is 28.8, while the predicted value of y is 118.91 .905(99.0) = 29.315. The residual for the first observation is  $y \hat{y} = 28.8 29.315 = -.515 \approx -.52$ . Similarly, for the second observation we have  $\hat{y} = 27.41$  and residual = 27.9 27.41 = .49.
- e. From software and the data provided, a point estimate of  $\sigma$  is s = .938. This represents the "typical" size of a deviation from the least squares line. More precisely, predictions from the least squares line are "typically"  $\pm$  .938% off from the actual porosity percentage.
- f. From software,  $r^2 = 97.4\%$  or .974, the proportion of observed variation in porosity that can be attributed to the approximate linear relationship between unit weight and porosity.
- **19.** n = 14,  $\Sigma x_i = 3300$ ,  $\Sigma y_i = 5010$ ,  $\Sigma x_i^2 = 913,750$ ,  $\Sigma y_i^2 = 2,207,100$ ,  $\Sigma x_i y_i = 1,413,500$ 
  - **a.**  $\hat{\beta}_1 = \frac{3,256,000}{1,902,500} = 1.71143233$ ,  $\hat{\beta}_0 = -45.55190543$ , so the equation of the least squares line is roughly  $\hat{y} = -45.5519 + 1.7114x$ .

rouging y holoors in it an

**b.** 
$$\hat{\mu}_{y,225} = -45.5519 + 1.7114(225) = 339.51$$
.

- Estimated expected change  $= -50\hat{\beta}_1 = -85.57$ .
- **d.** No, the value 500 is outside the range of x values for which observations were available (the danger of extrapolation).
- 21.
- **a.** Yes a scatter plot of the data shows a strong, linear pattern, and  $r^2 = 98.5\%$ .
- **b.** From the output, the estimated regression line is  $\hat{y} = 321.878 + 156.711x$ , where x = absorbance and y = resistance angle. For x = .300,  $\hat{y} = 321.878 + 156.711(.300) = 368.89$ .
- c. The estimated regression line serves as an estimate both for a single y at a given x-value and for the true average  $\mu_y$  at a given x-value. Hence, our estimate for  $\mu_y$  when x = .300 is also 368.89.

#### 23.

**a.** Using the given  $y_i$ 's and the formula  $\hat{y}_i = -45.5519 + 1.7114x_i$ ,

 $SSE = (150 - 125.6)^2 + ... + (670 - 639.0)^2 = 16,213.64$ . The computation formula gives SSE = 2,207,100 - (-45.55190543)(5010) - (1.71143233)(1,413,500) = 16,205.45

**b.** 
$$SST = 2,207,100 - \frac{(5010)^2}{14} = 414,235.71$$
 so  $r^2 = 1 - \frac{16,205.45}{414,235.71} = .961$ .

25. Substitution of β̂₀ = ∑y<sub>i</sub> - β̂<sub>i</sub>Σx<sub>i</sub>/n and β̂₁ for b₀ and b₁ on the left-hand side of the first normal equation yields n ∑y<sub>i</sub> - β̂<sub>i</sub>∑x<sub>i</sub> + β̂<sub>i</sub>∑x<sub>i</sub> = ∑y<sub>i</sub> - β̂<sub>i</sub>∑x<sub>i</sub> + β̂<sub>i</sub>∑x<sub>i</sub> = ∑y<sub>i</sub>, which verifies they satisfy the first one. The same substitution in the left-hand side of the second equation gives

(∑x<sub>i</sub>)(∑y<sub>i</sub> - β̂<sub>i</sub>∑x<sub>i</sub>)/n + (∑x<sub>i</sub><sup>2</sup>)β̂<sub>i</sub> = (∑x<sub>i</sub>)(∑y<sub>i</sub>) + β̂<sub>i</sub>(n∑x<sub>i</sub><sup>2</sup> - (∑x<sub>i</sub>)<sup>2</sup>)/n = (∑x<sub>i</sub>)(∑y<sub>i</sub>)/n + β̂<sub>i</sub>(∑x<sub>i</sub><sup>2</sup> - (∑x<sub>i</sub>)<sup>2</sup>/n). The last term in parentheses is S<sub>xx</sub>, so making that substitution along with Equation (12.2) we have
(∑x<sub>i</sub>)(∑y<sub>i</sub>)/n + Ŝ<sub>xx</sub>/S<sub>xx</sub> (S<sub>xx</sub>) = (∑x<sub>i</sub>)(∑y<sub>i</sub>)/n + S<sub>xy</sub>. By the definition of S<sub>xy</sub>, this last expression is exactly ∑ x<sub>i</sub>y<sub>i</sub>, which verifies that the slope and intercept formulas satisfy the second normal equation.

27. We wish to find b₁ to minimize f(b<sub>i</sub>) = ∑(y<sub>i</sub> - b<sub>i</sub>x<sub>i</sub>)<sup>2</sup>. Equating f'(b<sub>i</sub>) to 0 yields ∑[2(y<sub>i</sub> - b<sub>i</sub>x<sub>i</sub>)<sup>1</sup>(-x<sub>i</sub>)] = 0 ⇒ 2∑[-x<sub>i</sub>y<sub>i</sub> + b<sub>i</sub>x<sub>i</sub><sup>2</sup>] = 0 ⇒ ∑x<sub>i</sub>y<sub>i</sub> = b<sub>1</sub>∑x<sub>i</sub><sup>2</sup> and b<sub>1</sub> = ∑x<sub>i</sub>y<sub>i</sub><sup>2</sup>. The least squares

estimator of  $\hat{\beta}_1$  is thus  $\hat{\beta}_1 = \frac{\sum x_i Y_i}{\sum r_i^2}$ .

29. For data set #1,  $r^2 = .43$  and s = 4.03; for #2,  $r^2 = .99$  and s = 4.03; for #3,  $r^2 = .99$  and s = 1.90. In general, we hope for both large  $r^2$  (large % of variation explained) and small s (indicating that observations don't deviate much from the estimated line). Simple linear regression would thus seem to be most effective for data set #3 and least effective for data set #1.



163 © 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

WW.KOIIKULI

### Section 12.3

a.

31.

Software output from least squares regression on this data appears below. From the output, we see that  $r^2 = 89.26\%$  or .8926, meaning 89.26% of the observed variation in threshold stress (y) can be attributed to the (approximate) linear model relationship with yield strength (x).

Regression Equation

y = 211.655 - 0.175635 x

Coefficients

Coef SE Coef T 95% CI P Term Constant 211.655 15.0622 14.0521 0.000 (178.503, 244.807) -0.176 0.0184 -9.5618 0.000 (-0.216, -0.135) x

Summary of Model

S = 6.80578 R-Sq = 89.26% R-Sq(adj) = 88.28%

**b.** From the software output,  $\hat{\beta}_1 = -0.176$  and  $s_{\hat{\alpha}} = 0.0184$ . Alternatively, the residual standard deviation is s = 6.80578, and the sum of squared deviations of the x-values can be calculated to equal  $S_{xx} =$ 

 $\sum (x_i - \overline{x})^2 = 138095$ . From these,  $s_{\beta_i} = \frac{s}{\sqrt{S_{xx}}} = .0183$  (due to some slight rounding error).

From the software output, a 95% CI for  $\beta_1$  is (-0.216, -0.135). This is a fairly narrow interval, so  $\beta_1$ c. has indeed been precisely estimated. Alternatively, with n = 13 we may construct a 95% CI for  $\beta_1$  as  $\hat{\beta}_1 \pm t_{.025,12} s_{\hat{\alpha}} = -0.176 \pm 2.179(.0184) = (-0.216, -0.136).$ 

### 33.

- Error df = n 2 = 25,  $t_{.025,25} = 2.060$ , and so the desired confidence interval is a.  $\hat{\beta}_1 \pm t_{.025,25} \cdot s_{\hat{\beta}_1} = .10748 \pm (2.060)(.01280) = (.081, .134)$ . We are 95% confident that the true average change in strength associated with a 1 GPa increase in modulus of elasticity is between .081 MPa and .134 MPa.
- **b.** We wish to test  $H_0: \beta_1 \le .1$  versus  $H_a: \beta_1 > .1$ . The calculated test statistic is

 $t = \frac{\hat{\beta}_1 - .1}{s_{\hat{\beta}_1}} = \frac{.10748 - .1}{.01280} = .58$ , which yields a *P*-value of .277 at 25 df. Thus, we fail to reject  $H_0$ ; i.e.,

there is not enough evidence to contradict the prior belief.

35.

www.konkur.in

**a.** We want a 95% CI for  $\beta_1$ : Using the given summary statistics,  $S_{xx} = 3056.69 - \frac{(222.1)^2}{17} = 155.019$ ,

$$S_{xy} = 2759.6 - \frac{(222.1)(193)}{17} = 238.112$$
, and  $\hat{\beta}_1 = \frac{S_{xy}}{S_{xx}} = \frac{238.112}{115.019} = 1.536$ . We need

 $\hat{\beta}_0 = \frac{193 - (1.536)(222.1)}{17} = -8.715$  to calculate the SSE:

SSE = 
$$2975 - (-8.715)(193) - (1.536)(2759.6) = 418.2494$$
. Then  $s = \sqrt{\frac{418.2494}{15}} = 5.28$  and

 $s_{\hat{\beta}_i} = \frac{5.28}{\sqrt{155.019}} = .424$ . With  $t_{.025,15} = 2.131$ , our CI is  $1.536 \pm 2.131 \cdot (.424) = (.632, 2.440)$ . With

95% confidence, we estimate that the change in reported nausea percentage for every one-unit change in motion sickness dose is between .632 and 2.440.

**b.** We test the hypotheses  $H_0: \beta_1 = 0$  versus  $H_a: \beta_1 \neq 0$ , and the test statistic is  $t = \frac{1.536}{.424} = 3.6226$ . With

df = 15, the two-tailed *P*-value = 2P(T > 3.6226) = 2(.001) = .002. With a *P*-value of .002, we would reject the null hypothesis at most reasonable significance levels. This suggests that there is a useful linear relationship between motion sickness dose and reported nausea.

- c. No. A regression model is only useful for estimating values of nausea % when using dosages between 6.0 and 17.6, the range of values sampled.
- **d.** Removing the point (6.0, 2.50), the new summary stats are: n = 16,  $\Sigma x_i = 216.1$ ,  $\Sigma y_i = 191.5$ ,

 $\Sigma x_i^2 = 3020.69$ ,  $\Sigma y_i^2 = 2968.75$ ,  $\Sigma x_i y_i = 2744.6$ , and then  $\hat{\beta}_1 = 1.561$ ,  $\hat{\beta}_0 = -9.118$ , SSE = 430.5264, s = 5.55,  $s_{\hat{\beta}_1} = .551$ , and the new CI is  $1.561 \pm 2.145 \cdot (.551)$ , or (.379, 2.743). The interval is a little wider. But removing the one observation did not change it that much. The observation does not seem

to be exerting undue influence.

37.

- **a.** Let  $\mu_d$  = the true mean difference in velocity between the two planes. We have 23 pairs of data that we will use to test  $H_0$ :  $\mu_d = 0$  v.  $H_a$ :  $\mu_d \neq 0$ . From software,  $\overline{x}_d = 0.2913$  with  $s_d = 0.1748$ , and so t =
  - $\frac{0.2913-0}{0.1748} \approx 8$ , which has a two-sided *P*-value of 0.000 at 22 df. Hence, we strongly reject the null

hypothesis and conclude there is a statistically significant difference in true average velocity in the two planes. [*Note:* A normal probability plot of the differences shows one mild outlier, so we have slight concern about the results of the t procedure.]

**b.** Let  $\beta_1$  denote the true slope for the linear relationship between Level – velocity and Level – velocity. We wish to test  $H_0$ :  $\beta_1 = 1$  v.  $H_a$ :  $\beta_1 < 1$ . Using the relevant numbers provided,  $t = \frac{b_1 - 1}{s(b_1)} = \frac{0.65393 - 1}{0.05947}$ 

= -5.8, which has a one-sided *P*-value at 23-2 = 21 df of  $P(T < -5.8) \approx 0$ . Hence, we strongly reject the null hypothesis and conclude the same as the authors; i.e., the true slope of this regression relationship is significantly less than 1.

Source	df	SS	MS	f
Regr	1	31.860	31.860	18.0
Error	18	7.968	1.77	Sec. 12
Total	19	39.828		

**39.** SSE = 124,039.58- (72.958547)(1574.8) - (.04103377)(222657.88) = 7.9679, and SST = 39.828

At df = (1, 18),  $f = 18.0 > F_{.001,1,18} = 15.38$  implies that the *P*-value is less than .001. So,  $H_0: \beta_1 = 0$  is rejected and the model is judged useful. Also,  $s = \sqrt{1.77} = 1.33041347$  and  $S_{xx} = 18,921.8295$ , so

 $t = \frac{.04103377}{1.33041347 / \sqrt{18,921.8295}} = 4.2426$  and  $t^2 = (4.2426)^2 = 18.0 = f$ , showing the equivalence of the two tests.

41.

**a.** Under the regression model,  $E(Y_i) = \beta_0 + \beta_1 x_i$  and, hence,  $E(\overline{Y}) = \beta_0 + \beta_1 \overline{x}$ . Therefore,  $E(Y_i - \overline{Y}) = \beta_1(x_i - \overline{x})$ , and  $E(\hat{\beta}_1) = E\left[\frac{\sum (x_i - \overline{x})(Y_i - \overline{Y})}{\sum (x_i - \overline{x})^2}\right] = \frac{\sum (x_i - \overline{x})E[Y_i - \overline{Y}]}{\sum (x_i - \overline{x})^2}$  $= \frac{\sum (x_i - \overline{x})\beta_1(x_i - \overline{x})}{\sum (x_i - \overline{x})^2} = \beta_1 \frac{\sum (x_i - \overline{x})^2}{\sum (x_i - \overline{x})^2} = \beta_1.$ 

**b.** Here, we'll use the fact that  $\sum (x_i - \overline{x})(Y_i - \overline{Y}) = \sum (x_i - \overline{x})Y_i - \overline{Y}\sum (x_i - \overline{x}) = \sum (x_i - \overline{x})Y_i - \overline{Y}(0) = \sum (x_i - \overline{x})Y_i$ . With  $c = \sum (x_i - \overline{x})^2$ ,  $\hat{\beta}_i = \frac{1}{c}\sum (x_i - \overline{x})(Y_i - \overline{Y}) = \sum \frac{(x_i - \overline{x})}{c}Y_i \Rightarrow$  since the  $Y_i$ s are independent,  $V(\hat{\beta}_i) = \sum \left(\frac{x_i - \overline{x}}{c}\right)^2 V(Y_i) = \frac{1}{c^2}\sum (x_i - \overline{x})^2 \sigma^2 = \frac{\sigma^2}{c} = \frac{\sigma^2}{\sum (x_i - \overline{x})^2}$  or, equivalently,  $\frac{\sigma^2}{\sum x^2 - (\sum x_i)^2/n}$ , as desired.

43. The numerator of d is |1 - 2| = 1, and the denominator is  $\frac{4\sqrt{14}}{\sqrt{324.40}} = .831$ , so  $d = \frac{1}{.831} = 1.20$ . The approximate power curve is for n - 2 df = 13, and  $\beta$  is read from Table A.17 as approximately .1.

166

## Section 12.4

45.

www.konkur.in

**a.** We wish to find a 90% CI for  $\mu_{y,125}$ :  $\hat{y}_{125} = 78.088$ ,  $t_{.05,18} = 1.734$ , and

 $s_{\bar{p}} = s_{\sqrt{\frac{1}{20} + \frac{(125 - 140.895)^2}{18,921.8295}}} = .1674$ . Putting it together, we get  $78.088 \pm 1.734(.1674) = (77.797, 78.378)$ .

**b.** We want a 90% PI. Only the standard error changes:  $s_{y} = s \sqrt{1 + \frac{1}{20} + \frac{(125 - 140.895)^2}{18,921.8295}} = .6860$ , so the PI is

 $78.088 \pm 1.734(.6860) = (76.898, 79.277).$ 

- c. Because the x of 115 is farther away from  $\overline{x}$  than the previous value, the term  $(x^* \overline{x})^2$  will be larger, making the standard error larger, and thus the width of the interval is wider.
- **d.** We would be testing to see if the filtration rate were 125 kg-DS/m/h, would the average moisture content of the compressed pellets be less than 80%. The test statistic is  $t = \frac{78.088 80}{.1674} = -11.42$ , and with 18 df the *P*-value is  $P(T < -11.42) \approx 0.00$ . Hence, we reject  $H_0$ . There is significant evidence to prove that the true average moisture content when filtration rate is 125 is less than 80%.

47.

**a.**  $\hat{y}_{(40)} = -1.128 + .82697(40) = 31.95$ ,  $t_{.025,13} = 2.160$ ; a 95% PI for runoff is  $31.95 \pm 2.160\sqrt{(5.24)^2 + (1.44)^2} = 31.95 \pm 11.74 = (20.21, 43.69)$ . No, the resulting interval is very wide, therefore the available information is not very precise.

**b.**  $\Sigma x = 798, \Sigma x^2 = 63,040$  which gives  $S_{xx} = 20,586.4$ , which in turn gives

 $s_{\hat{y}_{(50)}} = 5.24 \sqrt{\frac{1}{15} + \frac{(50 - 53.20)^2}{20,586.4}} = 1.358$ , so the PI for runoff when x = 50 is

 $40.22 \pm 2.160\sqrt{(5.24)^2 + (1.358)^2} = 40.22 \pm 11.69 = (28.53,51.92)$ . The simultaneous prediction level for the two intervals is at least  $100(1-2\alpha)\% = 90\%$ .

**49.** 95% CI = (462.1, 597.7) 
$$\Rightarrow$$
 midpoint = 529.9;  $t_{.025,8} = 2.306 \Rightarrow 529.9 + (2.306) \hat{s}_{\hat{\beta}_0 + \hat{\beta}_1(15)} = 597.7 \Rightarrow$   
 $\hat{s}_{\hat{\alpha}_1 + \hat{\alpha}_1(15)} = 29.402 \Rightarrow 99\%$  CI = 529.9  $\pm t_{.005,8} (29.402) = 529.9 \pm (3.355)(29.402) = (431.3, 628.5)$ .

51.

**a.** 0.40 is closer to  $\overline{x}$ .

**b.** 
$$\hat{\beta}_0 + \hat{\beta}_1(0.40) \pm t_{a/2, n-2} \hat{s}_{\hat{n}_1 + \hat{n}_1(0.40)}$$
 or  $0.8104 \pm 2.101(0.0311) = (0.745, 0.876).$ 

c. 
$$\hat{\beta}_0 + \hat{\beta}_1 (1.20) \pm t_{a/2, n-2} \cdot \sqrt{s^2 + s^2}_{\hat{\beta}_0 + \hat{\beta}_1 (1.20)}$$
 or  $0.2912 \pm 2.101 \cdot \sqrt{(0.1049)^2 + (0.0352)^2} = (.059, .523).$ 

53.

Choice  $\mathbf{a}$  will be the smallest, with  $\mathbf{d}$  being largest. The width of interval  $\mathbf{a}$  is less than  $\mathbf{b}$  and  $\mathbf{c}$  (obviously), and  $\mathbf{b}$  and  $\mathbf{c}$  are both smaller than  $\mathbf{d}$ . Nothing can be said about the relationship between  $\mathbf{b}$  and  $\mathbf{c}$ .

55. 
$$\hat{\beta}_{0} + \hat{\beta}_{1}x^{*} = (\overline{Y} - \hat{\beta}_{1}\overline{x}) + \hat{\beta}_{1}x^{*} = \overline{Y} + (x^{*} - \overline{x})\hat{\beta}_{1} = \frac{1}{n}\sum Y_{i} + \frac{(x^{*} - \overline{x})\sum(x_{i} - \overline{x})Y_{i}}{S_{xx}} = \sum d_{i}Y_{i}, \text{ where}$$
$$d_{i} = \frac{1}{n} + \frac{(x^{*} - \overline{x})}{S_{xx}}(x_{i} - \overline{x}). \text{ Thus, since the } Y_{i}\text{s are independent,}$$
$$V(\hat{\beta}_{0} + \hat{\beta}_{1}x) = \sum d_{i}^{2}V(Y_{i}) = \sigma^{2}\sum d_{i}^{2}$$
$$= \sigma^{2}\sum \left[\frac{1}{n^{2}} + 2\frac{(x^{*} - \overline{x})(x_{i} - \overline{x})}{nS_{xx}} + \frac{(x^{*} - \overline{x})^{2}(x_{i} - \overline{x})^{2}}{nS_{xx}^{2}}\right]$$
$$= \sigma^{2}\left[n\frac{1}{n^{2}} + 2\frac{(x^{*} - \overline{x})\sum(x_{i} - \overline{x})}{nS_{xx}} + \frac{(x^{*} - \overline{x})^{2}\sum(x_{i} - \overline{x})^{2}}{S_{xx}^{2}}\right]$$
$$= \sigma^{2}\left[\frac{1}{n} + 2\frac{(x^{*} - \overline{x})\cdot 0}{nS_{xx}} + \frac{(x^{*} - \overline{x})^{2}S_{xx}}{S_{xx}^{2}}\right] = \sigma^{2}\left[\frac{1}{n} + \frac{(x^{*} - \overline{x})^{2}}{S_{xx}}\right]$$

### Section 12.5

a

57. Most people acquire a license as soon as they become eligible. If, for example, the minimum age for obtaining a license is 16, then the time since acquiring a license, y, is usually related to age by the equation  $y \approx x - 16$ , which is the equation of a straight line. In other words, the majority of people in a sample will have y values that closely follow the line y = x - 16.

59.

$$S_{xx} = 251,970 - \frac{(1950)^{2}}{18} = 40,720 , S_{yy} = 130.6074 - \frac{(47.92)^{2}}{18} = 3.033711 , \text{ and}$$
$$S_{xy} = 5530.92 - \frac{(1950)(47.92)}{18} = 339.586667 , \text{ so } r = \frac{339.586667}{\sqrt{40,720}\sqrt{3.033711}} = .9662 . \text{ There is a very}$$

strong, positive correlation between the two variables.

- **b.** Because the association between the variables is positive, the specimen with the larger shear force will tend to have a larger percent dry fiber weight.
- c. Changing the units of measurement on either (or both) variables will have no effect on the calculated value of r, because any change in units will affect both the numerator and denominator of r by exactly the same multiplicative constant.
- **d.**  $r^2 = .9662^2 = .933$ , or 93.3%.
- e. We wish to test  $H_0: \rho = 0$  v.  $H_a: \rho > 0$ . The test statistic is  $t = \frac{r\sqrt{n-2}}{\sqrt{1-r^2}} = \frac{.9662\sqrt{18-2}}{\sqrt{1-.9662^2}} = 14.94$ . This is

"off the charts" at 16 df, so the one-tailed *P*-value is less than .001. So,  $H_0$  should be rejected: the data indicate a positive linear relationship between the two variables.

**a.** We are testing  $H_0: \rho = 0$  v.  $H_a: \rho > 0$ . The correlation is  $r = \frac{7377.704}{\sqrt{36.9839}\sqrt{2,628,930.359}} = .7482$ , and

the test statistic is  $t = \frac{.7482\sqrt{12}}{\sqrt{1-.7482^2}} \approx 3.9$ . At 14 df, the *P*-value is roughly .001. Hence, we reject  $H_0$ :

there is evidence that a positive correlation exists between maximum lactate level and muscular endurance.

- **b.** We are looking for  $r^2$ , the coefficient of determination:  $r^2 = (.7482)^2 = .5598$ , or about 56%. It is the same no matter which variable is the predictor.
- 63. With the aid of software, the sample correlation coefficient is r = .7729. To test  $H_0: \rho = 0$  v.  $H_a: \rho \neq 0$ , the test statistic is  $t = \frac{(.7729)\sqrt{6-2}}{\sqrt{1-(.7729)^2}} = 2.44$ . At 4 df, the 2-sided *P*-value is about 2(.035) = .07 (software gives

a *P*-value of .072). Hence, we fail to reject  $H_0$ : the data do not indicate that the population correlation coefficient differs from 0. This result may seem surprising due to the relatively large size of r (.77), however, it can be attributed to a small sample size (n = 6).

#### 65.

www.konkur.in

61.

**a.** From the summary statistics provided, a point estimate for the population correlation coefficient  $\rho$  is  $r \sum_{i=1}^{n} \sum_{j=1}^{n} \sum_{i=1}^{n} \sum_{i=1}^{n}$ 

$$=\frac{\sum (x_i - \bar{x})(y_i - \bar{y})}{\sqrt{\sum (x_i - \bar{x})^2 \sum (y_i - \bar{y})^2}} = \frac{44,185.67}{\sqrt{(64,732.83)(130,566.96)}} = .486$$

**b.** The hypotheses are  $H_0$ :  $\rho = 0$  versus  $H_a$ :  $\rho \neq 0$ . Assuming bivariate normality, the test statistic value is

 $t = \frac{r\sqrt{n-2}}{\sqrt{1-r^2}} = \frac{.4806\sqrt{15-2}}{\sqrt{1-.4806^2}} = 1.98. \text{ At df} = 15-2 = 13, \text{ the two-tailed } P \text{-value for this } t \text{ test is } 2P(T_{13})$ 

 $\geq 1.98$ )  $\approx 2P(T_{13} \geq 2.0) = 2(.033) = .066$ . Hence, we fail to reject  $H_0$  at the .01 level; there is not sufficient evidence to conclude that the population correlation coefficient between internal and external rotation velocity is not zero.

- **c.** If we tested  $H_0: \rho = 0$  versus  $H_a: \rho > 0$ , the one-sided *P*-value would be .033. We would still fail to reject  $H_0$  at the .01 level, lacking sufficient evidence to conclude a positive true correlation coefficient. However, for a one-sided test at the .05 level, we would reject  $H_0$  since *P*-value = .033 < .05. We have evidence at the .05 level that the true population correlation coefficient between internal and external rotation velocity is positive.
- 67.

**a.** Because *P*-value =  $.00032 < \alpha = .001$ ,  $H_0$  should be rejected at this significance level.

**b.** Not necessarily. For such a large *n*, the test statistic *t* has approximately a standard normal distribution when  $H_0$ :  $\rho = 0$  is true, and a *P*-value of .00032 corresponds to  $z = \pm 3.60$ . Solving  $\pm 3.60 = \frac{r\sqrt{500-2}}{\sqrt{1-r^2}}$ 

for r yields  $r = \pm .159$ . That is, with n = 500 we'd obtain this P-value with  $r = \pm .159$ . Such an r value suggests only a weak linear relationship between x and y, one that would typically have little practical importance.

c. The test statistic value would be  $t = \frac{.022\sqrt{10,000-2}}{\sqrt{1-.022^2}} = 2.20$ ; since the test statistic is again

approximately normal, the 2-sided *P*-value would be roughly  $2[1 - \Phi(2.20)] = .0278 < .05$ , so  $H_0$  is rejected in favor of  $H_a$  at the .05 significance level. The value t = 2.20 is statistically significant — it cannot be attributed just to sampling variability in the case  $\rho = 0$ . But with this enormous n, r = .022 implies  $\rho \approx .022$ , indicating an extremely weak relationship.

### Supplementary Exercises

Use available software for all calculations.

61.9

a.

69.

- We want a confidence interval for  $\beta_1$ . From software,  $b_1 = 0.987$  and  $s(b_1) = 0.047$ , so the corresponding 95% CI is  $0.987 \pm t_{.025,17}(0.047) = 0.987 \pm 2.110(0.047) = (0.888, 1.086)$ . We are 95% confident that the true average change in sale price associated with a one-foot increase in truss height is between \$0.89 per square foot and \$1.09 per square foot.
- **b.** Using software, a 95% CI for  $\mu_{y,25}$  is (47.730, 49.172). We are 95% confident that the true average sale price for all warehouses with 25-foot truss height is between \$47.73/ft<sup>2</sup> and \$49.17/ft<sup>2</sup>.
- c. Again using software, a 95% PI for Y when x = 25 is (45.378, 51.524). We are 95% confident that the sale price for a single warehouse with 25-foot truss height will be between \$45.38/ft<sup>2</sup> and \$51.52/ft<sup>2</sup>.
- **d.** Since x = 25 is nearer the mean than x = 30, a PI at x = 30 would be wider.
- e. From software,  $r^2 = SSR/SST = 890.36/924.44 = .963$ . Hence,  $r = \sqrt{.963} = .981$ .
- 71. Use software whenever possible.
  - **a.** From software, the estimated coefficients are  $\hat{\beta}_1 = 16.0593$  and  $\hat{\beta}_0 = 0.1925$ .

**b.** Test  $H_0$ :  $\beta_1 = 0$  versus  $H_a$ :  $\beta_1 \neq 0$ . From software, the test statistic is  $t = \frac{16.0593 - 0}{0.2965} = 54.15$ ; even at just 7 df, this is "off the charts" and the *P*-value is  $\approx 0$ . Hence, we strongly reject  $H_0$  and conclude that

c. From software or by direct computation, residual sd = s = .2626,  $\overline{x} = .408$  and  $S_{xx} = .784$ . When  $x = x^* = .2$ ,  $\hat{y} = 0.1925 + 16.0593(.2) = 3.404$  with an estimated standard deviation of

$$s_{\hat{y}} = s_{\sqrt{\frac{1}{n} + \frac{(x^* - \overline{x})^2}{S_{xx}}}} = .2626\sqrt{\frac{1}{9} + \frac{(.2 - .408)^2}{.784}} = .107$$
. The analogous calculations when  $x = x^* = .4$ 

result in  $\hat{y} = 6.616$  and  $s_{\hat{y}} = .088$ , confirming what's claimed. Prediction error is larger when x = .2 because .2 is farther from the sample mean of .408 than is x = .4.

**d.** A 95% CI for  $\mu_{y,4}$  is  $\hat{y} \pm t_{.025,9-2} s_{\hat{y}} = 6.616 \pm 2.365(.088) = (6.41, 6.82).$ 

a statistically significant relationship exists between the variables.

e. A 95% PI for Y when x = .4 is  $\hat{y} \pm t_{.025.9-2} \sqrt{s^2 + s_{\hat{y}}^2} = (5.96, 7.27).$ 

**a.** From the output,  $r^2 = .5073$ .

www.konkur.ir

73.

75.

**b.** 
$$r = (\text{sign of slope}) \cdot \sqrt{r^2} = +\sqrt{.5073} = .7122.$$

- c. We test  $H_0: \beta_1 = 0$  versus  $H_a: \beta_1 \neq 0$ . The test statistic t = 3.93 gives *P*-value = .0013, which is < .01, the given level of significance, therefore we reject  $H_0$  and conclude that the model is useful.
- **d.** We use a 95% CI for  $\mu_{\gamma.50}$ .  $\hat{y}(50) = .787218 + .007570(50) = 1.165718; t.025, 15 = 2.131;$

s = "Root MSE" = .20308  $\Rightarrow s_{\hat{y}} = .20308 \sqrt{\frac{1}{17} + \frac{17(50 - 42.33)^2}{17(41,575) - (719.60)^2}} = .051422$ . The resulting 95% CI is 1.165718 ± 2.131(.051422) = 1.165718 ± .109581 = (1.056137, 1.275299).

- e. Our prediction is  $\hat{y}(30) = .787218 + .007570(30) = 1.0143$ , with a corresponding residual of  $y \hat{y} = .80 1.0143 = -.2143$ .
- **a.** With y = stride rate and x = speed, we have  $\hat{\beta}_1 = \frac{S_{xy}}{S_{xx}} = \frac{660.130 (205.4)(35.16)/11}{3880.08 (205.4)^2/11} = \frac{3.597}{44.702} =$

0.080466 and  $\hat{\beta}_0 = \bar{y} - \hat{\beta}_1 \bar{x} = (35.16/11) - 0.080466(205.4)/11 = 1.694$ . So, the least squares line for predicting stride rate from speed is  $\hat{y} = 1.694 + 0.080466x$ .

**b.** With  $y = \text{speed and } x = \text{stride rate, we have } \hat{\beta}_1 = \frac{S_{xy}}{S_{xx}} = \frac{660.130 - (35.16)(205.4)/11}{112.681 - (35.16)^2/11} = \frac{3.597}{0.297} =$ 

12.117 and  $\hat{\beta}_0 = \overline{y} - \hat{\beta}_1 \overline{x} = (205.4)/11 - 12.117(35.16/11) = -20.058$ . So, the least squares line for predicting speed from stride rate is  $\hat{y} = -20.058 + 12.117x$ .

- c. The fastest way to find  $r^2$  from the available information is  $r^2 = \hat{\beta}_1^2 \frac{S_{xx}}{S_{yy}}$ . For the first regression, this gives  $r^2 = (0.080466)^2 \frac{44.702}{0.297} \approx .97$ . For the second regression,  $r^2 = (12.117)^2 \frac{0.297}{44.702} \approx .97$  as well. In fact, rounding error notwithstanding, these two  $r^2$  values should be exactly the same.
- 77.
- **a.** Yes: the accompanying scatterplot suggests an extremely strong, positive, linear relationship between the amount of oil added to the wheat straw and the amount recovered.

171

**b.** Pieces of Minitab output appear below. From the output,  $r^2 = 99.6\%$  or .996. That is, 99.6% of the total variation in the amount of oil recovered in the wheat straw can be explained by a linear regression on the amount of oil added to it.

Coef SE Coef Т Predictor -0.5234 0.1453 -3.60 0.003 Constant 0.87825 0.01610 54.56 0.000 x S = 0.311816 R-Sq = 99.6% R-Sq(adj) = 99.5% Predicted Values for New Observations New Obs Fit SE Fit 95% CI 95% PI 3.8678 0.0901 (3.6732, 4.0625) (3.1666, 4.5690)

- Refer to the preceding Minitab output. A test of  $H_0$ :  $\beta_1 = 0$  versus  $H_a$ :  $\beta_1 \neq 0$  returns a test statistic of c. t = 54.56 and a P-value of  $\approx 0$ , from which we can strongly reject  $H_0$  and conclude that a statistically significant linear relationship exists between the variables. (No surprise, based on the scatterplot!)
- d. The last line of the preceding Minitab output comes from requesting predictions at x = 5.0 g. The resulting 95% PI is (3.1666, 4.5690). So, at a 95% prediction level, the amount of oil recovered from wheat straw when the amount added was 5.0 g will fall between 3.1666 g and 4.5690 g.
- A formal test of  $H_0$ :  $\rho = 0$  versus  $H_a$ :  $\rho \neq 0$  is completely equivalent to the t test for slope conducted in e. c. That is, the test statistic and P-value would once again be t = 54.56 and  $P \approx 0$ , leading to the conclusion that  $\rho \neq 0$ .

79. Start with the alternative formula SSE = 
$$\Sigma y^2 - \hat{\beta}_0 \Sigma y - \hat{\beta}_1 \Sigma x y$$
. Substituting  $\hat{\beta}_0 = \frac{\Sigma y - \hat{\beta}_1 \Sigma x}{n}$ ,  
SSE =  $\Sigma y^2 - \frac{\Sigma y - \hat{\beta}_1 \Sigma x}{n} \Sigma y - \hat{\beta}_1 \Sigma x y = \Sigma y^2 - \frac{(\Sigma y)^2}{n} + \frac{\hat{\beta}_1 \Sigma x \Sigma y}{n} - \hat{\beta}_1 \Sigma x y = \left[ \Sigma y^2 - \frac{(\Sigma y)^2}{n} \right] - \hat{\beta}_1 \left[ \Sigma x y - \frac{\Sigma x \Sigma y}{n} \right]$ 

1

 $= S_{w} - \beta_1 S_{w}$ 

**a.** Recall that  $r = \frac{S_{xy}}{\sqrt{S_{xx}S_{yy}}}$ ,  $s_x^2 = \frac{\Sigma(x_i - \overline{x})^2}{n-1} = \frac{S_{xx}}{n-1}$ , and similarly  $s_y^2 = \frac{S_{yy}}{n-1}$ . Using these formulas,  $\hat{\beta}_1 = \frac{S_{xy}}{S_{xy}} = \frac{r \cdot \sqrt{S_{xx}S_{yy}}}{S_{xy}} = r \cdot \sqrt{\frac{S_{yy}}{S_{yy}}} = r \cdot \sqrt{\frac{(n-1)s_y^2}{(n-1)s_y^2}} = r \cdot \frac{s_y}{s_y}.$  Using the fact that  $\hat{\beta}_0 = \overline{y} - \hat{\beta}_1 \overline{x}$ , the least

squares equation becomes  $\hat{y} = \hat{\beta}_0 + \hat{\beta}_1 x = \overline{y} + \hat{\beta}_1 (x - \overline{x}) = \overline{y} + r \cdot \frac{s_y}{s} (x - \overline{x})$ , as desired.

b. In Exercise 64, r = .700. So, a specimen whose UV transparency index is 1 standard deviation below average is predicted to have a maximum prevalence of infection that is .7 standard deviations below average.

Remember that SST =  $S_{yy}$  and use Exercise 79 to write SSE =  $S_{yy} - \hat{\beta}_1 S_{xy} = S_{yy} - S_{xy}^2 / S_{xx}$ . Then 83.

$$r^{2} = \frac{S_{xy}^{2}}{S_{xx}S_{yy}} = \frac{S_{xy}^{2}/S_{xx}}{S_{yy}} = \frac{S_{yy}-SSE}{S_{yy}} = 1 - \frac{SSE}{S_{yy}} = 1 - \frac{SSE}{SST}.$$

#### Using Minitab, we create a scatterplot to see if a linear regression model is appropriate. 85.



A linear model is reasonable; although it appears that the variance in y gets larger as x increases. The Minitab output follows:

The regressio blood glucose	n equation level = 3	is .70 + 0.0379	time		
Predictor Constant time	Coef 3.6965 0.037895	StDev 0.2159 0.006137	T 17.12 6.17	P 0.000 0.000	
s = 0.5525	R-Sq =	63.4% R-S	sq(adj) = 6	1.7%	
Analysis of V	ariance				
Source Regression Residual Erro Total	DF 1 0r 22 23	SS 11.638 6.716 18.353	MS 11.638 0.305	F 38.12	P 0.000

The coefficient of determination of 63.4% indicates that only a moderate percentage of the variation in y can be explained by the change in x. A test of model utility indicates that time is a significant predictor of blood glucose level. ( $t = 6.17, P \approx 0$ ). A point estimate for blood glucose level when time = 30 minutes is 4.833%. We would expect the average blood glucose level at 30 minutes to be between 4.599 and 5.067, with 95% confidence.

From the SAS output in Exercise 73,  $n_1 = 17$ , SSE<sub>1</sub> = 0.61860,  $\hat{\beta}_1 = 0.007570$ ; by direct computation, 87. SS<sub>x1</sub> = 11,114.6. The pooled estimated variance is  $\hat{\sigma}^2 = \frac{.61860 + .51350}{17 + 15 - 4} = .040432$ , and the calculated test

statistic for testing  $H_0$ :  $\beta_1 = \gamma_1$  is

 $\sqrt{.040432}\sqrt{\frac{1}{11114.6} + \frac{1}{7152.5578}} \approx 0.24$ . At 28 df, the two-tailed *P*-value is roughly 2(.39) = .78. h such a large *P*-value word t =

With such a large *P*-value, we do not reject  $H_0$  at any reasonable level (in particular, .78 > .05). The data do not provide evidence that the expected change in wear loss associated with a 1% increase in austentite content is different for the two types of abrasive — it is plausible that  $\beta_1 = \gamma_1$ .

# **CHAPTER 13**

## Section 13.1

- **a.**  $\overline{x} = 15$  and  $\sum (x_i \overline{x})^2 = 250$ , so the standard deviation of the residual  $Y_i \hat{Y}_i$  is  $10\sqrt{1 \frac{1}{5} \frac{(x_i 15)^2}{250}} = 6.32, 8.37, 8.94, 8.37$ , and 6.32 for i = 1, 2, 3, 4, 5.
- **b.** Now  $\overline{x} = 20$  and  $\sum_{i=1,2,3,4,5}^{\infty} (x_i \overline{x})^2 = 1250$ , giving residual standard deviations 7.87, 8.49, 8.83, 8.94, and 2.83 for i = 1, 2, 3, 4, 5.
- c. The deviation from the estimated line is likely to be much smaller for the observation made in the experiment of b for x = 50 than for the experiment of a when x = 25. That is, the observation (50, Y) is more likely to fall close to the least squares line than is (25, Y).

#### 3.

1.

a. This plot indicates there are no outliers, the variance of ε is reasonably constant, and the ε are normally distributed. A straight-line regression function is a reasonable choice for a model.



**b.** We need  $S_{xx} = \sum (x_i - \overline{x})^2 = 415,914.85 - \frac{(2817.9)^2}{20} = 18,886.8295$ . Then each  $e_i^*$  can be calculated as follows:  $e_i^* = \frac{e_i}{.4427\sqrt{1 + \frac{1}{20} + \frac{(x_i - 140.895)^2}{18,886.8295}}}$ . The table below shows the values.

Notice that if  $e_i^* \approx e_i / s$ , then  $e_i / e_i^* \approx s$ . All of the  $e_i / e_i^*$ 's range between .57 and .65, which are close to s.
standardized residuals	$e_i/e_i^*$	standardized residuals	$e_i / e_i^*$
-0.31064	0.644053	0.6175	0.64218
-0 30593	0.614697	0.09062	0.64802
0.4791	0.578669	1.16776	0.565003
1 2307	0.647714	-1.50205	0.646461
1 15021	0.648002	0.96313	0.648257
0 34881	0.643706	0.019	0.643881
0.09972	0.633428	0.65644	0.584858
-0.09872	0.630420	-2.1562	0.647182
-1.39034	0.640035	-0.79038	0.642113
-0.15998	0.621857	1.73943	0.631795

c. This plot looks very much the same as the one in part a.



5.

a. 97.7% of the variation in ice thickness can be explained by the linear relationship between it and elapsed time. Based on this value, it is tempting to assume an approximately linear relationship; however, r<sup>2</sup> does not measure the aptness of the linear model.

b. The residual plot shows a curve in the data, suggesting a non-linear relationship exists. One observation (5.5, -3.14) is extreme.



175

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

a. From software and the data provided, the least squares line is  $\hat{y} = 84.4 - 290x$ . Also from software, the coefficient of determination is  $r^2 = 77.6\%$  or .776.

#### Regression Analysis: y versus x

The regression equation is y = 84.4 - 290 x Predictor Coef SE Coef T P Constant 84.38 11.64 7.25 0.000 x -289.79 43.12 -6.72 0.000 S = 2.72669 R-Sq = 77.6% R-Sq(adj) = 75.9%

**b.** The accompanying scatterplot exhibits substantial curvature, which suggests that a straight-line model is not actually a good fit.



c. Fits, residuals, and standardized residuals were computed using software and the accompanying plot was created. The residual-versus-fit plot indicates very strong curvature but not a lack of constant variance. This implies that a linear model is inadequate, and a quadratic (parabolic) model relationship might be suitable for x and y.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

9. Both a scatter plot and residual plot (based on the simple linear regression model) for the first data set suggest that a simple linear regression model is reasonable, with no pattern or influential data points which would indicate that the model should be modified. However, scatter plots for the other three data sets reveal difficulties.



For data set #2, a quadratic function would clearly provide a much better fit. For data set #3, the relationship is perfectly linear except one outlier, which has obviously greatly influenced the fit even though its x value is not unusually large or small. One might investigate this observation to see whether it was mistyped and/or it merits deletion. For data set #4 it is clear that the slope of the least squares line has been determined entirely by the outlier, so this point is extremely influential. A linear model is completely inappropriate for data set #4.

177

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

www.konkur.in

**a.** 
$$Y_i - \hat{Y}_i = Y_i - \overline{Y} - \hat{\beta}_i \left(x_i - \overline{x}\right) = Y_i - \frac{1}{n} \sum_j Y_j - \frac{\left(x_i - \overline{x}\right) \sum_j \left(x_j - \overline{x}\right) Y_j}{\sum_j \left(x_j - \overline{x}\right)^2} = \sum_j c_j Y_j$$
, where  
 $c_j = 1 - \frac{1}{n} - \frac{\left(x_i - \overline{x}\right)^2}{n\Sigma \left(x_j - \overline{x}\right)^2}$  for  $j = i$  and  $c_j = 1 - \frac{1}{n} - \frac{\left(x_i - \overline{x}\right) \left(x_j - \overline{x}\right)}{\Sigma \left(x_j - \overline{x}\right)^2}$  for  $j \neq i$ . Thus  
 $V\left(Y_i - \hat{Y}_i\right) = \Sigma V\left(c_j Y_j\right)$  (since the  $Y_j$ 's are independent) =  $\sigma^2 \Sigma c_j^2$  which, after some algebra, gives

13.

Equation (13.2).

**b.** 
$$\sigma^2 = V(Y_i) = V\left(\hat{Y}_i + \left(Y_i - \hat{Y}_i\right)\right) = V(\hat{Y}_i) + V\left(Y_i - \hat{Y}_i\right)$$
, so  
 $V\left(Y_i - \hat{Y}_i\right) = \sigma^2 - V(\hat{Y}_i) = \sigma^2 - \sigma^2 \left[\frac{1}{n} + \frac{(x_i - \overline{x})^2}{\Sigma\left(x_j - \overline{x}\right)^2}\right]$ , which is exactly (13.2)

c. As  $x_i$  moves further from  $\overline{x}$ ,  $(x_i - \overline{x})^2$  grows larger, so  $V(\hat{Y}_i)$  increases since  $(x_i - \overline{x})^2$  has a positive sign in  $V(\hat{Y}_i)$ , but  $V(Y_i - \hat{Y}_i)$  decreases since  $(x_i - \overline{x})^2$  has a negative sign in that expression.

The distribution of any particular standardized residual is also a *t* distribution with n - 2 d.f., since  $e_i^*$  is obtained by taking standard normal variable  $\frac{Y_i - \hat{Y}_i}{\sigma_{Y_i - \hat{Y}_i}}$  and substituting the estimate of  $\sigma$  in the denominator (exactly as in the predicted value case). With  $E_i^*$  denoting the *i*<sup>th</sup> standardized residual as a random

variable, when  $n = 25 E_i^*$  has a *t* distribution with 23 df and  $t_{.01,23} = 2.50$ , so  $P(E_i^* \text{ outside } (-2.50, 2.50)) = P(E_i^* \ge 2.50) + P(E_i^* \le -2.50) = .01 + .01 = .02$ .

## Chapter 13: Nonlinear and Multiple Regression

## Section 13.2

- a. The scatterplot of y versus x below, left has a curved pattern. A linear model would <u>not</u> be appropriate.
- **b.** The scatterplot of ln(y) versus ln(x) below, right exhibits a strong linear pattern.



- c. The linear pattern in **b** above would indicate that a transformed regression using the natural log of both x and y would be appropriate. The probabilistic model is then  $y = \alpha x^{\beta} \cdot \varepsilon$ , the power function with an error term.
- **d.** A regression of  $\ln(y)$  on  $\ln(x)$  yields the equation  $\ln(y) = 4.6384 1.04920 \ln(x)$ . Using Minitab we can get a PI for y when x = 20 by first transforming the x value:  $\ln(20) = 2.996$ . The computer generated 95% PI for  $\ln(y)$  when  $\ln(x) = 2.996$  is (1.1188, 1.8712). We must now take the antilog to return to the original units of y:  $(e^{1.1188}, e^{1.8712}) = (3.06, 6.50)$ .
- e. A computer generated residual analysis:



Looking at the residual vs. fits (bottom right), one standardized residual, corresponding to the third observation, is a bit large. There are only two positive standardized residuals, but two others are essentially 0. The patterns in the residual plot and the normal probability plot (upper left) are marginally acceptable.

- **a.**  $\Sigma x'_i = 15.501$ ,  $\Sigma y'_i = 13.352$ ,  $\Sigma {x'_i}^2 = 20.228$ ,  $\Sigma {y'_i}^2 = 16.572$ ,  $\Sigma {x'_i} y'_i = 18.109$ , from which  $\hat{\beta}_1 = 1.254$  and  $\hat{\beta}_0 = -.468$  so  $\hat{\beta} = \hat{\beta}_1 = 1.254$  and  $\hat{\alpha} = e^{-.468} = .626$ .
- **b.** The plots give strong support to this choice of model; in addition,  $r^2 = .960$  for the transformed data.
- c. SSE = .11536 (computer printout), s = .1024, and the estimated sd of  $\hat{\beta}_1$  is .0775, so  $t = \frac{1.25 - \frac{4}{3}}{.0775} = -1.02$ , for a *P*-value at 11 df of  $P(T \le -1.02) \approx .16$ . Since .16 > .05,  $H_0$  cannot be rejected in favor of  $H_a$ .
- **d.** The claim that  $\mu_{\gamma,5} = 2\mu_{\gamma,2,5}$  is equivalent to  $\alpha \cdot 5^{\beta} = 2\alpha(2.5)^{\beta}$ , or that  $\beta = 1$ . Thus we wish test  $H_0: \beta = 1$  versus  $H_a: \beta \neq 1$ . With  $t = \frac{1.25 1}{.0775} = 3.28$ , the 2-sided *P*-value at 11 df is roughly 2(.004) = .008. Since  $.008 \le .01$ ,  $H_0$  is rejected at level .01.

19.

a. No, there is definite curvature in the plot.

- **b.** With x = temperature and y = lifetime, a linear relationship between ln(lifetime) and 1/temperature implies a model  $y = \exp(\alpha + \beta/x + \varepsilon)$ . Let x' = 1/temperature and y' = ln(lifetime). Plotting y' vs. x' gives a plot which has a pronounced linear appearance (and, in fact,  $r^2 = .954$  for the straight line fit).
- **c.**  $\Sigma x'_i = .082273$ ,  $\Sigma y'_i = 123.64$ ,  $\Sigma {x'_i}^2 = .00037813$ ,  $\Sigma {y'_i}^2 = 879.88$ ,  $\Sigma {x'_i} {y'_i} = .57295$ , from which  $\hat{\beta} = 3735.4485$  and  $\hat{\alpha} = -10.2045$  (values read from computer output). With x = 220, x' = .004545 so  $\hat{y}' = -10.2045 + 3735.4485(.004545) = 6.7748$  and thus  $\hat{y} = e^{\hat{y}} = 875.50$ .
- **d.** For the transformed data, SSE = 1.39857, and  $n_1 = n_2 = n_3 = 6$ ,  $\overline{y}'_{1.} = 8.44695$ ,  $\overline{y}'_{2.} = 6.83157$ ,  $\overline{y}'_{3.} = 5.32891$ , from which SSPE = 1.36594, SSLF = .02993,  $f = \frac{.02993/1}{1.36594/15} = .33$ . Comparing this to the *F* distribution with df = (1, 15), it is clear that  $H_0$  cannot be rejected.

Lordane of the resource variable and rightly a security of a stiller concernments with every concernment is the large of base are only the many depicted and the stiller concernments whe under as many of the proceeding december do not the second depicted of a stiller is an impacted premomently accurated.

#### 21.

ww.konkur.in

- a. The accompanying scatterplot, left, shows a very strong <u>non</u>-linear association between the variables. The corresponding residual plot would look somewhat like a downward-facing parabola.
- **b.** The right scatterplot shows y versus 1/x and exhibits a much more linear pattern. We'd anticipate an  $r^2$  value very near 1 based on the plot. (In fact,  $r^2 = .998$ .)



- c. With the aid of software, a 95% PI for y when x = 100, aka x' = 1/x = 1/100 = .01, can be generated. Using Minitab, the 95% PI is (83.89, 87.33). That is, at a 95% prediction level, the nitrogen extraction percentage for a single run when leaching time equals 100 h is between 83.89 and 87.33.
- 23.  $V(Y) = V(\alpha e^{\beta x} \cdot \varepsilon) = [\alpha e^{\beta x}]^2 \cdot V(\varepsilon) = \alpha^2 e^{2\beta x} \cdot \tau^2$  where we have set  $V(\varepsilon) = \tau^2$ . If  $\beta > 0$ , this is an increasing function of x so we expect more spread in y for large x than for small x, while the situation is reversed if  $\beta < 0$ . It is important to realize that a scatter plot of data generated from this model will not spread out uniformly about the exponential regression function throughout the range of x values; the spread will only be uniform on the transformed scale. Similar results hold for the multiplicative power model.
- 25. First, the test statistic for the hypotheses  $H_0: \beta_1 = 0$  versus  $H_a: \beta_1 \neq 0$  is z = -4.58 with a corresponding *P*-value of .000, suggesting noise level has a highly statistically significant relationship with people's perception of the acceptability of the work environment. The negative value indicates that the likelihood of finding work environment acceptable <u>decreases</u> as the noise level increases (not surprisingly). We estimate that a 1 dBA increase in noise level decreases the <u>odds</u> of finding the work environment acceptable by a multiplicative factor of .70 (95% CI: .60 to .81).

The accompanying plot shows  $\hat{\pi} = \frac{e^{b_0 + b_1 x}}{1 + e^{b_0 + b_1 x}} = \frac{e^{23.2 - .359 x}}{1 + e^{23.2 - .359 x}}$ . Notice that the estimate probability of finding

work environment acceptable decreases as noise level, x, increases.



181

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

## Section 13.3

a.

27.

www.konkur.in

A scatter plot of the data indicated a quadratic regression model might be appropriate.



**b.**  $\hat{y} = 84.482 - 15.875(6) + 1.7679(6)^2 = 52.88$ ; residual =  $y_6 - \hat{y}_6 = 53 - 52.88 = .12$ 

c. 
$$SST = \Sigma y_i^2 - \frac{(\Sigma y_i)^2}{n} = 586.88$$
, so  $R^2 = 1 - \frac{61.77}{586.88} = .895$ .

d. None of the standardized residuals exceeds 2 in magnitude, suggesting none of the observations are outliers. The ordered z percentiles needed for the normal probability plot are -1.53, -.89, -.49, -.16, .16, .49, .89, and 1.53. The normal probability plot below does not exhibit any troublesome features.



- e.  $\hat{\mu}_{Y-6} = 52.88$  (from b) and  $t_{.025,n-3} = t_{.025,5} = 2.571$ , so the CI is  $52.88 \pm (2.571)(1.69) = 52.88 \pm 4.34 = (48.54,57.22)$ .
- f. SSE = 61.77, so  $s^2 = \frac{61.77}{5} = 12.35$  and  $s \{ \text{pred} \} \sqrt{12.35 + (1.69)^2} = 3.90$ . The PI is  $52.88 \pm (2.571)(3.90) = 52.88 \pm 10.03 = (42.85, 62.91)$ .

29.

a. The table below displays the y-values, fits, and residuals. From this,  $SSE = \sum e^2 = 16.8$ ,  $s^2 = SSE/(n-3) = 4.2$ , and s = 2.05.

y	ŷ	$e = y - \hat{y}$
81	82.1342	-1.13420
83	80.7771	2.22292
79	79.8502	-0.85022
75	72.8583	2.14174
70	72.1567	-2.15670
43	43.6398	-0.63985
22	21.5837	0.41630

- **b.** SST =  $\sum (y \overline{y})^2 = \sum (y 64.71)^2 = 3233.4$ , so  $R^2 = 1 \text{SSE/SST} = 1 16.8/3233.4 = .995$ , or 99.5%. 995% of the variation in free-flow can be explained by the quadratic regression relationship with viscosity.
- c. We want to test the hypotheses  $H_0$ :  $\beta_2 = 0$  v.  $H_a$ :  $\beta_2 \neq 0$ . Assuming all inference assumptions are met, the relevant *t* statistic is  $t = \frac{-.0031662 - 0}{.0004835} = -6.55$ . At n - 3 = 4 df, the corresponding *P*-value is 2P(T > 6.55) < .004. At any reasonable significance level, we would reject  $H_0$  and conclude that the quadratic predictor indeed belongs in the regression model.
- **d.** Two intervals with at least 95% simultaneous confidence requires individual confidence equal to 100% 5%/2 = 97.5%. To use the *t*-table, round up to 98%:  $t_{.01,4} = 3.747$ . The two confidence intervals are  $2.1885 \pm 3.747(.4050) = (.671, 3.706)$  for  $\beta_1$  and  $-.0031662 \pm 3.747(.0004835) = (-.00498, -.00135)$  for  $\beta_2$ . [In fact, we are at least 96% confident  $\beta_1$  and  $\beta_2$  lie in these intervals.]
- e. Plug into the regression equation to get  $\hat{y} = 72.858$ . Then a 95% CI for  $\mu_{Y,400}$  is  $72.858 \pm 3.747(1.198)$ = (69.531, 76.186). For the PI,  $s\{\text{pred}\} = \sqrt{s^2 + s_{\hat{Y}}^2} = \sqrt{4.2 + (1.198)^2} = 2.374$ , so a 95% PI for Y when x = 400 is  $72.858 \pm 3.747(2.374) = (66.271, 79.446)$ .

31.

- a.  $R^2 = 98.0\%$  or .980. This means 98.0% of the observed variation in energy output can be attributed to the model relationship.
- **b.** For a quadratic model, adjusted  $R^2 = \frac{(n-1)R^2 k}{n-1-k} = \frac{(24-1)(.780) 2}{24-1-2} = .759$ , or 75.9%. (A more precise answer, from software, is 75.95%.) The adjusted  $R^2$  value for the <u>cubic</u> model is 97.7%, as seen in the output. This suggests that the cubic term greatly improves the model: the cost of adding an extra
  - in the output. This suggests that the cubic term greatly improves the model: the cost of adding an extra parameter is more than compensated for by the improved fit. H = 0 warraw H = 0. From the Minitab
- c. To test the utility of the cubic term, the hypotheses are  $H_0$ :  $\beta_3 = 0$  versus  $H_0$ :  $\beta_3 \neq 0$ . From the Minitab output, the test statistic is t = 14.18 with a *P*-value of .000. We strongly reject  $H_0$  and conclude that the cubic term is a statistically significant predictor of energy output, even in the presence of the lower terms.
- **d.** Plug x = 30 into the cubic estimated model equation to get  $\hat{y} = 6.44$ . From software, a 95% CI for  $\mu_{Y30}$  is (6.31, 6.57). Alternatively,  $\hat{y} \pm t_{.025,20} s_{\hat{y}} = 6.44 \pm 2.086(.0611)$  also gives (6.31, 6.57). Next, a 95% PI

for Y-30 is (6.06, 6.81) from software. Or, using the information provided,  $\hat{y} \pm t_{.025,20} \sqrt{s^2 + s_v^2}$  $6.44 \pm 2.086 \sqrt{(.1684)^2 + (.0611)^2}$  also gives (6.06, 6.81). The value of s comes from the Minitab output, where s = .168354.

The null hypothesis states that the true mean energy output when the temperature difference is 35°K is e. equal to 5W; the alternative hypothesis says this isn't true.

Plug x = 35 into the cubic regression equation to get  $\hat{y} = 4.709$ . Then the test statistic is

 $t = \frac{4.709-5}{.0523} \approx -5.6$ , and the two-tailed *P*-value at df = 20 is approximately 2(.000) = .000. Hence, we

strongly reject  $H_0$  (in particular, .000 < .05) and conclude that  $\mu_{Y35} \neq 5$ .

Alternatively, software or direct calculation provides a 95% CI for µy35 of (4.60, 4.82). Since this CI does not include 5, we can reject  $H_0$  at the .05 level.



- **a.**  $\overline{x} = 20$  and  $s_x = 10.8012$  so  $x' = \frac{x 20}{10.8012}$ . For x = 20, x' = 0, and  $\hat{y} = \hat{\beta}_0^* = .9671$ . For x = 25, x' = 0.4629, so  $\hat{y} = .9671 - .0502(.4629) - .0176(.4629)^2 + .0062(.4629)^3 = .9407$ .
- **b.**  $\hat{y} = .9671 .0502 \left( \frac{x 20}{10.8012} \right) .0176 \left( \frac{x 20}{10.8012} \right)^2 + .0062 \left( \frac{x 20}{10.8012} \right)^3$

 $= .00000492x^3 - .000446058x^2 + .007290688x + .96034944.$ 

c.  $t = \frac{.0062}{.0031} = 2.00$ . At df = n - 4 = 3, the *P*-value is 2(.070) = .140 > .05. Therefore, we cannot reject  $H_0$ ; the cubic term should be deleted.

- **d.** SSE =  $\Sigma (y_i \hat{y}_i)^2$  and the  $\hat{y}_i$ 's are the same from the standardized as from the unstandardized model, so SSE, SST, and  $R^2$  will be identical for the two models.
- e.  $\Sigma y_i^2 = 6.355538$ ,  $\Sigma y_i = 6.664$ , so SST = .011410. For the quadratic model,  $R^2 = .987$ , and for the cubic model,  $R^2 = .994$ . The two  $R^2$  values are very close, suggesting intuitively that the cubic term is relatively unimportant.

35.  $Y' = \ln(Y) = \ln \alpha + \beta x + \gamma x^2 + \ln(\varepsilon) = \beta_0 + \beta_1 x + \beta_2 x^2 + \varepsilon' \text{ where } \varepsilon' = \ln(\varepsilon), \ \beta_0 = \ln(\alpha), \ \beta_1 = \beta \text{ , and}$  $\beta_2 = \gamma$ . That is, we should fit a quadratic to  $(x, \ln(y))$ . The resulting estimated quadratic (from computer output) is  $2.00397 + .1799x - .0022x^2$ , so  $\hat{\beta} = .1799$ ,  $\hat{\gamma} = -.0022$ , and  $\hat{\alpha} = e^{2.0397} = 7.6883$ . [The ln(y)'s are 3.6136, 4.2499, 4.6977, 5.1773, and 5.4189, and the summary quantities can then be computed as before.]

www.konkur.in

### Section 13.4

#### 37.

- **a.** The mean value of y when  $x_1 = 50$  and  $x_2 = 3$  is  $\mu_{y-50,3} = -.800 + .060(50) + .900(3) = 4.9$  hours.
- **b.** When the number of deliveries  $(x_2)$  is held fixed, then average change in travel time associated with a one-mile (i.e., one unit) increase in distance traveled  $(x_1)$  is .060 hours. Similarly, when distance traveled  $(x_1)$  is held fixed, then the average change in travel time associated with on extra delivery (i.e., a one unit increase in  $x_2$ ) is .900 hours.
- c. Under the assumption that Y follows a normal distribution, the mean and standard deviation of this distribution are 4.9 (because  $x_1 = 50$  and  $x_2 = 3$ ) and  $\sigma = .5$  (since the standard deviation is assumed to be constant regardless of the values of  $x_1$  and  $x_2$ ). Therefore,

 $P(Y \le 6) = P\left(Z \le \frac{6-4.9}{.5}\right) = P(Z \le 2.20) = .9861$ . That is, in the long run, about 98.6% of all days

will result in a travel time of at most 6 hours.

#### 39.

- **a.** For  $x_1 = 2$ ,  $x_2 = 8$  (remember the units of  $x_2$  are in 1000s), and  $x_3 = 1$  (since the outlet has a drive-up window), the average sales are  $\hat{y} = 10.00 1.2(2) + 6.8(8) + 15.3(1) = 77.3$  (i.e., \$77,300).
- **b.** For  $x_1 = 3$ ,  $x_2 = 5$ , and  $x_3 = 0$  the average sales are  $\hat{y} = 10.00 1.2(3) + 6.8(5) + 15.3(0) = 40.4$  (i.e., \$40,400).
- c. When the number of competing outlets  $(x_1)$  and the number of people within a 1-mile radius  $(x_2)$  remain fixed, the expected sales will increase by \$15,300 when an outlet has a drive-up window.

#### 41.

- a.  $R^2 = .834$  means that 83.4% of the total variation in cone cell packing density (y) can be explained by a linear regression on eccentricity (x<sub>1</sub>) and axial length (x<sub>2</sub>). For  $H_0$ :  $\beta_1 = \beta_2 = 0$  vs.  $H_a$ : at least one  $\beta \neq 0$ , the test statistic is  $F = \frac{R^2 / k}{(1 R^2) / (n k 1)} = \frac{.834 / 2}{(1 .834) / (192 2 1)} \approx 475$ , and the associated *P*-value at df = (2, 189) is essentially 0. Hence,  $H_0$  is rejected and the model is judged useful.
- **b.**  $\hat{y} = 35821.792 6294.729(1) 348.037(25) = 20,826.138 \text{ cells/mm}^2$ .
- c. For a fixed axial length  $(x_2)$ , a 1-mm increase in eccentricity is associated with an estimated <u>decrease</u> in mean/predicted cell density of 6294.729 cells/mm<sup>2</sup>.
- **d.** The error df = n k 1 = 192 3 = 189, so the critical CI value is  $t_{.025,189} \approx z_{.025} = 1.96$ . A 95% CI for  $\beta_1$  is  $-6294.729 \pm 1.96(203.702) = (-6694.020, -5895.438)$ .
- e. The test statistic is  $t = \frac{-348.037 0}{134.350} = -2.59$ ; at 189 df, the 2-tailed *P*-value is roughly  $2P(T \le -2.59) \approx 2\Phi(-2.59) = 2(.0048) \approx .01$ . Since .01 < .05, we reject  $H_0$ . After adjusting for the effect of eccentricity  $(x_1)$ , there is a statistically significant relationship between axial length  $(x_2)$  and cell density (y). Therefore, we should retain  $x_2$  in the model.

www.konkur.in

 $\hat{y} = 185.49 - 45.97(2.6) - 0.3015(250) + 0.0888(2.6)(250) = 48.313$ .

- **b.** No, it is not legitimate to interpret  $\beta_1$  in this way. It is not possible to increase the cobalt content,  $x_1$ , while keeping the interaction predictor,  $x_3$ , fixed. When  $x_1$  changes, so does  $x_3$ , since  $x_3 = x_1x_2$ .
- c. Yes, there appears to be a useful linear relationship between y and the predictors. We determine this by observing that the *P*-value corresponding to the model utility test is < .0001 (*F* test statistic = 18.924).
- **d.** We wish to test  $H_0: \beta_3 = 0$  vs.  $H_a: \beta_3 \neq 0$ . The test statistic is t = 3.496, with a corresponding *P*-value of .0030. Since the *P*-value is  $< \alpha = .01$ , we reject  $H_0$  and conclude that the interaction predictor does provide useful information about *y*.
- e. A 95% CI for the mean value of surface area under the stated circumstances requires the following quantities:  $\hat{y} = 185.49 45.97(2) 0.3015(500) + 0.0888(2)(500) = 31.598$ . Next,  $t_{.025,16} = 2.120$ , so the 95% confidence interval is  $31.598 \pm (2.120)(4.69) = 31.598 \pm 9.9428 = (21.6552, 41.5408)$ .

45.

a. The hypotheses are  $H_0$ :  $\beta_1 = \beta_2 = \beta_3 = \beta_4 = 0$  vs.  $H_a$ : at least one  $\beta_i \neq 0$ . The test statistic is  $f = \frac{R^2 / k}{(1 - R^2)/(n - k - 1)} = \frac{.946 / 4}{(1 - .946) / 20} = 87.6 \ge F_{.001,4,20} = 7.10$  (the smallest available *F*-value from

Table A.9), so the *P*-value is < .001 and we can reject  $H_0$  at any significance level. We conclude that at least one of the four predictor variables appears to provide useful information about tenacity.

- **b.** The adjusted  $R^2$  value is  $1 \frac{n-1}{n-(k+1)} \left(\frac{SSE}{SST}\right) = 1 \frac{n-1}{n-(k+1)} (1-R^2) = 1 \frac{24}{20} (1-.946) = .935$ , which does not differ much from  $R^2 = .946$ .
  - . The estimated average tenacity when  $x_1 = 16.5$ ,  $x_2 = 50$ ,  $x_3 = 3$ , and  $x_4 = 5$  is  $\hat{y} = 6.121 - .082(16.5) + .113(50) + .256(3) - .219(5) = 10.091$ . For a 99% CI,  $t_{.005,20} = 2.845$ , so the interval is  $10.091 \pm 2.845(.350) = (9.095, 11.087)$ . Therefore, when the four predictors are as specified in this problem, the true average tenacity is estimated to be between 9.095 and 11.087.

#### 47.

- a. For a 1% increase in the percentage plastics, we would expect a 28.9 kcal/kg increase in energy content. Also, for a 1% increase in the moisture, we would expect a 37.4 kcal/kg decrease in energy content. Both of these assume we have accounted for the linear effects of the other three variables.
- **b.** The appropriate hypotheses are  $H_0$ :  $\beta_1 = \beta_2 = \beta_3 = \beta_4 = 0$  vs.  $H_a$ : at least one  $\beta \neq 0$ . The value of the *F*-test statistic is 167.71, with a corresponding *P*-value that is  $\approx 0$ . So, we reject  $H_0$  and conclude that at least one of the four predictors is useful in predicting energy content, using a linear model.
- c.  $H_0: \beta_3 = 0$  v.  $H_a: \beta_3 \neq 0$ . The value of the t test statistic is t = 2.24, with a corresponding *P*-value of .034, which is less than the significance level of .05. So we can reject  $H_0$  and conclude that percentage garbage provides useful information about energy consumption, given that the other three predictors remain in the model.

- **d.**  $\hat{y} = 2244.9 + 28.925(20) + 7.644(25) + 4.297(40) 37.354(45) = 1505.5$ , and  $t_{.025,25} = 2.060$ . So a 95% CI for the true average energy content under these circumstances is  $1505.5 \pm (2.060)(12.47) = 1505.5 \pm 25.69 = (1479.8,1531.1)$ . Because the interval is reasonably narrow, we would conclude that the mean energy content has been precisely estimated.
- e. A 95% prediction interval for the energy content of a waste sample having the specified characteristics is  $1505.5 \pm (2.060)\sqrt{(31.48)^2 + (12.47)^2} = 1505.5 \pm 69.75 = (1435.7,1575.2).$

49.

- **a.** Use the ANOVA table in the output to test  $H_0$ :  $\beta_1 = \beta_2 = \beta_3 = 0$  vs.  $H_a$ : at least one  $\beta_j \neq 0$ . With f = 17.31 and *P*-value = 0.000, so we reject  $H_0$  at any reasonable significance level and conclude that the model is useful.
- **b.** Use the *t* test information associated with  $x_3$  to test  $H_0$ :  $\beta_3 = 0$  vs.  $H_a$ :  $\beta_3 \neq 0$ . With t = 3.96 and *P*-value = .002 < .05, we reject  $H_0$  at the .05 level and conclude that the interaction term should be retained.
- c. The predicted value of y when  $x_1 = 3$  and  $x_2 = 6$  is  $\hat{y} = 17.279 6.368(3) 3.658(6) + 1.7067(3)(6) = 6.946$ . With error df = 11,  $t_{.025,11} = 2.201$ , and the CI is  $6.946 \pm 2.201(.555) = (5.73, 8.17)$ .

**d.** Our point prediction remains the same, but the SE is now  $\sqrt{s^2 + s_{\hat{\gamma}}^2} = \sqrt{1.72225^2 + .555^2} = 1.809$ . The resulting 95% PI is 6.946 ± 2.201(1.809) = (2.97, 10.93).

- 51.
- **a.** Associated with  $x_3$  = drilling depth are the test statistic t = 0.30 and *P*-value = .777, so we certainly do not reject  $H_0$ :  $\beta_3 = 0$  at any reasonably significance level. Thus, we should remove  $x_3$  from the model.
- **b.** To test  $H_0$ :  $\beta_1 = \beta_2 = 0$  vs.  $H_a$ : at least one  $\beta \neq 0$ , use  $R^2$ :  $f = \frac{R^2 / k}{(1 R^2) / (n k 1)} = \frac{.836 / 2}{(1 .836) / (9 2 1)}$ = 15.29; at df = (2, 6), 10.92 < 15.29 < 27.00  $\Rightarrow$  the *P*-value is between .001 and .01. (Software gives .004.) In particular, *P*-value  $\leq .05 \Rightarrow$  reject  $H_0$  at the  $\alpha = .05$  level: the model based on  $x_1$  and  $x_2$  is

useful in predicting y.

- c. With error df = 6,  $t_{.025,6}$  = 2.447, and from the Minitab output we can construct a 95% CI for  $\beta_1$ : -0.006767 ± 2.447(0.002055) = (-0.01180, -0.00174). Hence, after adjusting for feed rate ( $x_2$ ), we are 95% confident that the true change in mean surface roughness associated with a 1rpm increase in spindle speed is between -.01180 µm and -.00174 µm.
- **d.** The point estimate is  $\hat{y} = 0.365 0.006767(400) + 45.67(.125) = 3.367$ . With the standard error provided, the 95% CI for  $\mu_Y$  is  $3.367 \pm 2.447(.180) = (2.93, 3.81)$ .
- e. A normal probability plot of the  $e^*$  values is quite straight, supporting the assumption of normally distributed errors. Also, plots of the  $e^*$  values against  $x_1$  and  $x_2$  show no discernible pattern, supporting the assumptions of linearity and equal variance. Together, these validate the regression model.

- Some possible questions might be: 53.
  - Is this model useful in predicting deposition of poly-aromatic hydrocarbons? A test of model (1)utility gives us an F = 84.39, with a *P*-value of 0.000. Thus, the model is useful.
  - Is  $x_1$  a significant predictor of y in the presence of  $x_2$ ? A test of  $H_0$ :  $\beta_1 = 0$  v.  $H_a$ :  $\beta_1 \neq 0$  gives us a t (2)= 6.98 with a P-value of 0.000, so this predictor is significant.
  - A similar question, and solution for testing x2 as a predictor yields a similar conclusion: with a P-(3)value of 0.046, we would accept this predictor as significant if our significance level were anything larger than 0.046.

### Section 13.5

a.

Ĵ

55.

To test  $H_0$ :  $\beta_1 = \beta_2 = \beta_3 = 0$  vs.  $H_a$ : at least one  $\beta \neq 0$ , use  $R^2$ :

$$=\frac{R^2/k}{(1-R^2)/(n-k-1)} = \frac{.706/3}{(1-.706)/(12-3-1)} = 6.40. \text{ At } df = (3,8), 4.06 < 6.40 < 7.59 \Rightarrow \text{the } P$$

value is between .05 and .01. In particular, P-value  $< .05 \Rightarrow$  reject  $H_0$  at the .05 level. We conclude that the given model is statistically useful for predicting tool productivity.

- No: the large *P*-value (.510) associated with  $\ln(x_3)$  implies that we should not reject  $H_0$ :  $\beta_3 = 0$ , and b. hence we need not retain  $\ln(x_3)$  in the model that already includes  $\ln(x_1)$ .
- Part of the Minitab output from regression ln(y) on  $ln(x_1)$  appears below. The estimated regression c. equation is  $\ln(y) = 3.55 + 0.844 \ln(x_1)$ . As for utility, t = 4.69 and P-value = .001 imply that we should reject  $H_0$ :  $\beta_1 = 0$  — the stated model is useful.

The regression equation is ln(y) = 3.55 + 0.844 ln(x1)Coef SE Coef T Predictor 266.06 0.000 Constant 3.55493 0.01336 4.69 0.001 0.1799 0.8439 ln(x1)

d. The residual plot shows pronounced curvature, rather than "random scatter." This suggests that the functional form of the relationship might not be correctly modeled — that is,  $\ln(y)$  might have a <u>non</u>linear relationship with  $ln(x_1)$ . [Obviously, one should investigate this further, rather than blindly continuing with the given model!]

P



e. First, for the model utility test of  $\ln(x_1)$  and  $\ln^2(x_1)$  as predictors, we again rely on  $\mathbb{R}^2$ :

 $f = \frac{R^2 / k}{(1 - R^2) / (n - k - 1)} = \frac{.819 / 2}{(1 - .819) / (12 - 2 - 1)} = 20.36.$  Since this is greater than  $F_{.001,2,9} = 16.39$ , the

*P*-value is < .001 and we strongly reject the null hypothesis of no model utility (i.e., the utility of this model is confirmed). Notice also the *P*-value associated with  $\ln^2(x_1)$  is .031, indicating that this "quadratic" term adds to the model.

Next, notice that when  $x_1 = 1$ ,  $\ln(x_1) = 0$  [and  $\ln^2(x_1) = 0^2 = 0$ ], so we're really looking at the information associated with the intercept. Using that plus the critical value  $t_{.025,9} = 2.262$ , a 95% PI for

the response,  $\ln(Y)$ , when  $x_1 = 1$  is  $3.5189 \pm 2.262 \sqrt{.0361358^2 + .0178^2} = (3.4277, 3.6099)$ . Lastly, to create a 95% PI for Y itself, exponentiate the endpoints: at the 95% prediction level, a new value of Y when  $x_1 = 1$  will fall in the interval  $(e^{3.4277}, e^{3.6099}) = (30.81, 36.97)$ .

_	-	
_		
_	1.1	
~		

www.konkur.in

k	$R^2$	$R_a^2$	$C_k = \frac{SSE_k}{s^2} + 2(k+1) - n$
1	.676	.647	138.2
2	.979	.975	2.7
3	.9819	.976	3.2
4	.9824		4
0005			

where  $s^2 = 5.9825$ 

**a.** Clearly the model with k = 2 is recommended on all counts.

**b.** No. Forward selection would let  $x_4$  enter first and would not delete it at the next stage.

#### 59.

a. The choice of a "best" model seems reasonably clear-cut. The model with 4 variables including all but the summerwood fiber variable would seem best.  $R^2$  is as large as any of the models, including the 5-variable model.  $R^2$  adjusted is at its maximum and CP is at its minimum. As a second choice, one might consider the model with k = 3 which excludes the summerwood fiber and springwood % variables.

#### b. Backwards Stepping:

Step 1: A model with all 5 variables is fit; the smallest *t*-ratio is t = .12, associated with variable  $x_2$  (summerwood fiber %). Since t = .12 < 2, the variable  $x_2$  was eliminated.

- Step 2: A model with all variables except  $x_2$  was fit. Variable  $x_4$  (springwood light absorption) has the smallest *t*-ratio (t = -1.76), whose magnitude is smaller than 2. Therefore,  $x_4$  is the next variable to be eliminated.
- Step 3: A model with variables  $x_3$  and  $x_5$  is fit. Both *t*-ratios have magnitudes that exceed 2, so both variables are kept and the backwards stepping procedure stops at this step. The final model identified by the backwards stepping method is the one containing  $x_3$  and  $x_5$ .

Forward Stepping:

- Step 1: After fitting all five 1-variable models, the model with  $x_3$  had the *t*-ratio with the largest magnitude (t = -4.82). Because the absolute value of this *t*-ratio exceeds 2,  $x_3$  was the first variable to enter the model.
- Step 2: All four 2-variable models that include  $x_3$  were fit. That is, the models  $\{x_3, x_1\}, \{x_3, x_2\}, \{x_3, x_4\}, \{x_3, x_5\}$  were all fit. Of all 4 models, the *t*-ratio 2.12 (for variable  $x_5$ ) was largest in absolute value. Because this *t*-ratio exceeds 2,  $x_5$  is the next variable to enter the model.
- Step 3: (not printed): All possible 3-variable models involving  $x_3$  and  $x_5$  and another predictor. None of the *t*-ratios for the added variables has absolute values that exceed 2, so no more variables are added. There is no need to print anything in this case, so the results of these tests are not shown.
- Note: Both the forwards and backwards stepping methods arrived at the same final model,  $\{x_3, x_5\}$ , in this problem. This often happens, but not always. There are cases when the different stepwise methods will arrive at slightly different collections of predictor variables.
- 61. If multicollinearity were present, at least one of the four  $R^2$  values would be very close to 1, which is not the case. Therefore, we conclude that multicollinearity is not a problem in this data.
- 63. Before removing any observations, we should investigate their source (e.g., were measurements on that observation misread?) and their impact on the regression. To begin, Observation #7 deviates significantly from the pattern of the rest of the data (standardized residual = -2.62); if there's concern the PAH deposition was not measured properly, we might consider removing that point to improve the overall fit. If the observation was <u>not</u> mis-recorded, we should <u>not</u> remove the point.

We should also investigate Observation #6: Minitab gives  $h_{66} = .846 > 3(2+1)/17$ , indicating this observation has very high leverage. However, the standardized residual for #6 is not large, suggesting that it follows the regression pattern specified by the other observations. Its "influence" only comes from having a comparatively large  $x_1$  value.

## **Supplementary Exercises**

65.

a.

nkur.in

WWWWW.

Boxplots of ppv by prism quality (means are indicated by solid circles)

A two-sample t confidence interval, generated by Minitab: Two sample T for ppv

nriem mi	N	Mean	StDev	SE Mean	
prioked	12	827	295	85	
not cracke	18	483	234	55	
not cracke	10	10.56			

95% CI for mu (cracked ) - mu (not cracke): ( 132, 557)

**b.** The simple linear regression results in a significant model,  $r^2$  is .577, but we have an extreme observation, with std resid = -4.11. Minitab output is below. Also run, but not included here was a model with an indicator for cracked/ not cracked, and for a model with the indicator and an interaction term. Neither improved the fit significantly.

The regression equation is ratio = 1.00 -0.000018 ppv

 Predictor
 Coef
 StDev
 T
 P

 Constant
 1.00161
 0.00204
 491.18
 0.000

 ppv
 -0.00001827
 0.00000295
 -6.19
 0.000

S = 0.004892 R-Sq = 57.7% R-Sq(adj) = 56.2%

Analysis of Variance

Source Regression Residual Er Total	DF 1 ror 28 29	0.0009	SS 91571 57016 58587	0.000	MS 91571 02393	F 38.26	0.000		
Unusual Obs Obs 29 1	ervations ppv 1 144 0.9	ratio 62000	0.980	Fit 704	StDev F 0.0017	it Res 86 -0.0	idual 18704	St	Resid -4.11R

R denotes an observation with a large standardized residual

67.

ww.konkur.in

- a. After accounting for all the other variables in the regression, we would expect the VO<sub>2</sub>max to decrease by .0996, on average for each one-minute increase in the one-mile walk time.
- b. After accounting for all the other variables in the regression, we expect males to have a VO<sub>2</sub>max that is .6566 L/min higher than females, on average.
- c.  $\hat{y} = 3.5959 + .6566(1) + .0096(170) .0996(11) .0880(140) = 3.67$ . The residual is  $\hat{y} = (3.15 3.67) = -.52$ .
- d.  $R^2 = 1 \frac{SSE}{SST} = 1 \frac{30.1033}{102.3922} = .706$ , or 70.6% of the observed variation in VO<sub>2</sub>max can be attributed to the model relationship.
- e. To test  $H_0$ :  $\beta_1 = \beta_2 = \beta_3 = \beta_4 = 0$  vs.  $H_a$ : at least one  $\beta \neq 0$ , use  $R^2$ :

 $f = \frac{R^2/k}{(1-R^2)/(n-k-1)} = \frac{.706/4}{(1-.706)/(20-4-1)} = 9.005.$  At df = (4, 15), 9.005 > 8.25  $\Rightarrow$  the *P*-value is less than .05, so  $H_0$  is rejected. It appears that the model specifies a useful relationship between VO<sub>2</sub>max and at least one of the other predictors.

69.

a. Based on a scatter plot (below), a simple linear regression model would not be appropriate. Because of the slight, but obvious curvature, a quadratic model would probably be more appropriate.



 b. Using a quadratic model, a Minitab generated regression equation is ŷ = 35.423+1.7191x-.0024753x<sup>2</sup>, and a point estimate of temperature when pressure is 200 is ŷ = 280.23. Minitab will also generate a 95% prediction interval of (256.25, 304.22). That is, we are confident that when pressure is 200 psi, a single value of temperature will be between 256.25 and 304.22°F.

71.

www.konkur.in

- a. Using Minitab to generate the first order regression model, we test the model utility (to see if any of the predictors are useful), and with f = 21.03 and a *P*-value of .000, we determine that at least one of the predictors is useful in predicting palladium content. Looking at the individual predictors, the *P*-value associated with the pH predictor has value .169, which would indicate that this predictor is unimportant in the presence of the others.
- **b.** We wish to test  $H_0: \beta_1 = ... = \beta_{20} = 0$  vs.  $H_a:$  at least one  $\beta \neq 0$ . With calculated statistic f = 6.29 and *P*-value .002, this model is also useful at any reasonable significance level.
- c. Testing  $H_0: \beta_6 = ... = \beta_{20} = 0$  vs.  $H_a$ : at least one of the listed  $\beta$ 's  $\neq 0$ , the test statistic is

$$f = \frac{(716.10 - 290.27)/(20 - 5)}{290.27(32 - 20 - 1)} = 1.07 < F_{.05,15,11} = 2.72.$$
 Thus, *P*-value > .05, so we fail to reject H<sub>0</sub>

and conclude that all the quadratic and interaction terms should not be included in the model. They do not add enough information to make this model significantly better than the simple first order model.

d. Partial output from Minitab follows, which shows all predictors as significant at level .05:

```
The regression equation is

pdconc = - 305 + 0.405 niconc + 69.3 pH - 0.161 temp + 0.993 currdens

+ 0.355 pallcont - 4.14 pHsq
```

Predictor	Coef	StDev	Т	P
Constant	-304.85	93.98	-3.24	0.003
niconc	0.40484	0.09432	4.29	0.000
oH	69.27	21.96	3.15	0.004
temp	-0.16134	0.07055	-2.29	0.031
currdens	0,9929	0.3570	2.78	0.010
oallcont	0.35460	0.03381	10.49	0.000
oHsa	-4.138	1.293	-3.20	0.004
a second				

73.

**a.** We wish to test  $H_0$ :  $\beta_1 = \beta_2 = 0$  vs.  $H_a$ : either  $\beta_1$  or  $\beta_2 \neq 0$ . With  $R^2 = 1 - \frac{.29}{202.88} = .9986$ , the test

statistic is  $f = \frac{R^2 / k}{(1 - R^2) / (n - k - 1)} = \frac{.9986 / 2}{(1 - .9986) / (8 - 2 - 1)} = 1783$ , where k = 2 for the quadratic model.

Clearly the *P*-value at df = (2,5) is effectively zero, so we strongly reject  $H_0$  and conclude that the quadratic model is clearly useful.

**b.** The relevant hypotheses are  $H_a$ :  $\beta_2 = 0$  vs.  $H_a$ :  $\beta_2 \neq 0$ . The test statistic value is

$$t = \frac{\hat{\beta}_2}{s_a} = \frac{-.00163141 - 0}{.00003391} = -48.1$$
; at 5 df, the *P*-value is  $2P(T \ge |-48.1|) \approx 0$ . Therefore,  $H_0$  is rejected.

The quadratic predictor should be retained.

- c. No.  $R^2$  is extremely high for the quadratic model, so the marginal benefit of including the cubic predictor would be essentially nil and a scatter plot doesn't show the type of curvature associated with a cubic model.
- **d.**  $t_{.025,5} = 2.571$ , and  $\hat{\beta}_0 + \hat{\beta}_1(100) + \hat{\beta}_2(100)^2 = 21.36$ , so the CI is  $21.36 \pm 2.571(.1141) = 21.36 \pm .29$ = (21.07,21.65).

First, we need to figure out  $s^2$  based on the information we have been given:  $s^2 = MSE = SSE/df =$ .29/5 = .058. Then, the 95% PI is  $21.36 \pm 2.571 \sqrt{.058 + (.1141)^2} = 21.36 \pm 0.685 = (20.675, 22.045)$ .

75.

77.

```
a. To test H_0: \beta_1 = \beta_2 = 0 vs. H_a: either \beta_1 or \beta_2 \neq 0, first find R^2: SST = \sum y^2 - (\sum y)^2 / n = 264.5 \implies R^2 = 264.5
      1 - \text{SSE/SST} = 1 - 26.98/264.5 = .898. Next, f = \frac{.898/2}{(1 - .898)/(10 - 2 - 1)} = 30.8, which at df = (2,7)
```

corresponds to a P-value of  $\approx 0$ . Thus,  $H_0$  is rejected at significance level .01 and the quadratic model is judged useful.

**b.** The hypotheses are  $H_0: \beta_2 = 0$  vs.  $H_a: \beta_2 \neq 0$ . The test statistic value is t = (-2.3621 - 0)/.3073 = -7.69, and at 7 df the P-value is  $2P(T \ge |-7.69|) \approx 0$ . So,  $H_0$  is rejected at level .001. The quadratic predictor should not be eliminated.

c. 
$$x = 1$$
 here,  $\hat{\mu}_{Y_{11}} = \hat{\beta}_0 + \hat{\beta}_1(1) + \hat{\beta}_2(1)^2 = 45.96$ , and  $t_{.025,7} = 1.895$ , giving the CI  
45.96 + (1.895)(1.031) = (44.01.47.91).

- a.
  - The hypotheses are  $H_0$ :  $\beta_1 = \beta_2 = \beta_3 = \beta_4 = 0$  versus  $H_a$ : at least one  $\beta_i \neq 0$ . From the output, the Fstatistic is f = 4.06 with a P-value of .029. Thus, at the .05 level we reject  $H_0$  and conclude that at least one of the explanatory variables is a significant predictor of power.
- **b.** Yes, a model with  $R^2 = .834$  would appear to be useful. A formal model utility test can be performed:

 $f = \frac{R^2 / k}{(1 - R^2) / [n - (k + 1)]} = \frac{.834 / 3}{(1 - .834) / [16 - 4]} = 20.1$ , which is much greater than  $F_{.05,3,12} = 3.49$ . Thus, the mode including  $\{x_3, x_4, x_3x_4\}$  is usef

We cannot use an F test to compare this model with the first-order model in (a), because neither model is a "subset" of the other. Compare  $\{x_1, x_2, x_3, x_4\}$  to  $\{x_3, x_4, x_3x_4\}$ .

The hypotheses are  $H_0: \beta_5 = ... = \beta_{10} = 0$  versus  $H_a$ : at least one of these  $\beta_i \neq 0$ , where  $\beta_5$  through  $\beta_{10}$  are c. the coefficients for the six interaction terms. The "partial F test" statistic is

 $f = \frac{(\text{SSE}_l - \text{SSE}_k)/(k-l)}{\text{SSE}_k/[n-(k+1)]} = \frac{(R_k^2 - R_l^2)/(k-l)}{(1 - R_k^2)/[n-(k+1)]} = \frac{(.960 - .596)/(10 - 4)}{(1 - .960)/[16 - (10 + 1)]} = 7.58$ , which is greater

than  $F_{.05,6,5} = 4.95$ . Hence, we reject  $H_0$  at the .05 level and conclude that at least one of the interaction terms is a statistically significant predictor of power, in the presence of the first-order terms.

There are obviously several reasonable choices in each case. In a, the model with 6 carriers is a defensible 79. choice on all three grounds, as are those with 7 and 8 carriers. The models with 7, 8, or 9 carriers in b merit serious consideration. These models merit consideration because  $R_k^2$ ,  $MSE_k$ , and  $C_k$  meet the variable selection criteria given in Section 13.5.

81.

The relevant hypotheses are  $H_0: \beta_1 = ... = \beta_5 = 0$  vs.  $H_a$ : at least one among  $\beta_1, ..., \beta_5 \neq 0$ . f =

 $\frac{.827/5}{.173/11} = 106.1 \ge F_{.05,5,111} \approx 2.29$ , so *P*-value < .05. Hence,  $H_0$  is rejected in favor of the conclusion that there is a useful linear relationship between Y and at least one of the predictors.

194

**b.**  $t_{.05,111} = 1.66$ , so the CI is  $.041 \pm (1.66)(.016) = .041 \pm .027 = (.014, .068)$ .  $\beta_1$  is the expected change in mortality rate associated with a one-unit increase in the particle reading when the other four predictors are held fixed; we can be 90% confident that  $.014 < \beta_1 < .068$ .

c. In testing  $H_0: \beta_4 = 0$  versus  $H_a: \beta_4 \neq 0$ ,  $t = \frac{\hat{\beta}_4 - 0}{s_{\hat{\beta}_4}} = \frac{.047}{.007} = 5.9$ , with an associated *P*-value of  $\approx 0$ . So,

 $H_0$  is rejected and this predictor is judged important.

- **d.**  $\hat{y} = 19.607 + .041(166) + .071(60) + .001(788) + .041(68) + .687(.95) = 99.514$ , and the corresponding residual is 103 99.514 = 3.486.
- Taking logs, the regression model is  $\ln(Y) = \beta_0 + \beta_1 \ln(x_1) + \beta_2 \ln(x_2) + \varepsilon'$ , where  $\beta_0 = \ln(\alpha)$ . Relevant Minitab output appears below.
  - **a.** From the output,  $\hat{\beta}_0 = 10.8764$ ,  $\hat{\beta}_1 = -1.2060$ ,  $\hat{\beta}_2 = -1.3988$ . In the original model, solving for  $\alpha$  returns  $\hat{\alpha} = \exp(\hat{\beta}_0) = e^{10.8764} = 52,912.77$ .
  - b. From the output, R<sup>2</sup> = 78.2%, so 78.2% of the total variation in ln(wear life) can be explained by a linear regression on ln(speed) and ln(load). From the ANOVA table, a test of H<sub>0</sub>: β<sub>1</sub> = β<sub>2</sub> = 0 versus H<sub>a</sub>: at least one of these β's ≠ 0 produces f = 42.95 and P-value = 0.000, so we strongly reject H<sub>0</sub> and conclude that the model is useful.
  - c. Yes: the variability utility *t*-tests for the two variables have t = -7.05, P = 0.000 and t = -6.01, P = 0.000. These indicate that each variable is highly statistically significant.
  - d. With ln(50) ≈ 3.912 and ln(5) ≈ 1.609 substituted for the transformed x values, Minitab produced the accompanying output. A 95% PI for ln(Y) at those settings is (2.652, 5.162). Solving for Y itself, the 95% PI of interest is (e<sup>2.652</sup>, e<sup>5.162</sup>) = (14.18, 174.51).

```
The regression equation is
ln(y) = 10.9 - 1.21 ln(x1) - 1.40 ln(x2)
Predictor Coef SE Coef T P
```

onstant	10.8764	0.7872	13.82	0.000	
n(v1)	-1.2060	0.1710	-7.05	0.000	
$\ln(x^2)$	-1.3988	0.2327	-6.01	0.000	
The Jacob V.					

S = 0.596553 R-Sq = 78.2% R-Sq(adj) = 76.3%

Analysis of Variance

```
        Source
        DF
        SS
        MS
        F
        P

        Regression
        2
        30.568
        15.284
        42.95
        0.000

        Residual Error
        24
        8.541
        0.356

        Total
        26
        39.109
```

Predicted Values for New Observations

New Obs Fit SE Fit 95% CI 95% PI 1 3.907 0.118 (3.663, 4.151) (2.652, 5.162)

83.

WWW.K

nkur.in

## **CHAPTER 14**

## Section 14.1

1.

For each part, we reject  $H_0$  if the *P*-value is  $\leq \alpha$ , which occurs if and only the calculated  $\chi^2$  value is greater than or equal to the value  $\chi^2_{\alpha,k-1}$  from Table A.7.

a. Since  $12.25 \ge \chi^2_{.05,4} = 9.488$ , *P*-value  $\le .05$  and we would reject  $H_0$ .

- **b.** Since  $8.54 < \chi^2_{.013} = 11.344$ , *P*-value > .01 and we would fail to reject  $H_0$ .
- c. Since  $4.36 < \chi^2_{10.2} = 4.605$ , P-value > .10 and we would fail to reject  $H_0$ .
- d. Since  $10.20 < \chi^2_{015} = 15.085$ , *P*-value > .01 we would fail to reject  $H_0$ .

3. The uniform hypothesis implies that  $p_{i0} = \frac{1}{8} = .125$  for i = 1, ..., 8, so the null hypothesis is  $H_0: p_{10} = p_{20} = ... = p_{80} = .125$ . Each expected count is  $np_{i0} = 120(.125) = 15$ , so

 $\chi^{2} = \left[\frac{\left(12-15\right)^{2}}{15} + \dots + \frac{\left(10-15\right)^{2}}{15}\right] = 4.80 \text{ . At } df = 8 - 1 = 7, 4.80 < 12.10 \Rightarrow P \text{-value} > .10 \Rightarrow \text{ we fail to}$ 

reject  $H_0$ . There is not enough evidence to disprove the claim.

The observed values, expected values, and corresponding  $\chi^2$  terms are :

Obs	4	15	23	25	38	21	32	14	10	8
Exp	6.67	13.33	20	26.67	33.33	33.33	26.67	20	13.33	6.67
$\chi^2$	1.069	.209	.450	.105	.654	.163	1.065	1.800	.832	.265

 $\chi^2 = 1.069 + ... + .265 = 6.612$ . With df = 10 - 1 = 9, 6.612 < 14.68  $\Rightarrow$  P-value > .10  $\Rightarrow$  we cannot reject  $H_0$ . There is no significant evidence that the data is not consistent with the previously determined proportions.

7.

5.

We test  $H_0: p_1 = p_2 = p_3 = p_4 = .25$  vs.  $H_a$ : at least one proportion  $\neq .25$ , and df = 3.

Cell	1	2	3	4
Observed	328	334	372	327
Expected	340.25	340.25	340.25	34.025
$\chi^2$ term	.4410	.1148	2.9627	.5160

 $\chi^2 = 4.0345$ , and with 3 df, *P*-value > .10, so we fail to reject  $H_0$ . The data fails to indicate a seasonal relationship with incidence of violent crime.

**a.** Denoting the 5 intervals by  $[0, c_1), [c_1, c_2), ..., [c_4, \infty)$ , we wish  $c_1$  for which

$$2 = P(0 \le X \le c_1) = \int_{-\infty}^{c_1} e^{-x} dx = 1 - e^{-c_1}$$
, so  $c_1 = -\ln(.8) = .2231$ . Then

 $.2 = P(c_1 \le X \le c_2) \Rightarrow .4 = P(0 \le X_1 \le c_2) = 1 - e^{-c_2}$ , so  $c_2 = -\ln(.6) = .5108$ . Similarly,  $c_3 = -\ln(.4) = .0163$  and  $c_4 = -\ln(.2) = 1.6094$ . The resulting intervals are [0, .2231), [.2231, .5108), [.5108, .9163), [.9163, 1.6094), and [1.6094,  $\infty$ ).

**b.** Each expected cell count is 40(.2) = 8, and the observed cell counts are 6, 8, 10, 7, and 9, so

$$\chi^{2} = \left[\frac{(6-8)^{2}}{8} + \dots + \frac{(9-8)^{2}}{8}\right] = 1.25. \text{ Because } 1.25 < \chi^{2}_{.10,4} = 7.779 \text{ , even at level .10 } H_{0} \text{ cannot be}$$

rejected; the data is quite consistent with the specified exponential distribution.

#### 11.

9.

unkur.in

WWW.K

a. The six intervals must be symmetric about 0, so denote the 4<sup>th</sup>, 5<sup>th</sup> and 6<sup>th</sup> intervals by [0, a), [a, b), [b, ∞). The constant a must be such that Φ(a) = .6667(<sup>1</sup>/<sub>2</sub> + <sup>1</sup>/<sub>6</sub>), which from Table A.3 gives a ≈ .43. Similarly, Φ(b) = .8333 implies b ≈ .97, so the six intervals are (-∞, -.97), [-.97, -.43), [-.43, 0), [0, .43), [.43, .97), and [.97, ∞).

- b. The six intervals are symmetric about the mean of .5. From a, the fourth interval should extend from the mean to .43 standard deviations above the mean, i.e., from .5 to .5 + .43(.002), which gives [.5, .50086). Thus the third interval is [.5 .00086, .5) = [.49914, .5). Similarly, the upper endpoint of the fifth interval is .5 + .97(.002) = .50194, and the lower endpoint of the second interval is .5 .00194 = .49806. The resulting intervals are (-∞, .49806), [.49806, .49914), [.49914, .5), [.5, .50086), [.50086, .50194), and [.50194, ∞).
- c. Each expected count is 45(1/6) = 7.5, and the observed counts are 13, 6, 6, 8, 7, and 5, so  $\chi^2 = 5.53$ . With 5 df, the *P*-value > .10, so we would fail to reject  $H_0$  at any of the usual levels of significance. There is no significant evidence to suggest that the bolt diameters are not normally distributed with  $\mu = .5$  and  $\sigma = .002$ .

### Section 14.2

13. According to the stated model, the three cell probabilities are  $(1-p)^2$ , 2p(1-p), and  $p^2$ , so we wish the value of p which maximizes  $(1-p)^{2n_i} [2p(1-p)]^{n_2} p^{2n_3}$ . Proceeding as in Example 14.6 gives

$$\hat{p} = \frac{n_2 + 2n_3}{2n} = \frac{234}{2776} = .0843 \text{ . The estimated expected cell counts are then } n(1-\hat{p})^2 = 1163.85 \text{ ,}$$

$$n[2\hat{p}(1-\hat{p})]^2 = 214.29 \text{ , } n\hat{p}^2 = 9.86 \text{ . This gives}$$

$$\chi^2 = \left[\frac{(1212 - 1163.85)^2}{1163.85} + \frac{(118 - 214.29)^2}{214.29} + \frac{(58 - 9.86)^2}{9.86}\right] = 280.3 \text{ . With } df = 4 - 1 - 1 = 2, 280.3 > 13.8$$

 $\Rightarrow$  *P*-value < .001  $\Rightarrow$  *H*<sub>0</sub> is soundly rejected. The stated model is strongly contradicted by the data.

15.

KUL.IL

The part of the likelihood involving  $\theta$  is  $[(1-\theta)^4]^{n_1} \cdot [\theta(1-\theta)^3]^{n_2} \cdot [\theta^2(1-\theta)^2]^{n_3} \cdot [\theta^3(1-\theta)]^{n_4} \cdot [\theta^4]^{n_5} = \theta^{n_2+2n_3+3n_4+4n_5}(1-\theta)^{4n_1+3n_2+2n_3+n_4} = \theta^{233}(1-\theta)^{367}$ , so the log-likelihood is 233 ln  $\theta$  + 367 ln(1- $\theta$ ). Differentiating and equating to 0 yields  $\hat{\theta} = \frac{233}{600} = .3883$ , and  $(1-\hat{\theta}) = .6117$  [note that the exponent on  $\theta$  is simply the total # of successes (defectives here) in the n = 4(150) = 600 trials]. Substituting this  $\hat{\theta}$  into the formula for  $p_i$  yields estimated cell probabilities .1400, .3555, .3385, .1433, and .0227. Multiplication by 150 yields the estimated expected cell counts are 21.00, 53.33, 50.78, 21.50, and 3.41. the last estimated expected cell count is less than 5, so we combine the last two categories into a single one ( $\geq$  3 defectives), yielding estimated counts 21.00, 53.33, 50.78, 24.91, observed counts 26, 51, 47, 26, and  $\chi^2 = 1.62$ . With df = 4 - 1 - 1 = 2, since  $1.62 < \chi^2_{.10,2} = 4.605$ , the *P*-value > .10, and we do not reject  $H_0$ . The data suggests that the stated binomial distribution is plausible.

17. 
$$\hat{\mu} = \overline{x} = \frac{(0)(6) + (1)(24) + (2)(42) + \dots + (8)(6) + (9)(2)}{300} = \frac{1163}{300} = 3.88$$
, so the estimated cell probabilities are computed from  $\hat{p} = e^{-3.88} \frac{(3.88)^x}{x!}$ .

x	0	1	2	3	4	5	6	7	≥8
np(x)	6.2	24.0	46.6	60.3	58.5	45.4	29.4	16.3	13.3
obs	6	24	42	59	62	44	41	14	8

This gives  $\chi^2 = 7.789$ . At df = 9 - 1 - 1 = 7, 7.789 < 12.01  $\Rightarrow$  *P*-value > .10  $\Rightarrow$  we fail to reject  $H_0$ . The Poisson model does provide a good fit.

19. With 
$$A = 2n_1 + n_4 + n_5$$
,  $B = 2n_2 + n_4 + n_6$ , and  $C = 2n_3 + n_5 + n_6$ , the likelihood is proportional to  $\theta_1^A \theta_2^B (1 - \theta_1 - \theta_2)^C$ . Taking the natural log and equating both  $\frac{\partial}{\partial \theta_1}$  and  $\frac{\partial}{\partial \theta_2}$  to zero gives  $\frac{A}{\theta_1} = \frac{C}{1 - \theta_1 - \theta_2}$  and  $\frac{B}{\theta_2} = \frac{C}{1 - \theta_1 - \theta_2}$ , whence  $\theta_2 = \frac{B\theta_1}{A}$ . Substituting this into the first equation gives  $\theta_1 = \frac{A}{A + B + C}$ , and then  $\theta_2 = \frac{B}{A + B + C}$ . Thus  $\hat{\theta}_1 = \frac{2n_1 + n_4 + n_5}{2n}$ ,  $\hat{\theta}_2 = \frac{2n_2 + n_4 + n_6}{2n}$ , and  $(1 - \hat{\theta}_1 - \hat{\theta}_2) = \frac{2n_3 + n_5 + n_6}{2n}$ . Substituting the observed  $n_i$ 's yields  $\hat{\theta}_1 = \frac{2(49) + 20 + 53}{400} = .4275$ ,  $\hat{\theta}_2 = \frac{110}{400} = .2750$ , and  $(1 - \hat{\theta}_1 - \hat{\theta}_2) = .2975$ , from which  $\hat{p}_1 = (.4275)^2 = .183$ ,  $\hat{p}_2 = .076$ ,  $\hat{p}_3 = .089$ ,  $\hat{p}_4 = 2(.4275)(.275) = .235$ ,  $\hat{p}_5 = .254$ ,  $\hat{p}_6 = .164$ .

Category	1	2	3	4	5	6
пр	36.6	15.2	17.8	47.0	50.8	32.8
observed	49	26	14	20	53	38

This gives  $\chi^2 = 29.1$ . At df = 6 - 1 - 2 = 3, this gives a *P*-value less than .001. Hence, we reject  $H_0$ .

- 21. The Ryan-Joiner test *P*-value is larger than .10, so we conclude that the null hypothesis of normality cannot be rejected. This data could reasonably have come from a normal population. This means that it would be legitimate to use a one-sample *t* test to test hypotheses about the true average ratio.
- 23. Minitab gives r = .967, though the hand calculated value may be slightly different because when there are ties among the  $x_{(i)}$ 's, Minitab uses the same  $y_i$  for each  $x_{(i)}$  in a group of tied values.  $c_{10} = .9707$ , and  $c_{.05} = .9639$ , so .05 < P-value < .10. At the 5% significance level, one would have to consider population normality plausible.

### Section 14.3

www.konkur.in

- 25. The hypotheses are  $H_0$ : there is <u>no</u> association between extent of binge drinking and age group vs.  $H_a$ : there <u>is</u> an association between extent of binge drinking and age group. With the aid of software, the calculated test statistic value is  $\chi^2 = 212.907$ . With all expected counts well above 5, we can compare this value to a chi-squared distribution with df = (4-1)(3-1) = 6. The resulting *P*-value is  $\approx 0$ , and so we strongly reject  $H_0$  at any reasonable level (including .01). There is strong evidence of an association between age and binge drinking for college-age males. In particular, comparing the observed and expected counts shows that <u>younger</u> men tend to binge drink <u>more</u> than expected if  $H_0$  were true.
- 27. With i = 1 identified with men and i = 2 identified with women, and j = 1, 2, 3 denoting the 3 categories L>R, L=R, L<R, we wish to test  $H_0$ :  $p_{1j} = p_{2j}$  for j = 1, 2, 3 vs.  $H_a$ :  $p_{1j} \neq p_{2j}$  for at least one j. The estimated cell counts for men are 17.95, 8.82, and 13.23 and for women are 39.05, 19.18, 28.77, resulting in a test statistic of  $\chi^2 = 44.98$ . With (2 1)(3 1) = 2 degrees of freedom, the *P*-value is < .001, which strongly suggests that  $H_0$  should be rejected.

#### 29.

a. The null hypothesis is  $H_0$ :  $p_{1j} = p_{2j} = p_{3j}$  for j = 1, 2, 3, 4, where  $p_{ij}$  is the proportion of the *i*th population (natural scientists, social scientists, non-academics with graduate degrees) whose degree of spirituality falls into the *j*th category (very, moderate, slightly, not at all).

From the accompanying Minitab output, the test statistic value is  $\chi^2 = 213.212$  with df = (3–1)(4–1) = 6, with an associated *P*-value of 0.000. Hence, we strongly reject  $H_0$ . These three populations are <u>not</u> homogeneous with respect to their degree of spirituality.

#### Chi-Square Test: Very, Moderate, Slightly, Not At All

Expected counts are printed below observed counts Chi-Square contributions are printed below expected counts

	Very	Moderate	Slightly	Not At All	Total
1	56	162	198	211	627
	78.60	195.25	183.16	170.00	
	6.497	5.662	1.203	9.889	
2	56	223	243	239	761
	95.39	236.98	222.30	206.33	
	16.269	0.824	1.928	5.173	
3	109	164	74	28	375
	47.01	116.78	109.54	101.67	
	81.752	19.098	11.533	53.384	
Total	221	549	515	478	1763

Chi-Sq = 213.212, DF = 6, P-Value = 0.000

**b.** We're now testing  $H_0: p_{1j} = p_{2j}$  for j = 1, 2, 3, 4 under the same notation. The accompanying Minitab output shows  $\chi^2 = 3.091$  with df = (2-1)(4-1) = 3 and an associated *P*-value of 0.378. Since this is larger than any reasonable significance level, we fail to reject  $H_0$ . The data provides no statistically significant evidence that the populations of social and natural scientists differ with respect to degree of spirituality.

#### Chi-Square Test: Very, Moderate, Slightly, Not At All

Expected counts are printed below observed counts Chi-Square contributions are printed below expected counts

	Verv	Moderate	Slightly	Not At All	Total	
1	56	162	198	211	627	
	50.59	173.92	199.21	203.28		
	0.578	0.816	0.007	0.293		
2	56	223	243	239	761	
	61.41	211.08	241.79	246.72		
	0.476	0.673	0.006	0.242		
Total	112	385	441	450	1388	
chi-Sa	= 3.09	1. $DF = 3$ .	P-Value =	0.378		

Telegram: @uni\_k

#### 31.

**a.** The accompanying table shows the proportions of male and female smokers in the sample who began smoking at the ages specified. (The male proportions were calculated by dividing the counts by the total of 96; for females, we divided by 93.) The patterns of the proportions seems to be different, suggesting there does exist an association between gender and age at first smoking.

		Gender				
	ALC: NO	Male	Female			
	<16	0.26	0.11			
Age	16-17	0.25	0.34			
-8-	18-20	0.29	0.18			
	>20	0.20	0.37			

**b.** The hypotheses, in words, are  $H_0$ : gender and age at first smoking are independent, versus  $H_a$ : gender and age at first smoking are associated. The accompanying Minitab output provides a test statistic value of  $\chi^2 = 14.462$  at df = (2-1)(4-1) = 3, with an associated *P*-value of 0.002. Hence, we would reject  $H_0$  at both the .05 and .01 levels. We have evidence to suggest an association between gender and age at first smoking.

### Chi-Square Test: Male, Female

Expected counts are printed below observed counts Chi-Square contributions are printed below expected counts Male Female Total

	Male	Female	Total	
1	25	10	35	
	17.78	17.22		
	2.934	3.029		
2	24	32	56	
	28.44	27.56		
	0.694	0.717		
3	28	17	45	
	22.86	22.14		
	1.157	1.194		
4	19	34	53	
	26.92	26.08		
	2.330	2.406		
otal	96	93	189	

Chi-Sq = 14.462, DF = 3, P-Value = 0.002

33.

χ

$${}^{2} = \Sigma \Sigma \frac{\left(N_{ij} - \hat{E}_{ij}\right)^{2}}{\hat{E}_{ij}} = \Sigma \Sigma \frac{N_{ij}^{2} - 2\hat{E}_{ij}N_{ij} + \hat{E}_{ij}^{2}}{\hat{E}_{ij}} = \Sigma \Sigma \frac{N_{ij}^{2}}{\hat{E}_{ij}} - 2\Sigma \Sigma N_{ij} + \Sigma \Sigma \hat{E}_{ij}, \text{ but } \Sigma \Sigma \hat{E}_{ij} = \Sigma \Sigma N_{ij} = n, \text{ so}$$

 $\chi^2 = \Sigma \Sigma \frac{N_{ij}^2}{\hat{E}_{ij}} - n$ . This formula is computationally efficient because there is only one subtraction to be

performed, which can be done as the last step in the calculation.

35.

With  $p_{ij}$  denoting the common value of  $p_{ij1}$ ,  $p_{ij2}$ ,  $p_{ij3}$ , and  $p_{ij4}$  under  $H_0$ ,  $\hat{p}_{ij} = \frac{n_{ij.}}{n}$  and  $\hat{E}_{ijk} = \frac{n_k n_{ij.}}{n}$ , where

$$n_{ij} = \sum_{k=1}^{4} n_{ijk}$$
 and  $n = \sum_{k=1}^{4} n_k$ . With four different tables (one for each region), there are  $4(9-1) = 32$ 

freely determined cell counts. Under  $H_0$ , the nine parameters  $p_{11}, \ldots, p_{33}$  must be estimated, but  $\Sigma\Sigma p_{ij} = 1$ , so only 8 independent parameters are estimated, giving  $\chi^2 df = 32 - 8 = 24$ . Note: this is really a test of homogeneity for 4 strata, each with 3x3=9 categories. Hence, df = (4-1)(9-1) = 24.

### Supplementary Exercises

37. There are 3 categories here – firstborn, middleborn, (2<sup>nd</sup> or 3<sup>rd</sup> born), and lastborn. With  $p_1$ ,  $p_2$ , and  $p_3$  denoting the category probabilities, we wish to test  $H_0$ :  $p_1 = .25$ ,  $p_2 = .50$ ,  $p_3 = .25$  because  $p_2 = P(2^{nd} \text{ or } 3^{rd} \text{ born}) = .25 + .25 = .50$ . The expected counts are (31)(.25) = 7.75, (31)(.50) = 15.5, and 7.75, so  $\chi^2 = \frac{(12 - 7.75)^2}{7.75} + \frac{(11 - 15.5)^2}{15.5} + \frac{(8 - 7.75)^2}{7.75} = 3.65$ . At df = 3 - 1 = 2,  $3.65 < 5.992 \Rightarrow P$ -value > .05  $\Rightarrow H_0$  is not rejected. The hypothesis of equiprobable birth order appears plausible.

39.

- a. For that top-left cell, the estimated expected count is (row total)(column total)/(grand total) = (189)(406)/(852) = 90.06. Next, the chi-squared contribution is  $(O E)^2/E = (83 90.06)^2/90.06 = 0.554$ .
- b. <u>No</u>: From the software output, the *P*-value is .023 > .01. Hence, we fail to reject the null hypothesis of "no association" at the .01 level. We have insufficient evidence to conclude that an association exists between cognitive state and drug status. [*Note*: We would arrive at a different conclusion for  $\alpha = .05$ .]
- 41. The null hypothesis  $H_0: p_{ij} = p_i p_j$  states that level of parental use and level of student use are independent in the population of interest. The test is based on (3-1)(3-1) = 4 df.

Estimated ex	pecieu counts	
57.6	58.1	235
33.9	40.3	163
11.5	11.6	47
109	110	445
	57.6 33.9 11.5 109	57.6         58.1           33.9         40.3           11.5         11.6           109         110

Estimated expected counts

The calculated test statistic value is  $\chi^2 = 22.4$ ; at df = (3-1)(3-1) = 4, the *P*-value is < .001, so  $H_0$  should be rejected at any reasonable significance level. Parental and student use level do not appear to be independent.

43.

ikur.in

This is a test of homogeneity:  $H_0$ :  $p_{1j} = p_{2j} = p_{3j}$  for j = 1, 2, 3, 4, 5. The given SPSS output reports the calculated  $\chi^2 = 70.64156$  and accompanying *P*-value (significance) of .0000. We reject  $H_0$  at any significance level. The data strongly supports that there are differences in perception of odors among the three areas.

$$(n_1 - np_{10})^2 = (np_{10} - n_1)^2 = (n - n_1 - n(1 - p_{10}))^* = (n_2 - np_{20})^2 . \text{ Therefore}$$

$$\chi^2 = \frac{(n_1 - np_{10})^2}{np_{10}} + \frac{(n_2 - np_{20})^2}{np_{20}} = \frac{(n_1 - np_{10})^2}{n_2} \left(\frac{n}{p_{10}} + \frac{n}{p_{20}}\right)$$

$$= \left(\frac{n_1}{n} - p_{10}\right)^2 \cdot \left(\frac{n}{p_{10}p_{20}}\right) = \frac{(\hat{p}_1 - p_{10})^2}{p_{10}p_{20} / n} = z^2$$

47.

a. Our hypotheses are H<sub>0</sub>: no difference in proportion of concussions among the three groups v. H<sub>a</sub>: there is a difference in proportion of concussions among the three groups.

Observed	Concussion	No Concussion	Total
Soccer	45	46	91
Non Soccer	28	68	96
Control	8	45	53
Total	81	159	240
Expected	Concussion	No Concussion	Total
Expected	30 7125	60.2875	91
Non Soccer	32.4	63.6	96
Control	17 8875	37.1125	53
Total	81	159	240

$$\chi^{2} = \frac{(45 - 30.7125)^{2}}{30.7125} + \frac{(46 - 60.2875)^{2}}{60.2875} + \frac{(28 - 32.4)^{2}}{32.4} + \frac{(68 - 63.6)^{2}}{63.6}$$

 $+\frac{(8-17.8875)^2}{17.8875} + \frac{(45-37.1125)^2}{37.1125} = 19.1842$ . The df for this test is (I-1)(J-1) = 2, so the *P*-value is less than .001 and we reject H<sub>0</sub>. There is a difference in the proportion of concussions based on whether a person plays soccer.

**b.** The sample correlation of r = -.220 indicates a weak negative association between "soccer exposure" and immediate memory recall. We can formally test the hypotheses  $H_0$ :  $\rho = 0$  vs  $H_a$ :  $\rho < 0$ . The test

statistic is  $t = \frac{r\sqrt{n-2}}{\sqrt{1-r^2}} = \frac{-.22\sqrt{89}}{\sqrt{1-.22^2}} = -2.13$ . At significance level  $\alpha = .01$ , we would fail to reject  $H_0$ 

and conclude that there is no significant evidence of negative association in the population.

- c. We will test to see if the average score on a controlled word association test is the same for soccer and non-soccer athletes. H<sub>0</sub>: μ<sub>1</sub> = μ<sub>2</sub> vs H<sub>a</sub>: μ<sub>1</sub> ≠ μ<sub>2</sub>. Since the two sample standard deviations are very close, we will use a pooled-variance two-sample t test. From Minitab, the test statistic is t = -0.91, with an associated P-value of 0.366 at 80 df. We clearly fail to reject H<sub>0</sub> and conclude that there is no statistically significant difference in the average score on the test for the two groups of athletes.
- **d.** Our hypotheses for ANOVA are  $H_0$ : all means are equal vs  $H_a$ : not all means are equal. The test statistic is  $f = \frac{MSTr}{MST}$ .

$$MSE$$
  

$$SSTr = 91(.30 - .35)^{2} + 96(.49 - .35)^{2} + 53(.19 - .35)^{2} = 3.4659 \ MSTr = \frac{3.4659}{2} = 1.73295$$
  

$$SSE = 90(.67)^{2} + 95(.87)^{2} + 52(.48)^{2} = 124.2873 \text{ and } MSE = \frac{124.2873}{237} = .5244.$$

Now,  $f = \frac{1.73295}{.5244} = 3.30$ . Using df = (2,200) from Table A.9, the *P*-value is between .01 and .05. At

.5244 significance level .05, we reject the null hypothesis. There is sufficient evidence to conclude that there is a difference in the average number of prior non-soccer concussions between the three groups.

**49.** According to Benford's law, the probability a lead digit equals x is given by  $\log_{10}(1 + 1/x)$  for x = 1, ..., 9. Let  $p_i$  = the proportion of Fibonacci numbers whose lead digit is i (i = 1, ..., 9). We wish to perform a goodness-of-fit test  $H_0$ :  $p_i = \log_{10}(1 + 1/i)$  for i = 1, ..., 9. (The alternative hypothesis is that Benford's formula is incorrect for at least one category.) The table below summarizes the results of the test.

Digit	1	2	3	4	5	6	7	8	9
Obs. #	25	16	11	7	7	5	4	6	4
Exp. #	25.59	14.97	10.62	8.24	6.73	5.69	4.93	4.35	3.89

Expected counts are calculated by  $np_i = 85 \log_{10}(1 + 1/i)$ . Some of the expected counts are too small, so combine 6 and 7 into one category (obs = 9, exp = 10.62); do the same to 8 and 9 (obs = 10, exp = 8.24).

The resulting chi-squared statistic is  $\chi^2 = \frac{(25-25.59)^2}{25.59} + \dots + \frac{(10-8.24)^2}{8.24} = 0.92$  at df = 7 - 1 = 6 (since there are 7 categories after the earlier combining). Software provides a *P*-value of .988!

We certainly do not reject  $H_0$  — the lead digits of the Fibonacci sequence are highly consistent with Benford's law.

# CHAPTER 15

### Section 15.1

5.

www.konkur.in

- 1. Refer to Table A.13. a. With n = 12,  $P_0(S_+ \ge 56) = .102$ .
  - **b.** With n = 12,  $61 < 62 < 64 \Rightarrow P_0(S_+ \ge 62)$  is between .046 and .026.
  - c. With n = 12 and a lower-tailed test, P-value =  $P_0(S_+ \ge n(n+1)/2 s_+) = P_0(S_+ \ge 12(13)/2 20) = P_0(S_+ \ge 58)$ . Since 56 < 58 < 60, the P-value is between .055 and .102.
  - **d.** With n = 14 and a two-tailed test, P-value =  $2P_0(S_+ \ge \max\{21, 14(15)/2 21\}) = 2P_0(S_+ \ge 84) = .025$ .
  - e. With n = 25 being "off the chart," use the large-sample approximation:  $z = \frac{s_{+} - n(n+1)/4}{\sqrt{n(n+1)(2n+1)/24}} = \frac{300 - 25(26)/4}{\sqrt{25(26)(51)/24}} = 3.7 \Rightarrow \text{two-tailed } P \text{-value} = 2P(Z \ge 3.7) \approx 0.$
- 3. We test  $H_0: \mu = 7.39$  vs.  $H_a: \mu \neq 7.39$ , so a two tailed test is appropriate. The  $(x_i 7.39)$ 's are -.37, -.04, -.05, -.22, -.11, .38, -.30, -.17, .06, -.44, .01, -.29, -.07, and -.25, from which the ranks of the three positive differences are 1, 4, and 13. Thus  $s_+ = 1 + 4 + 13 = 18$ , and the two-tailed *P*-value is given by  $2P_0(S_+ \ge \max\{18, 14(15)/2 18\}) = 2P_0(S_+ \ge 87)$ , which is between 2(.025) and 2(.010) or .05 and .02. In particular, since *P*-value < .05,  $H_0$  is rejected at level .05.

$d_i$	3	2.8	3.9	.6	1.2	-1.1	2.9	1.8	.5	2.3	.9	2.5
rank	1	10*	12*	3*	6*	5	11*	7*	2*	8*	4*	9*

7. The data are paired, and we wish to test  $H_0$ :  $\mu_D = .20$  vs.  $H_a$ :  $\mu_D > .20$  where  $\mu_D = \mu_{outdoor} - \mu_{indoor}$ . Because n = 33, we'll use the large-sample test.

d.	$d_i - 2$	rank	d,	$d_i2$	rank	di	$d_i2$	rank	
0.22	0.02	2	0.15	-0.05	5.5	0.63	0.43	23	
0.01	_0.19	17	1.37	1.17	32	0.23	0.03	4	
0.38	0.18	16	0.48	0.28	21	0.96	0.76	31	
0.30	0.10	10	0.10	-0.09	8	0.2	0	1	
0.42	0.44	20	0.03	-0.17	15	-0.02	-0.22	18	
0.85	0.03	3	0.83	0.63	28	0.03	-0.17	14	
0.25	0.05	13	1 30	1.19	33	0.87	0.67	30	
0.30	0.10	15	0.68	0.48	25	0.3	0.1	9.5	
0.7	0.5	20	0.00	0.10	9.5	0.31	0.11	11	
0.71	0.51	21	0.11	_0.31	22	0.45	0.25	20	
0.13	-0.07	5.5	0.31	0.11	12	-0.26	-0.46	24	

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

#### Chapter 15: Distribution-Free Procedures

From the table,  $s_{+} = 424$ , so  $z = \frac{s_{+} - n(n+1)/4}{\sqrt{n(n+1)(2n+1)/24}} = \frac{424 - 280.5}{\sqrt{3132.25}} = \frac{143.5}{55.9665} = 2.56$ . The upper-tailed

*P*-value is  $P(Z \ge 2.56) = .0052 < .05$ , so we reject H0. There is statistically significant evidence that the true mean difference between outdoor and indoor concentrations exceeds .20 nanograms/m<sup>3</sup>.

$R_1$	1	1	1	1	1	1	2	2	2	2	2	2
$R_2$	2	2	3	3	4	4	1	1	3	3	4	4
$R_3$	3	4	2	4	2	3	3	4	1	4	1	3
$R_4$	4	3	4	2	3	2	4	3	4	1	3	1
D	0	2	2	6	6	8	2	4	6	12	10	14
$R_1$	3	3	3	3	3	3	4	4	4	4	4	4
$R_2$	1	1	2	2	4	4	1	1	2	2	3	3
$R_3$	2	4	1	4	1	2	2	3	1	3	1	2
$R_4$	4	2	4	1	2	1	3	2	3	1	2	1
D	6	10	8	14	16	18	12	14	14	18	18	20

When  $H_0$  is true, each of the above 24 rank sequences is equally likely, which yields the distribution of D:

d	0	2	4	6	8	10	12	14	16	18	20
p(d)	1/24	3/24	1/24	4/24	2/24	2/24	2/24	4/24	1/24	3/24	1/24

Then c = 0 yields  $\alpha = 1/24 = .042$  (too small) while c = 2 implies  $\alpha = 1/24 + 3/24 = .167$ , and this is the closest we can come to achieving a .10 significance level.

### Section 15.2

9.

- 11. The ordered combined sample is 163(y), 179(y), 213(y), 225(y), 229(x), 245(x), 247(y), 250(x), 286(x), and 299(x), so w = 5 + 6 + 8 + 9 + 10 = 38. With m = n = 5, Table A.14 gives *P*-value =  $P_0(W \ge 38)$ , which is between .008 and .028. In particular, *P*-value < .05, so  $H_0$  is rejected in favor of  $H_a$ .
- 13. Identifying x with unpolluted region (m = 5) and y with polluted region (n = 7), we wish to test the hypotheses  $H_0: \mu_1 \mu_2 = 0$  vs.  $H_a: \mu_1 \mu_2 < 0$ . The x ranks are 1, 5, 4, 6, 9, so w = 25. In this particular order, the test is lower-tailed, so P-value  $= P_0(W \ge 5(5+7+1)-25) = P_0(W \ge 40) > .053$ . So, we fail to reject  $H_0$  at the .05 level: there is insufficient evidence to conclude that the true average fluoride level is higher in polluted areas.
- 15. Let  $\mu_1$  and  $\mu_2$  denote true average cotanine levels in unexposed and exposed infants, respectively. The hypotheses of interest are  $H_0: \mu_1 \mu_2 = -25$  vs.  $H_0: \mu_1 \mu_2 < -25$ . Before ranking, -25 is subtracted from each  $x_i$  (i.e. 25 is added to each), giving 33, 36, 37, 39, 45, 68, and 136. The corresponding x ranks in the combined set of 15 observations are 1, 3, 4, 5, 6, 8, and 12, from which w = 1 + 3 + ... + 12 = 39. With m = 7 and n = 8, P-value  $= P_0(W \ge 7(7 + 8 + 1) 39) = P_0(W \ge 73) = .027$ . Therefore,  $H_0$  is rejected at the .05 level. The true average level for exposed infants appears to exceed that for unexposed infants by more than 25 (note that  $H_0$  would not be rejected using level .01).

ww.konkur.in

### Section 15.3

www.konkur.in

- 17. n = 8, so from Table A.15, a 95% CI (actually 94.5%) has the form  $(\overline{x}_{(36-32+1)}, \overline{x}_{(32)}) = (\overline{x}_{(5)}, \overline{x}_{(32)})$ . It is easily verified that the 5 smallest pairwise averages are  $\frac{5.0 + 5.0}{2} = 5.00$ ,  $\frac{5.0 + 11.8}{2} = 8.40$ ,  $\frac{5.0 + 12.2}{2} = 8.60$ ,  $\frac{5.0 + 17.0}{2} = 11.00$ , and  $\frac{5.0 + 17.3}{2} = 11.15$  (the smallest average not involving 5.0 is  $\overline{x}_{(6)} = \frac{11.8 + 11.8}{2} = 11.8$ ), and the 5 largest averages are 30.6, 26.0, 24.7, 23.95, and 23.80, so the confidence interval is (11.15, 23.80).
- 19. First, we must recognize this as a paired design; the eight <u>differences</u> (Method 1 minus Method 2) are -0.33, -0.41, -0.71, 0.19, -0.52, 0.20, -0.65, and -0.14. With n = 8, Table A.15 gives c = 32, and a 95% CI for  $\mu_D$  is  $(\overline{x}_{8(8+1)/2-32+1}, \overline{x}_{32}) = (\overline{x}_{(5)}, \overline{x}_{(32)})$ .

Of the 36 pairwise averages created from these 8 differences, the 5<sup>th</sup> smallest is  $\overline{x}_{(5)} = -0.585$ , and the 5<sup>th</sup>-largest (aka the 32<sup>nd</sup>-smallest) is  $\overline{x}_{(32)} = 0.025$ . Therefore, we are 94.5% confident the true mean difference in extracted creosote between the two solvents,  $\mu_D$ , lies in the interval (-.585, .025).

21. m = n = 5 and from Table A.16, c = 21 and the 90% (actually 90.5%) interval is  $(d_{ij(5)}, d_{ij(21)})$ . The five smallest  $x_i - y_j$  differences are -18, -2, 3, 4, 16 while the five largest differences are 136, 123, 120, 107, 87 (construct a table like Table 15.5), so the desired interval is (16, 87).

### Section 15.4

23. Below we record in parentheses beside each observation the rank of that observation in the combined sample.

1:	5.8(3)	6.1(5)	6.4(6)	6.5(7)	7.7(10)	$r_{1.} = 31$
2:	7.1(9)	8.8(12)	9.9(14)	10.5(16)	11.2(17)	$r_{2.} = 68$
3:	5.1(1)	5.7(2)	5.9(4)	6.6(8)	8.2(11)	$r_{3.} = 26$
4:	9.5(13)	1.0.3(15)	11.7(18)	12.1(19)	12.4(20)	<i>r</i> <sub>4.</sub> = 85

The computed value of k is  $k = \frac{12}{20(21)} \left[ \frac{31^2 + 68^2 + 26^2 + 85^2}{5} \right] - 3(21) = 14.06$ . At 3 df, the P-value is <

.005, so we reject  $H_0$ .

25. The ranks are 1, 3, 4, 5, 6, 7, 8, 9, 12, 14 for the first sample; 11, 13, 15, 16, 17, 18 for the second; 2, 10, 19, 20, 21, 22 for the third; so the rank totals are 69, 90, and 94.

 $k = \frac{12}{22(23)} \left[ \frac{69^2}{10} + \frac{90^2}{6} + \frac{94^2}{5} \right] - 3(23) = 9.23; \text{ at } 2 \text{ df, the } P \text{-value is roughly .01. Therefore, we reject}$ 

 $H_0: \mu_1 = \mu_2 = \mu_3$  at the .05 level.

### Chapter 15: Distribution-Free Procedures

	1	2	3	4	5	6	7	8	9	10	$r_i$	$r_i^2$
I	1	2	3	3	2	1	1	3	1	2	19	361
H	2	1	1	2	1	2	2	1	2	3	17	289
с	3	3	2	1	3	3	3	2	3	1	24	576
H.C.	1											1226

The computed value of  $F_r$  is  $\frac{12}{10(3)(4)}(1226)-3(10)(4)=2.60$ . At 2 df, *P*-value > .10, and so we don't reject  $H_0$  at the .05 level.

Supplementary Exercises

- 29. Friedman's test is appropriate here. It is easily verified that  $r_1 = 28$ ,  $r_2 = 29$ ,  $r_3 = 16$ ,  $r_4 = 17$ , from which the defining formula gives  $f_r = 9.62$  and the computing formula gives  $f_r = 9.67$ . Either way, at 3 df the *P*-value is < .025, and so we reject  $H_0: \alpha_1 = \alpha_2 = \alpha_3 = \alpha_4 = 0$  at the .05 level. We conclude that there are effects due to different years.
- **31.** From Table A.16, m = n = 5 implies that c = 22 for a confidence level of 95%, so mn c + 1 = 25 22 = 1 = 4. Thus the confidence interval extends from the 4<sup>th</sup> smallest difference to the 4<sup>th</sup> largest difference. The 4 smallest differences are -7.1, -6.5, -6.1, -5.9, and the 4 largest are -3.8, -3.7, -3.4, -3.2, so the CI is (-5.9, -3.8).

#### 33.

- a. With "success" as defined, then Y is binomial with n = 20. To determine the binomial proportion p, we realize that since 25 is the hypothesized median, 50% of the distribution should be above 25, thus when  $H_0$  is true p = .50. The upper-tailed P-value is  $P(Y \ge 15$  when  $Y \sim Bin(20, .5)) = 1 B(14; 20, .5) = .021$ .
- **b.** For the given data, y = (# of sample observations that exceed 25) = 12. Analogous to **a**, the *P*-value is then  $P(Y \ge 12 \text{ when } Y \sim \text{Bin}(20, .5)) = 1 B(11; 20, .5) = .252$ . Since the *P*-value is large, we fail to reject  $H_0$  we have insufficient evidence to conclude that the population median exceeds 25.
- 35.

Sample:	у	x	y	y	x	x	x	y	y
Observations:	3.7	4.0	4.1	4.3	4.4	4.8	4.9	5.1	5.6
Rank:	1	3	5	7	9	8	6	4	2

The value of W' for this data is w' = 3 + 6 + 8 + 9 = 26. With m = 4 and n = 5, he upper-tailed P-value is  $P_0(W \ge 26) > .056$ . Thus,  $H_0$  cannot be rejected at level .05.

www.konkur.in

# **CHAPTER 16**

### Section 16.1

5.

www.konkur.in

 All ten values of the quality statistic are between the two control limits, so no out-of-control signal is generated.

3.  $P(10 \text{ successive points inside the limits}) = P(1^{\text{st}} \text{ inside}) \times P(2^{\text{nd}} \text{ inside}) \times \dots \times P(10^{\text{th}} \text{ inside}) = (.998)^{10} = .9802$ .  $P(25 \text{ successive points inside the limits}) = (.998)^{25} = .9512$ .  $(.998)^{52} = .9011$ , but  $(.998)^{53} = .8993$ , so for 53 successive points the probability that at least one will fall outside the control limits when the process is in control is 1 - .8993 = .1007 > .10.

**a.** For the case of 4(a), with  $\sigma = .02$ ,  $C_p = \frac{\text{USL} - \text{LSL}}{6\sigma} = \frac{3.1 - 2.9}{6(.02)} = 1.67$ . This is indeed a very good capability index. In contrast, the case of 4(b) with  $\sigma = .05$  has a capability index of  $C_p = \frac{3.1 - 2.9}{6(.05)}$ 0.67. This is quite a bit less than 1, the dividing line for "marginal capability."

**b.** For the case of 4(a), with  $\mu = 3.04$  and  $\sigma = .02$ ,  $\frac{USL - \mu}{3\sigma} = \frac{3.1 - 3.04}{3(.02)} = 1$  and  $\frac{\mu - LSL}{3\sigma} = \frac{3.04 - 2.9}{3(.02)} = 2.33$ , so  $C_{pk} = \min\{1, 2.33\} = 1$ . For the case of 4(b), with  $\mu = 3.00$  and  $\sigma = .05$ ,  $\frac{USL - \mu}{3\sigma} = \frac{3.1 - 3.00}{3(.05)} = .67$  and  $\frac{\mu - LSL}{3\sigma} = \frac{3.00 - 2.9}{3(.05)} = .67$ , so  $C_{pk} = \min\{.67, .67\} = .67$ . Even using this mean-adjusted capability index, process (a) is more "capable" than process (b), though  $C_{pk}$  for process (a) is now right at the "marginal capability" threshold. **c.** In general,  $C_{pk} \le C_p$ , and they are equal iff  $\mu = \frac{LSL + USL}{2}$ , i.e. the process mean is the midpoint of the spec limits. To demonstrate this, suppose first that  $\mu = \frac{LSL + USL}{2}$ . Then  $\frac{USL - \mu}{3\sigma} = \frac{USL - (LSL + USL)/2}{3\sigma} = \frac{2USL - (LSL + USL)}{6\sigma} = \frac{USL - LSL}{6\sigma} = C_p$ , and similarly  $\frac{\mu - LSL}{3\sigma} = C_p$ . In that case,  $C_{pk} = \min\{C_p, C_p\} = C_p$ . Otherwise, suppose  $\mu$  is closer to the lower spec limit than to the upper spec limit (but between the two), so that  $\mu - LSL < USL - \mu$ . In such a case,  $C_{pk} = \frac{\mu - LSL}{3\sigma}$ . However, in this same case  $\mu < \frac{LSL + USL}{2}$ , from which  $\frac{\mu - LSL}{3\sigma} < \frac{(LSL + USL)/2 - LSL}{3\sigma} = \frac{USL - LSL}{6\sigma} = C_p$ . That is,  $C_{pk} < C_p$ .

Analogous arguments for all other possible values of  $\mu$  also yield  $C_{pk} < C_p$ .

209

### Section 16.2

7.

- a. P(point falls outside the limits when  $\mu = \mu_0 + .5\sigma$ ) =  $1 P\left(\mu_0 \frac{3\sigma}{\sqrt{n}} < \overline{X} < \mu_0 + \frac{3\sigma}{\sqrt{n}} \text{ when } \mu = \mu_0 + .5\sigma\right)$ =  $1 - P\left(-3 - .5\sqrt{n} < Z < 3 - .5\sqrt{n}\right)$  =  $1 - P\left(-4.12 < Z < 1.882\right) = 1 - .9699 = .0301$ .
- **b.**  $1 P\left(\mu_0 \frac{3\sigma}{\sqrt{n}} < \overline{X} < \mu_0 + \frac{3\sigma}{\sqrt{n}} \text{ when } \mu = \mu_0 \sigma\right) = 1 P\left(-3 + \sqrt{n} < Z < 3 + \sqrt{n}\right)$ =  $1 - P\left(-.76 < Z < 5.24\right) = .2236$

c. 
$$1 - P(-3 - 2\sqrt{n} < Z < 3 - 2\sqrt{n}) = 1 - P(-7.47 < Z < -1.47) = .9292$$

$$\overline{x} = 12.95$$
 and  $\overline{s} = .526$ , so with  $a_5 = .940$ , the control limits are  
 $12.95 \pm 3 \frac{.526}{.940\sqrt{5}} = 12.95 \pm .75 = 12.20, 13.70$ . Again, every point  $(\overline{x})$  is between these limits, so there is no

evidence of an out-of-control process.

9.

$$=\frac{2517.07}{24}=96.54$$
,  $\overline{s}=1.264$ , and  $a_6=.952$ , giving the control limits

 $96.54 \pm 3 \frac{1.264}{.952\sqrt{6}} = 96.54 \pm 1.63 = 94.91,98.17$ . The value of  $\overline{x}$  on the 22<sup>nd</sup> day lies above the UCL, so the process appears to be out of control at that time.

13.

$$P\left(\mu_0 - \frac{2.81\sigma}{\sqrt{n}} < \overline{X} < \mu_0 + \frac{2.81\sigma}{\sqrt{n}} \text{ when } \mu = \mu_0\right) = P(-2.81 < Z < 2.81) = .995, \text{ so the probability that a point falls outside the limits is .005 and ARL} = \frac{1}{.005} = 200.$$

**b.**  $P(\text{a point is <u>inside</u> the limits}) = P\left(\mu_0 - \frac{2.81\sigma}{\sqrt{n}} < \overline{X} < \mu_0 + \frac{2.81\sigma}{\sqrt{n}} \text{ when } \mu = \mu_0 + \sigma\right) = \dots = P\left(-2.81 - \sqrt{n} < Z < 2.81 - \sqrt{n}\right) = P(-4.81 < Z < .81) \text{ [when } n = 4\text{]} \approx \Phi(.81) = .7910 \Rightarrow p = P(\text{a point is outside the limits}) = 1 - .7910 = .209 \Rightarrow \text{ARL} = \frac{1}{2.090} = 4.78.$ 

c. Replace 2.81 with 3 above. For **a**, P(-3 < Z < 3) = .9974, so p = 1 - .9974 = .0026 and ARL =  $\frac{1}{.0026} = 385$  for an in-control process. When  $\mu = \mu_0 + \sigma$  as in **b**, the probability of an <u>out</u>-ofcontrol point is  $1 - P(-3 - \sqrt{n} < Z < 3 - \sqrt{n}) = 1 - P(-5 < Z < 1) \approx 1 - \Phi(1) = .1587$ , so ARL =  $\frac{1}{.1587} = 6.30$ .
15.  $\overline{\overline{x}} = 12.95$ , IQR = .4273,  $k_5 = .990$ . The control limits are  $12.95 \pm 3 \frac{.4273}{.990\sqrt{5}} = 12.37, 13.53$ .

#### Section 16.3

17.

www.konkur.in

$$\overline{r} = \frac{85.2}{30} = 2.84 \text{, } b_4 = 2.058 \text{, and } c_4 = .880. \text{ Since } n = 4, \text{ LCL} = 0 \text{ and UCL}$$
$$= 2.84 + \frac{3(.880)(2.84)}{2.058} = 2.84 + 3.64 = 6.48.$$

**b.** 
$$\overline{r} = 3.54$$
,  $b_8 = 2.844$ , and  $c_8 = .820$ , and the control limits are  
 $3.54 \pm \frac{3(.820)(3.54)}{2.844} = 3.54 \pm 3.06 = .48, 6.60$ .

19.

$$s = 1.2642$$
,  $a_6 = .952$ , and the control limits are  
 $1.2642 \pm \frac{3(1.2642)\sqrt{1-(.952)^2}}{.952} = 1.2642 \pm 1.2194 = .045, 2.484$ . The smallest s<sub>1</sub> is s<sub>20</sub> = .75, and the largest  
is s<sub>12</sub> = 1.65, so every value is between .045 and 2.434. The process appears to be in control with respect to  
variability.

#### Section 16.4

21.

2

$$\overline{p} = \Sigma \frac{\hat{p}_i}{k} \text{ where } \Sigma \hat{p}_i = \frac{x_1}{n} + \dots + \frac{x_k}{n} = \frac{x_1 + \dots + x_k}{n} = \frac{578}{100} = 5.78 \text{ . Thus } \overline{p} = \frac{5.78}{25} = .231$$
**a.** The control limits are  $.231 \pm 3\sqrt{\frac{(.231)(.769)}{100}} = .231 \pm .126 = .105,.357 \text{ .}$ 

**b.**  $\frac{13}{100} = .130$ , which is between the limits, but  $\frac{39}{100} = .390$ , which exceeds the upper control limit and therefore generates an out-of-control signal.

3. LCL > 0 when 
$$\overline{p} > 3\sqrt{\frac{\overline{p}(1-\overline{p})}{n}}$$
, i.e. (after squaring both sides)  $50\overline{p}^2 > 9\overline{p}(1-\overline{p})$ , i.e.  $50\overline{p} > 3(1-\overline{p})$ , i.e.  
 $53\overline{p} > 3 \Rightarrow \overline{p} = \frac{3}{53} = .0566$ .

25.  $\Sigma x_i = 102$ ,  $\overline{x} = 4.08$ , and  $\overline{x} \pm 3\sqrt{\overline{x}} = 4.08 \pm 6.06 \approx (-2.0, 10.1)$ . Thus LCL = 0 and UCL = 10.1. Because no  $x_i$  exceeds 10.1, the process is judged to be in control.

27. With 
$$u_i = \frac{x_i}{g_i}$$
, the  $u_i$ 's are 3.75, 3.33, 3.75, 2.50, 5.00, 5.00, 12.50, 12.00, 6.67, 3.33, 1.67, 3.75, 6.25, 4.00,  
6.00, 12.00, 3.75, 5.00, 8.33, and 1.67 for  $i = 1, ..., 20$ , giving  $\overline{u} = 5.5125$ . For  $g_i = .6$ ,  
 $\overline{u} \pm 3\sqrt{\frac{\overline{u}}{g_i}} = 5.5125 \pm 9.0933$ , LCL = 0, UCL = 14.6. For  $g_i = .8$ ,  $\overline{u} \pm 3\sqrt{\frac{\overline{u}}{g_i}} = 5.5125 \pm 7.857$ , LCL = 0,  
UCL = 13.4. For  $g_i = 1.0$ ,  $\overline{u} \pm 3\sqrt{\frac{\overline{u}}{g_i}} = 5.5125 \pm 7.0436$ , LCL = 0, UCL = 12.6. Several  $u_i$ 's are close to  
the corresponding LICL's but none exceed them, so the process is judged to be in control.

## Section 16.5

1

the conc

$$\mu_{0} = 16, \ k = \frac{\Delta}{2} = 0.05, \ h = .20, \ d_{i} = \max\left(0, d_{i-1} + (\overline{x}_{i} - 16.05)\right), \ e_{i} = \max\left(0, e_{i-1} + (\overline{x}_{i} - 15.95)\right).$$

$$\frac{i \quad \overline{x}_{i} - 16.05}{1} \quad \frac{d_{i} \quad \overline{x}_{i} - 15.95}{2} \quad \frac{e_{i}}{0}$$

$$\frac{i \quad \overline{x}_{i} - 16.05}{2} \quad 0.001 \quad 0.001 \quad 0.101 \quad 0$$

$$3 \quad 0.016 \quad 0.017 \quad 0.116 \quad 0$$

$$4 \quad -0.138 \quad 0 \quad -0.038 \quad 0.038$$

$$5 \quad -0.020 \quad 0 \quad 0.080 \quad 0$$

$$6 \quad 0.010 \quad 0.010 \quad 0.110 \quad 0$$

$$7 \quad -0.068 \quad 0 \quad 0.032 \quad 0$$

$$8 \quad -0.151 \quad 0 \quad -0.054 \quad 0.054$$

$$9 \quad -0.012 \quad 0 \quad 0.088 \quad 0$$

$$10 \quad 0.024 \quad 0.024 \quad 0.124 \quad 0$$

$$11 \quad -0.021 \quad 0.003 \quad 0.079 \quad 0$$

$$12 \quad -0.115 \quad 0 \quad -0.015 \quad 0.015$$

$$13 \quad -0.018 \quad 0 \quad 0.082 \quad 0$$

$$14 \quad -0.090 \quad 0 \quad 0.010 \quad 0$$

For no time r is it the case that  $d_r > .20$  or that  $e_r > .20$ , so no out-of-control signals are generated.

Connecting 600 on the in-control ARL scale to 4 on the out-of-control scale and extending to the k' scale gives k' = .87. Thus  $k' = \frac{\Delta/2}{\sigma/\sqrt{n}} = \frac{.002}{.005/\sqrt{n}}$  from which  $\sqrt{n} = 2.175 \Rightarrow n = 4.73 = s$ . Then connecting .87 31. on the k' scale to 600 on the out-of-control ARL scale and extending to h' gives h' = 2.8, so

$$h = \left(\frac{\sigma}{\sqrt{n}}\right) (2.8) = \left(\frac{.005}{\sqrt{5}}\right) (2.8) = .00626$$
.

### Section 16.6

For the binomial calculation, n = 50 and we wish 33.

$$P(X \le 2) = {\binom{50}{0}} p^0 (1-p)^{50} + {\binom{50}{1}} p^1 (1-p)^{49} + {\binom{50}{2}} p^2 (1-p)^{48} \text{ when } p = .01, .02, ..., .10. \text{ For the}$$
  
hypergeometric calculation,  $P(X \le 2) = {\binom{M}{0}} {\binom{500-M}{50}} + {\binom{M}{1}} {\binom{500-M}{49}} + {\binom{M}{2}} {\binom{500-M}{48}}, \text{ to be}$ 

calculated for M = 5, 10, 15, ..., 50. The resulting probabilities appear below.

	01	02	.03	.04	.05	.06	.07	.08	.09	.10
p	.01	.02	.00	0-10-	tus ter	10.17	2064	2110	.1464	.0994
Hypg.	.9919	.9317	.8182	.6775	.5343	.4047	.2904	.2110		
,10	1 1 2 3 4		0109	6767	5405	.4162	.3108	.2260	.1605	.1117
Bin.	.9862	.9216	.8108	.0707						

$P(X \le 2) = \begin{pmatrix} 10 \\ 0 \end{pmatrix}$	$\binom{0}{p^{0}}(1)$	$(-p)^{100} +$	-(100)	$p^{1}(1-p)$	$)^{99} + \begin{pmatrix} 10 \\ 2 \end{pmatrix}$	$\binom{10}{2}p^{2}(1)$	$(-p)^{98}$			
(0	)	02	03	04	.05	.06	.07	.08	.09	.10
р	.01	.02	.05				0250	0113	.0048	.0019
$P(X \leq 2)$	.9206	.6767	.4198	.2321	.1183	.0566	.0238	.0115		

For values of p quite close to 0, the probability of lot acceptance using this plan is larger than that for the previous plan, whereas for larger p this plan is less likely to result in an "accept the lot" decision (the dividing point between "close to zero" and "larger p" is someplace between .01 and .02). In this sense, the current plan is better.

37.

35.

 $P(\text{accepting the lot}) = P(X_1 = 0 \text{ or } 1) + P(X_1 = 2, X_2 = 0, 1, 2, \text{ or } 3) + P(X_1 = 3, X_2 = 0, 1, \text{ or } 2)$  $= P(X_1 = 0 \text{ or } 1) + P(X_1 = 2)P(X_2 = 0, 1, 2, \text{ or } 3) + P(X_1 = 3)P(X_2 = 0, 1, \text{ or } 2).$ p = .01: = .9106 + (.0756)(.9984) + (.0122)(.9862) = .9981 p = .05: = .2794 + (.2611)(.7604) + (.2199)(.5405) = .5968 p = .10: = .0338 + (.0779)(.2503) + (.1386)(.1117) = .0688

39.

2

$$AOQ = pP(A) = p[(1-p)^{50} + 50p(1-p)^{49} + 1225p^2(1-p)^{\infty}]$$

$$\frac{p}{AOQ} = .01 \quad .02 \quad .03 \quad .04 \quad .05 \quad .06 \quad .07 \quad .08 \quad .09 \quad .10$$

$$\frac{p}{AOQ} = .010 \quad .018 \quad .024 \quad .027 \quad .027 \quad .025 \quad .022 \quad .018 \quad .014 \quad .011$$

**b.** p = .0447, AOQL = .0447P(A) = .0274

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

## Chapter 16: Quality Control Methods

c. ATI = 50P(A) + 2000(1 - P(A))

	01	02	.03	.04	.05	.06	.07	.08	.09	.10
p	.01	.01 .02			1000		1202 (	1550 3	1686.1	1781.6
ATI	77.3	202.1	418.6	679.9	945.1	1188.8	1393.0	1559.5	1000.1	

# Supplementary Exercises

 $n = 6, k = 26, \Sigma \overline{x}_i = 10,980, \overline{\overline{x}} = 422.31, \Sigma s_i = 402, \overline{s} = 15.4615, \Sigma r_i = 1074, \overline{r} = 41.3077$ 41. *S* chart:  $15.4615 \pm \frac{3(15.4615)\sqrt{1-(.952)^2}}{.952} = 15.4615 \pm 14.9141 \approx .55, 30.37$ *R* chart:  $41.31 \pm \frac{3(.848)(41.31)}{2.536} = 41.31 \pm 41.44$ , so LCL = 0, UCL = 82.75  $\overline{X}$  chart based on  $\overline{s}$ :  $422.31 \pm \frac{3(15.4615)}{.952\sqrt{6}} = 402.42,442.20$  $\overline{X}$  chart based on  $\overline{r}$ :  $422.31 \pm \frac{3(41.3077)}{2.536\sqrt{6}} = 402.36,442.26$ 

13	i	$\overline{x}_i$	Si	ri
The state	1	50.83	1.172	2.2
	2	50.10	.854	1.7
	3	50.30	1.136	2.1
	4	50.23	1.097	2.1
	5	50.33	.666	1.3
	6	51.20	.854	1.7
	7	50.17	.416	.8
	8	50.70	.964	1.8
	9	49.93	1.159	2.1
	10	49.97	.473	.9
	11	50.13	.698	.9
	12	49.33	.833	1.6
	13	50.23	.839	1.5
	14	50.33	.404	.8
	15	49.30	.265	.5
	16	49.90	.854	1.7
	17	50.40	.781	1.4
	18	49.37	.902	1.8
	10	49.87	.643	1.2
	20	50.00	.794	1.5
	20	50.80	2.931	5.6
	21	50.43	.971	1.9

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

 $\Sigma_{\overline{s}_i} = 19.706$ ,  $\overline{s} = .8957$ ,  $\Sigma_{\overline{x}_i} = 1103.85$ ,  $\overline{\overline{x}} = 50.175$ ,  $a_3 = .886$ , from which an s chart has LCL = 0 and UCL =  $.8957 + \frac{3(.8957)\sqrt{1 - (.886)^2}}{.886} = 2.3020$ , and  $s_{21} = 2.931 > UCL$ . Since an assignable cause is assumed to have been identified we eliminate the 21<sup>st</sup> group. Then  $\Sigma s_i = 16.775$ ,  $\overline{s} = .7998$ ,  $\overline{\overline{x}} = 50.145$ . The resulting UCL for an s chart is 2.0529, and  $s_i < 2.0529$  for every remaining *i*. The  $\overline{x}$  chart based on  $\overline{s}$ 50 145+ has limits

Its 
$$50.145 \pm \frac{1}{.886\sqrt{3}}$$

$$\frac{3(.7988)}{.886\sqrt{3}} = 48.58,51.71$$
. All  $\overline{x}_i$  values are between these limits.

$$\Sigma n_i = 4(16) + (3)(4) = 76, \ \Sigma n_i \overline{x}_i = 32,729.4, \ \overline{x} = 430.65,$$

$$s^2 = \frac{\Sigma(n_i - 1)s_i^2}{\Sigma(n_i - 1)} = \frac{27,380.16 - 5661.4}{76 - 20} = 590.0279, \ \text{so} \ s = 24.2905. \ \text{For variation: when } n = 3,$$

$$UCL = 24.2905 + \frac{3(24.2905)\sqrt{1 - (.886)^2}}{.886} = 24.29 + 38.14 = 62.43; \ \text{when } n = 4,$$

$$UCL = 24.2905 + \frac{3(24.2905)\sqrt{1 - (.921)^2}}{.921} = 24.29 + 30.82 = 55.11. \ \text{For location: when } n = 3,$$

$$430.65 \pm 47.49 = 383.16,478.14; \ \text{when } n = 4, \ 430.65 \pm 39.56 = 391.09,470.21.$$

45.

www.cengage.com/math



To learn more about Cengage learning, visit www.cengage.com Purchase any of our products at your local college store or at our preferred Telegram: @unohime store www.cengagebrain.com

